

# Observed reductions of surface solar radiation at sites in the United States and worldwide from 1961 to 1990

Beate G. Liepert

Lamont-Doherty Earth Observatory of Columbia University, Palisades, New York, USA

Received 13 February 2002; revised 8 April 2002; accepted 11 April 2002; published 24 May 2002.

[1] Surface solar radiation revealed an estimated 7 W/m<sup>2</sup> or 4% decline at sites worldwide from 1961 to 1990. Here I find that the strongest declines occurred in the United States sites with 19 W/m<sup>2</sup> or 10%. The clear sky optical thickness effect accounts for -8 W/m<sup>2</sup> and the cloud optical thickness effect for -18 W/m<sup>2</sup> in three decades. If the observed increases in cloud cover frequencies are added to the clear sky and cloud optical thickness effect, the higher all sky reduction in solar radiation in the United States can be explained. It is shown that solar radiation declined below cloud-free sky because of the reduction of the cloud-free fraction of the sky itself and because of the reduction of clear sky optical thickness. Solar radiation exhibits no significant changes below cloud-covered sky because reduced cloud optical thickness is compensated by increased frequencies of hours with overcast skies. *INDEX TERMS*: 0305 Atmospheric Composition and Structure: Aerosols and particles (0345, 4801); 0320 Atmospheric Composition and Structure: Cloud physics and chemistry; 1655 Global Change: Water cycles (1836); 3359 Meteorology and Atmospheric Dynamics: Radiative processes

## 1. Introduction

[2] Measurements of solar radiation at the surface are important to discussions of climate change and global warming because they can indicate anthropogenic disturbances [Ramanathan *et al.*, 2001]. First, solar radiation is affected by clouds, which in turn are affected by global warming due to greenhouse gases. Second, increasing air pollution especially anthropogenic particulate matter such as sulfate, soot, etc. pollute the troposphere over major parts of the continents and oceans. These aerosol particles scatter sunlight back into space and hence reduce surface heating on a highly variable time and space scale [Haywood *et al.*, 1997]. Some aerosols (e.g. carbonaceous, dust aerosols) also absorb solar energy and therefore heat the atmospheric layer and not the surface itself. Third, aerosol particles can act as cloud condensation nuclei and modify cloud lifetime and cloud transmissivity altering surface solar radiation [Ackerman *et al.*, 1993]. In general, surface solar radiation is a most valuable but rarely used climate parameter. Broadband solar radiation is routinely recorded at many sites worldwide since the International Geophysical Year in 1957–58. From early on calibration and maintenance procedures have been standardized, which make a long-term study feasible even though the accuracy is debated and there is room for further improvement [World Climate Research Programme, 1991].

## 2. All Sky Solar Radiation


[3] Here I use a new quality-tested global database of monthly mean surface solar radiation time series, the Global Energy Balance Archive (GEBA), which is maintained by the World

Radiation Data Center [Gilgen *et al.*, 1998]. For this study 252 records covering the entire 1961 to 1990 time period and additionally 43 time series from the United States National Solar Radiation Database [Maxwell *et al.*, 1995] were selected. For each station I calculated monthly averages for the following three decades: 1961 to 1970, 1971 to 1980, and 1981 to 1990. Figure 1a shows the distribution of surface solar radiation for the 1981 to 1990 baseline period. The average surface solar radiation is 182 W/m<sup>2</sup>. Note that this average does not represent a global mean. The temporal variability of these time series reveal an averaged drop of 4 W/m<sup>2</sup> from the 1970s to the 1980s and 7 W/m<sup>2</sup> from the 1960s to the 1980s (Table 1), which means a reduction of about 4% in three decades. Here the differences of the decadal monthly means for the 1961 to 1970 or the 1971 to 1980, and the 1981 to 1990 baseline period were calculated for each station separately and the anomalies were only considered if all twelve months exist. The anomalies were then added to the 1981–90 baseline mean. The resulting decline is slightly more than what Stanhill and Cohen [2001] calculated for a reduced global data set of yearly averages. Their analysis shows globally averaged differences of 5 W/m<sup>2</sup> or 3% in surface solar radiation between the 1958 annual average and the 1992 average. Gilgen *et al.* [1998] who analyzed only spatial averages of varying temporal length report declining tendencies in many regions as well.

[4] Solar radiation declined most strongly in North America with all but one station showing reductions between the 1960s and the 1980s and 29 out of 34 showing reductions from the 1970s to the 1980s (see Figure 1b). This result is also illustrated in Figure 2a where the decadal mean cycles of the surface solar radiation for the United States sites are plotted. The averaged reduction of 19 W/m<sup>2</sup> from the 1960s to the 1980s with the strongest decline in the second decade is remarkable (Table 1). The surface solar radiation also declined in areas of the Former Soviet Union (FSU) as shown in Figure 2b. But the mean decrease during the 1960s to the 1980s time period is only 2 W/m<sup>2</sup>. See also Abakumova *et al.* [1996] and Russak [1990]. Similarly, declining tendencies have been found around the Mediterranean Sea in Israel, Egypt, and Turkey [Stanhill and Ianetz, 1997; Omran, 2000; Aksoy, 1997]. As indicated in Figure 1b and reported elsewhere [Liepert *et al.*, 1993; De Bruin *et al.*, 1995] the interdecadal trends in surface solar radiation in Western Europe are rather mixed. A critical point is that the stations are often located in vicinities of major cities where local pollution may not represent the desirable background conditions [see for example Jauregui and Luyando, 1999]. On the other hand, even extremely remote sites show reductions in solar radiation [see for example Dutton and Bodhaine, 2001].

## 3. Clear and Overcast Sky Solar Radiation

[5] Aerosol particles scatter and absorb sunlight in the cloud free atmosphere and hence reduce the solar energy reaching the surface. On the other hand, clouds are by far the strongest modifiers of surface solar radiation. Consequently investigations of the possible causes of the observed reductions require separating cloud effects from aerosol effects. Since hourly solar




**Figure 1.** Decadal means and changes of surface solar radiation. (a) 1981 to 1990 mean surface solar radiation. (b) Difference in surface solar radiation from 1961 to 1970 period minus 1981 to 1990 baseline period.

radiation recordings and cloud cover observations from 1961 to 1990 are available for the United States sites these data were utilized for separating solar radiation time series into all sky, clear sky and overcast sky composites for each ten-year period. An hour is defined as clear if the cloud cover is 0 or 1 tenths of sky coverage at the beginning and end of the hour. Overcast is defined at 10 tenths cloud coverage at the beginning and the end of an hour. The monthly mean diurnal cycles are calculated by averaging all means of solar radiation of each hour of the 24-hour day for a month. Composites of diurnal cycles avoid biases due to data sampling like overrepresentation of morning hours.

[6] The 1960s, 1970s, and 1980s monthly means of the clear-sky solar radiation are plotted in Figure 3a for the United States sites. Although clear sky solar radiation decreased by  $8\text{ W/m}^2$  in three decades (Table 1)—indicating an enhanced direct aerosol effect—the reduction is not as high as under all-sky conditions. Clear sky changes are mainly observed during the 1980s and not during the 1960s or 1970s. A rather different behavior is shown in Germany where during the 1970s to the 1980s positive trends have been observed (Table 1). These declines indicate reductions in the aerosol load and improvements of air pollution. The anthropogenic changes in aerosol concentrations and their effect on surface solar radiation have been calculated with a general circulation model by *Teegen et al.* [2000] and analyzed by *Liepert and Teegen* [2002]. For the United States sites of this study the model simulates a reduction of  $1\text{ W/m}^2$  in three decades. For the German sites the modeled increase in surface solar radiation is  $3\text{ W/m}^2$ , which is exactly what is observed.

[7] As shown in Figure 3b, the solar radiation under overcast conditions dropped drastically by  $18\text{ W/m}^2$  or  $14\%$  in three decades at the United States and  $5\text{ W/m}^2$  or  $11\%$  at the German sites (Table 1). The surface solar radiation decreased even more under the optically thicker “opaque” clouds with a decline of  $20\text{ W/m}^2$  or  $21\%$  in three decades in the United States. “Opaque” clouds are defined as clouds that “prevent observing the sky or higher cloud layers” [*Maxwell et al.*, 1995], which



**Figure 2.** Monthly mean surface solar radiation. Blue line represents 1961 to 1970, green 1971 to 1980, and red 1981 to 1990 for (a) United States and (b) Former Soviet Union sites.

excludes cirrus and lower level thin clouds. The declines may be due to increasing sulfate aerosol concentrations that increase cloud optical thickness and lifetime. But over the continental mid-latitudes the indirect aerosol effect is presumably not as dominant. In *Liepert and Lohmann* [2001] we investigate the indirect aerosol effect on surface solar radiation with a general circulation model that includes a coupled sulfur chemistry cycle and cloud scheme. The simulations show reductions of surface solar radiation at overcast skies of about  $4\text{ W/m}^2$  or  $4\%$  for the sites in the United States, which is only one fifth of what is observed.


#### 4. Cloud Cover Changes

[8] In this study surface solar radiation is analyzed for subsets of clear sky and overcast sky conditions. A clear sky effect of  $\Delta F_{cl} = 8\text{ W/m}^2$  and a cloud optical thickness effect of  $\Delta F_o = 18\text{ W/m}^2$  for three decades have been detected. The all sky solar radiation records, however, consist of a clear, and a

**Table 1.** Observed 10-Year Annual Means of Surface Solar Radiation ( $F$ ) For Various Regions and Cloud Cover Conditions

Stations	All	NH	FSU	United States			Germany			
	All sky $F$ ( $\text{W/m}^2$ )	All sky	All sky	All sky $F$ ( $\text{W/m}^2$ )	Clear sky $F_{cl}$ ( $\text{W/m}^2$ )	Overcast $F_o$ ( $\text{W/m}^2$ )	Opaque $F_q$ ( $\text{W/m}^2$ )	All sky $F$ ( $\text{W/m}^2$ )	Clear sky $F_{cl}$ ( $\text{W/m}^2$ )	Overcast $F_o$ ( $\text{W/m}^2$ )
1961–70	$189 \pm 6$	$163 \pm 3$	$125 \pm 4$	$200 \pm 4$	$243 \pm 3$	$128 \pm 10$	$95 \pm 8$	$124 \pm 5$	$190 \pm 4$	$47 \pm 4$
1971–80	$186 \pm 6$	$160 \pm 4$	$123 \pm 4$	$194 \pm 5$	$242 \pm 3$	$121 \pm 11$	$84 \pm 9$	$122 \pm 4$	$190 \pm 4$	$44 \pm 3$
1981–90	$182 \pm 6$	$156 \pm 4$	$123 \pm 4$	$181 \pm 7$	$235 \pm 6$	$110 \pm 14$	$75 \pm 11$	$120 \pm 4$	$193 \pm 3$	$42 \pm 3$

(The uncertainty ranges are based on the statistical mean error of the decadal average monthly and hourly means for each site.) All stations, NH northern hemisphere records, FSU Former Soviet Union records, United States and Germany.



**Figure 3.** Monthly mean surface solar radiation in United States. Blue line represents 1961 to 1970, green 1971 to 1980, and red 1981 to 1990 period for (a) clear sky and (b) overcast sky conditions.

overcast portion as defined in chapter 3 and a cloudy portion. The observed all sky reduction is  $\Delta F = 19 \text{ W/m}^2$ . Here I use the calculated averages of clear sky  $F_{cl}$  and overcast sky  $F_o$  solar radiation to estimate the cloudy sky solar radiation. Solar radiation  $F$  is separated into a portion below cloud-free sky and below cloud-covered sky. The linear partitioning is described in equation (1). Clear sky solar radiation decadal means  $F_{cl}$  are multiplied with the clear sky frequency  $f_{cl}$  plus  $f_{cd}(1 - A_c^*)$  to estimate the below cloud-free sky solar radiation  $F_{cf}$ . The below cloud-covered sky solar radiation  $F_{cc}$  is calculated with the overcast sky solar radiation decadal means  $F_o$  multiplied by overcast sky frequencies  $f_o$  plus  $f_{cd} A_c^*$ .  $f_{cd}$  is the cloudy sky frequency and  $A_c^*$  is the cloudy sky areal coverage as defined in equation (1).  $A_c^*$  is larger than the total cloud coverage  $A_c$  which does include the two extremes clear sky and overcast sky (Table 2). The cloud-covered portion of solar radiation  $F_{cc}$  is a lower limit because cloud optical thickness tends to increase

with increasing cloud coverage and  $F_{cc}$  is calculated with the solar radiation at overcast cloud coverage.

$$F = F_{cl} \cdot f_{cl} + F_o \cdot f_o + F_o \cdot f_{cd} \cdot A_c^* + F_{cl} \cdot f_{cd} \cdot (1 - A_c^*)$$

$$\text{with } A_c^* = \sum_{c_i} f_{c_i} \cdot c_i, \quad \text{with } c_i = i/10; i = 2, \dots, 9$$

$$f_{cd} = 1 - f_{cl} - f_o \quad (1)$$

[9] With equation (1) all sky changes of solar radiation can be estimated under different conditions. If exclusively the clear sky and overcast sky solar radiation changed over time the combined effect would be  $\Delta F = F_{1960s} - F_{1980s} = 13 \text{ W/m}^2$ . The modeled aerosol effect (direct plus first indirect) would account for only  $\Delta F = 2.6 \text{ W/m}^2$ . These results are not enough to explain the observed  $19 \text{ W/m}^2$  all-sky decline of this study. Note that the observed total cloud coverage  $A_c$  did not change at these United States sites during the three decades (Table 2). However, the frequencies of the occurrence of hours with overcast and clear skies  $f_{cl}$  and  $f_o$  changed.  $f_o$  increased from 23% to 31% and  $f_{cl}$  declined from 32% to 26% in thirty years. Consequently the combined effect  $\Delta F$  reaches a  $20 \text{ W/m}^2$  reduction in surface solar radiation in three decades in the United States when these cloud frequency changes are included. This result matches the independently calculated all-sky reduction of  $\Delta F = 19 \text{ W/m}^2$ . Noteworthy is that the below cloud-covered sky solar radiation does not change over time anymore because increasing cloud optical thickness and increasing cloud frequency effect cancel each other. Rather the below cloud-free sky solar radiation exhibits a strong decline due to accumulation of the increasing clear sky optical thickness and the declining clear sky frequency.

## 5. Conclusions

[10] The observed  $19 \text{ W/m}^2$  or 10% decline in surface solar radiation in the United States from the 1960s to the 1980s is strong compared to sites in other regions of the globe with a  $7 \text{ W/m}^2$  or 4% decline in three decades. Nonetheless, the uncertainty range of the data cannot be ignored. Increasing direct and indirect aerosol effects through increasing air pollution alone can explain about  $3 \text{ W/m}^2$  of the observed decline in the United States when simulated with state-of-the-art general circulation models. Increasing frequencies of overcast skies and declining frequencies of clear skies with simultaneously constant cloud coverage indicate more fractional or multi-layered cloud coverage. This is in line with the strongest reduction of surface solar radiation under optically thick “opaque” clouds and also the observed slight reduction of  $A_c^*$ . Similar results have been shown in a study of cloud type, cloud coverage and solar radiation changes for German sites [Liepert, 1997]. One possible explanation might be a combination of greenhouse gas and aerosol forcing. Tselioudis and Rossow [1994] suggest that greenhouse gas warming leads to increasing cloud optical thickness at overcast conditions (through increasing cloud height and liquid water content). Increasing optical thickness reduces surface solar radiation and cools the surface. Albeit, in

**Table 2.** Observed 10-year Annual Means of Cloud Cover Properties and Surface Solar Radiation for United States Sites

	$f_{cl}$	$f_o$	$A_c^*$	$A_c$	Cloud-free $F_{cf}$ ( $\text{W/m}^2$ ) ( $F_{cl} f_{cl} + F_o f_{cd}(1-A_c^*)$ )	Cloud-covered $F_{cc}$ ( $\text{W/m}^2$ ) ( $F_o f_o + F_o f_{cd} A_c^*$ )	Combined F ( $\text{W/m}^2$ )
1961–70	$0.32 \pm 0.13$	$0.23 \pm 0.11$	0.59	0.56	124 (80 + 44)	61 (28 + 33)	185
1971–80	$0.29 \pm 0.13$	$0.27 \pm 0.11$	0.58	0.56	116 (70 + 46)	62 (31 + 31)	178
1981–90	$0.26 \pm 0.13$	$0.31 \pm 0.11$	0.58	0.56	105 (63 + 42)	60 (33 + 27)	165

Frequencies of clear  $f_{cl}$ , overcast  $f_o$  sky, total cloud coverage  $A_c$ , and cloudy sky areal coverage  $A_c^*$ , cloud-free sky portion  $F_{cf}$  and cloud covered sky portion  $F_{cc}$  and combined F surface solar radiation.

general the troposphere is still heated through greenhouse gas warming and water vapor is still augmented. Furthermore, absorbing aerosols that heat the atmospheric layer may enhance this warming. Consequently tropospheric warming and surface cooling might occur (see also *Roeckner et al.*, [1999]). It is worth noting that the reductions in surface solar radiation are simultaneous with an observed missing increase in surface air temperature in the United States from 1961 to 1990 [*Hansen et al.*, 2001]. More model simulations are needed and strongly encouraged to explain the reductions of surface solar radiation.

[11] **Acknowledgments.** I thank the anonymous reviewers for their insightful comments. The data were provided by NOAA and WRDC who maintain these valuable data bases. This work was sponsored by NASA Global Aerosol Climatology Project NAG 5-7687.

## References

- Abakumova, G. M., E. M. Feigelson, V. Russak, and V. V. Stadnik, Evaluation of long-term changes in radiation, cloudiness, and surface temperature on the territory of the former Soviet Union, *J. Climate*, 9, 1319–1327, 1996.
- Ackerman, A., O. B. Toon, and P. V. Hobbs, Dissipation of marine stratiform clouds and collapse of the marine boundary layer due to the depletion of cloud condensation nuclei by clouds, *Science*, 262, 226–228, 1993.
- Aksoy, B., Variations and trends in global solar radiation for Turkey, *Theor. Appl. Climatol.*, 58, 71–77, 1997.
- De Bruin, H. A. R., B. J. J. M. Van den Hurk, and D. Welgraven, A series of global radiation at Wageningen for 1928–1992, *Int. J. Climatol.*, 15, 1253–1272, 1995.
- Dutton, E., and B. Bodhaine, Solar irradiance anomalies caused by clear-sky transmission variations above Mauna Loa: 1958–99, *J. Climate*, 14, 3255–3262, 2001.
- Gilgen, H., M. Wild, and A. Ohmura, Means and trends of shortwave irradiance at the surface estimated from global energy balance archive data, *J. Climate*, 11, 2042–2061, 1998.
- Hansen, J. E., et al., A closer look at United States and global surface temperature change, *J. Geophys. Res.*, 106(D20), 23,947–23,963, 2001.
- Haywood, J. M., D. L. Roberts, A. Slingo, J. M. Edwards, and K. P. Shine, General circulation model calculations of the direct radiative forcing by anthropogenic sulfate and fossil-fuel soot aerosol, *J. Climate*, 10, 1562, 1997.
- Jauregui, E., and E. Luyando, Global radiation attenuation by air pollution and its effects on the thermal climate in Mexico City, *Int. J. Climatol.*, 19, 683–693, 1999.
- Liepert, B. G., Recent Changes in Solar Radiation under Cloudy Conditions in Germany, *Int. J. Climatol.*, 17, 1581–1593, 1997.
- Liepert, B. G., and G. J. Kukla, Declines in global solar radiation with increased horizontal visibility in Germany between 1964 and 1990, *J. Climate*, 10, 2391–2401, 1997.
- Liepert, B. G., and U. Lohmann, A comparison of surface observations and ECHAM4-GCM experiments and its relevance to the indirect aerosol effect, *J. Climate*, 14, 1078–1091, 2001.
- Liepert, B. G., and I. Tegen, Multi-decadal solar radiation trends in the United States and Germany and direct tropospheric aerosol forcing, *J. Geophys. Res.*, in print, 2002.
- Maxwell, E. L., W. F. Marion, D. R. Myers, M. D. Rymes, and S. M. Wilcox, *Final Technical Report Volume 1 + 2: National Solar Radiation Database (1961–1990)*, National Renewable Energy Laboratory, Golden, Colorado, 1995.
- Omran, M. A., Analysis of solar radiation over Egypt, *Theor. Appl. Climatol.*, 67, 225–240, 2000.
- Ramanathan, V., P. J. Crutzen, J. T. Kiehl, and D. Rosenfeld, Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119–2124, 2001.
- Roeckner, E., L. Bengtsson, J. Feichter, J. Leveiveld, and H. Rodhe, Transient climate change simulations with a coupled atmosphere ocean GCM including the tropospheric sulfur cycle, *J. Climate*, 12, 3004–3032, 1999.
- Russak, V., Trends of solar radiation, cloudiness and atmospheric transparency during recent decades in Estonia, *Tellus*, 42B, 206–210, 1990.
- Stanhill, G., and A. Ianetz, Long-term trends in and the spatial variation of global irradiance in Israel, *Tellus*, 49B, 112–122, 1997.
- Stanhill, G., and S. Cohen, Global dimming, a review of the evidence for a widespread and significant reduction in global radiation with a discussion of its probable causes and possible agricultural consequences, *Agric. For. Meteorol.*, 107, 255–278, 2001.
- Tegen, I., D. Koch, A. A. Lacis, and M. Sato, Trends in tropospheric aerosol loads and corresponding impact on direct radiative forcing between 1950 and 1990: A model study, *J. Geophys. Res.*, 105(D22), 26,971–26,989, 2000.
- Tselioudis, G., and W. B. Rossow, Global, multiyear variations of optical thickness with temperature in low and cirrus clouds, *Geophys. Res. Lett.*, 21, 2211–2214, 1994.
- World Climate Research Programme (WCRP), Baseline Surface Radiation Network (BSRN). WCRP-54, WMO/TD-No. 406, WMO, 1991.

B. G. Liepert, presently at Max-Planck Institute for Meteorology, Bundesstrasse 55, Hamburg D-20146, Germany.