

	Hudson Riv	er Estuary	
Educ	eator Packet for A Day in the Life of	the Hudson Riv	ver (Snapshot Day)
	•	Year	(1
	http://www.ldeo.columbia.ed	u/edu/k12/sna	 nshotday/
The Packe	t is designed for educators & teachers wit		
	hat are a part of A Day in the Life of the H		
	an be completed as part of the day's event		
	Life website. Student data recording shee		
	sure to submit your results to Margie Turi		
mkt@ldeo	.columbia.edu) within 24-48 hours of colle	ection! Questions?	845-365-8494.
	BE SURE TO RECORD TIME & UNIT		
ITE	M SO THAT COMPARISONS CAN BE		HOUT THE RIVER
	ACTIVITY	PAGE	
	1. Site Background Information	Page 1	
	2. Tides & Currents 3. Weather & Wind	Page 2 & 3	
	4. Beaufort Chart (wind)	Page 4 Page 5	
	4. The Sampling Site Environment	Page 6-7	
	5. Site Sketch	Page 7	
	6. Other Physical Factors	Page 8	
	7. Chlorophyll Sampling	Page 9	
	8. Sediment Sampling	Page 10 -12	
	9. Chemical Measures	Page 13-14	
	10. Fish & Macroinvertebrates	Page 15-16	
	11. Hudson River Fish Checklist	Page 17-18	
	12. Other Observations - Shipping e	_	
G: D 1	13. Science Journaling/Almanac En	try Page 19	
Site Back	ground Information.		
1. Coordin	ator/contact person		
Organiz	ation		
Street			
	StateZip_		
phone	faxemail		
	group name		ict
	f teacher/group leader		
	C 1		_
City	StateZip		
	fay er		

Number of student participants grade level/age Number of Adults.

3. Please tell us where you are sampling. Be as specific as possible. (Example: swimming beach, Kingston Point, City of Kingston, Ulster County.)

	nattan is River Mile 0; the Federal Dam at Troy is River
Mile 153.)	lattan is River wine 0, the rederal Dam at 110y is River
River mile	
If you have a way to determine the latitude an	d longitude of your site, enter that data here.
GPS Latitude	Longitude
Activity I - Tides and currents	

TIDES: The tide is the up and down motion of the water, the rising & falling. There are several ways to measure tide.

Simple Measure: At a beach use two slender, strong sticks as tide markers. At the start, place one stick at the water's edge. Push it deep into the ground or pile rocks at its base to hold it in place. Every 15 or 30 minutes check your marker If the level has changed, place the second stick to mark the new position of the water's edge on a beach so through time you see the total change. Record on your sheet the water level as rising, falling, or unchanged. *Pier or bulkhead* choose a distinct, immoveable feature on or near the bulkhead to see whether the water level is rising or falling. If there are waves, use your judgment in deciding where the water's edge/surface is. Record on your sheet the water level as rising, falling, or unchanged.

Intermediate Measure: Use a dowel marked in 10cm increments to set firmly in the sediment in the water OR if measuring off a bulkhead use a tape measure to measure from the dock to the water surface. Have the students record the water level once the marker is set (record in column 3). Then every 15 to 30 minutes check your marker and record actual measures so that a total tidal change can be calculated over a total time available for the activity. NOTE: measuring down to the water means the distance will get smaller as the tide rises – be sure you talk through this with students.

Extra Activity: If your students have time at this station they can calculate how quickly the tide is rising or falling by dividing the change in height by the time between recordings. Think of the basic definition of speed as distance traveled divided by the time of travel. Calculation: Subtract the time from prior reading from the time of this reading for 'time of travel' (or time elapsed). Next calculate the change in height from your prior reading (for the distance traveled) by subtracting these two numbers. Now divide the change in height by the time elapsed. This is the rate (speed) of tidal change (column 4)

	TIDES							
Time	Rising, Falling, Unchanged	Height in cm (if recording)	Rate of Tidal Change (cm/min)					

CURRENTS: Currents are the internal movement in the water sometimes described as a push and pull in the water. After recording the tide level, determine the direction of the current. Using a mid sized stick (large enough so the wind can't easily push it), or an orange, toss it as far as you can out into the river. Note which direction it moves. The current moving downriver towards the sea is called the ebb; the current moving upriver is the flood. Don't confuse the direction of waves with the direction of the current; waves and current are different things.

Basic Measure: Every 60 minutes toss an orange (or stick) as far out into the water as you can. Record the direction of travel as North (towards Albany) or South (towards the Atlantic) in column 5. Next record incoming as Ebb, outgoing as Flood, or if there is no movement record it as Slack in column 6. Intermediate Measure: For a more exact measure you can calculate the speed (distance traveled divided by time traveled) by using a tape and a stopwatch to measure how fast the current is moving. Toss the stick or orange into the water at a marked starting point (use a student to align with the start). Stop your watch after 60 seconds on a stop watch and place a student at the end point. Now have your students measure the distance between the two student markers with a measuring tape. Record this in column 2. Calculate distance per second by dividing the total distance by 60 secs. Record in column 3. Extra Activity: Calculate KNOTS: If your students want to calculate the rate of travel in knots use the distance in cm for 60 seconds to compute this. Let's think this through.

1 kt. = 6076 ft. per hr. But we have cm so we need to convert ft. to cm. 1 ft. = 30.48 cm. so multiple these two to compute cm/hr or 185196.5 cm/hr. Now divide by 60 for cm per minute (3086.6 cm/min.) now by 60 again for cm/sec. What you find is that 1 kt = 51.44 cm/sec.

SO to compute Knots from cm/sec use the following equation:

kts = cm/sec divided by 51.4. Record this as knots in column 4.

Example: If the stick traveled 125 cms in 60 seconds divide 125/60 = 2.08 cm sec. /51.4 = .04 kts.

	CURRENT						
Time	Cm/60 sec	Cm/sec	Knots (cm/sec)/51.4	North/ South	Ebb/Flood/Still (E/F/S)		

Is there anything about the river or shoreline here that may cause the current near shore to flow in a different direction than the current out in the middle of the Hudson (a protected embayment, a pier jutting out causing an unusual swirling)?

DATA FROM THE MAIN CHANNEL IS THE BEST DATA TO RECORD SO PLEASE BE ALERT TO DIFFERENCES THAT MIGHT EXIST. IF YOU NOTICE THAT THE CURRENT APPEARS TO BE DIFFERENT IN THE MAIN CHANNEL THAN IT IS IN THE SHORELINE AREA PLEASE RECORD THIS NOTATION USING "S" FOR SHORELINE AND "C" FOR CHANNEL.

Activity II – Weather and Wind

Weather and wind are important pieces of physical data that help to provide context for the other data. Weather includes current conditions and conditions over the last few days that may have an impact on the data you collect today (such as rain, extremely hot or cold weather).

Wind levels can increase choppiness in the water thus adding oxygen and increasing levels of oxygen saturation. Wind can also affect movement on the top of the water surface which may make assessing currents difficult.

1. Record weather condition	ions at the start of	sampling. Record	changes every hour if	f possible.
a. Time	Air temperature	°F	° C	
bTime				
Cloud cover (check one) Any precipitation?			mostly cloudy	overcast
If the weather changes ov	er the time you ar	re sampling, please	note that here.	
Briefly describe the weath	ner for the last thr	ee days. Any rain,	wind, or unusual temp	eratures?
2. Wind speed: (PLEASE REFER TO I	BEAUFORT CH	HART ON PAGE	S 4-5)	
Using the Beaufort chart	record the FIRST	COLUMN as Beau	ıfort FORCE	<u></u> .
Optional additional inform	nation to record _	kts. and/or _	mph	
Using an anemometer to i	record wind recor	d (Be sure to	record as kts ,or mph	(kts preferred)
Record wind direction as until it hits your face ever		_ ,		_
Water Choppy	Water Ca	lm		

Beaufort Wind Scale 09/26/2006 10:56 PM

STORMFAX® WEATHER ALMANAC

Beaufort Wind Scale

Devised by British Rear-Admiral, Sir Francis Beaufort in 1805 based on observations of the effects of the wind

Beaufort	Wind	Speed	Wave	WMO*	ons of the effects of the wind	Effects
number (force)	knots	mph	height (feet)	description	Effects observed on the sea	observed on land
0	under 1	under 1	-	Calm	Sea is like a mirror	7
1	1 - 3	1 - 3	0.25	Light air	Ripples with appearance of scales; no foam crests	7
2	4 - 6	4 - 7	0.5 - 1	Light breeze	Small wavelets; crests of glassy appearance, not breaking	
3	7 - 10	8 - 12	2 - 3	Gentle breeze	Large wavelets; crests begin to break; scattered whitecaps	
4	11-16	13- 18	3½ - 5	Moderate breeze	Small waves, becoming longer; numerous whitecaps	
5	17-21	19- 24	6 - 8	Fresh breeze	Moderate waves, taking longer form; many whitecaps; some spray	
6	22-27	25- 31	9½-13	Strong breeze	Larger waves forming; whitecaps everywhere; more spray	
7	28-33	32- 38	13½- 19	Near gale	Sea heaps up; white foam from breaking waves begins to be blown in streaks	

http://www.stormfax.com/beaufort.htm

Activity III - The Environment at the Sampling Site

Surrounding Land Use:

Estimated % Beach

Estimated % urban/residential_____ Estimated % Forested _____

Estimated % Beach______Estimated % Industrial/Commercial_____

In completing the physical survey of your sample site please include a 200 ft. segment extending up and downriver from your sampling site (your site bisects the segment). Sketch a map of your sampling site on the next page and answer the following questions. If available, use a digital camera to photograph your site.

1. Using the list below describe the land at your site. Is it forested? Open and grassy? A parking lot? Used by people for picnics, launching boats, fishing, swimming, or other activities?

Estimated % Other	(specify)								
bulkheading - woo	den timbers or metal plate long the shore? Do any pi	arsh? Is it sandy, muddy, or rocky s that hold the shore in place? Has pes discharge into the river here?							
SHORELINE AP	SHORELINE APPEARANCE: CHECK ALL THAT APPLY								
Beach area	Pier	Shore with Bulkhead (wood timbers/metal plates)	Shoreline RipRap (large rocks)						
Covered with	Debris in the Area	Piping entering the river -	Brick Pieces						
vegetation	such as broken	(size)							
vegetation	concrete, docking	(North or South or sampling_	Charcoal						
	ocarrott, document	site & estimate distance)	Slag						
the water. Wate	r chestnut REMOVES ox it is photosynthesizing. Us	nents and nutrients, and can assist vygen from under its beds while was se the <i>Hudson River Field Guide to</i> find growing in the water. List the	ter celery adds oxygen to Plants of Freshwater						
Are there plants growing in or on the water? Do they cover more than half of the area you are sampling? Less than half?									
What percent of yo	our entire sampling area	is covered with plants in the water	?						
Check if present a	and list estimated percent	age of the total plant population f	or each plant:						
Tidal Shallows:									
Water Milfoils	% vegetation	Water Celery	% vegetation						

Water Chestnut% vegetation	Other % vegetation
Tidal Marshes:	
Arrow Arum % vegetation	Purple Loosestrife % vegetation
Arrowhead% vegetation	Reed Grass % vegetation
Big Cordgrass % vegetation	Saltwater Cordgrass % vegetation
Broad leaved Cattail % vegetation	Spatterdock% vegetation
Bur-reed% vegetation	Swamp Rose-Mallow % vegetation
Golden Club % vegetation	Sweet Flag % vegetation
Jewelweed % vegetation	Wildrice % vegetation
Mud Wort% vegetation	Yellow Flag % vegetation
Narrow-leaved Cattail % vegetation	Yellow Pond Lily% vegetation
Pickerelweed% vegetation	OTHER % Vegetation
Phragmites /Common Reed% vegetation	

Activity IV - Sketch Map of the Sampling Site

Include a compass rose (N, S, E, W) and rough scale. Label landmarks or notable features. Indicate specific locations where you sampled.

Activity V - Other Physical Factors

1. Water temperature

Water temperature is important for understand the amount of dissolved oxygen the water can hold, and for the fish communities that will use the area. Students will better understand Fahrenheit temperatures, but in science it is important to become familiar with Celsius, so if possible, record water temperature in BOTH degrees Celsius and degrees Fahrenheit. Have them look at the comparison between the two. If you don't have both °C and °F thermometers, students can convert between the two using the following formulas:

$$^{\circ}$$
C = 0.556 x ($^{\circ}$ F - 32) $^{\circ}$ F = (1.8 x $^{\circ}$ C) + 32

It is best to take the water temperature several times in succession and then average. Over the day, you might want to see if there's any change, especially in shallow water and backwater areas, which may show more variation through the day due to sunlight, tide or current changes.

Time	Reading 1	Reading 2	Reading 3	Average
	°F			
	°F °C			
	°F			
	°F			

2. Turbidity

Turbidity is water clarity, an important feature of an estuary. In the Hudson River turbidity is made up of small bits of plankton, pieces of detritus or decomposing plant and animal matter, salt and suspended bits of sediment. Different techniques for determining turbidity use different units of measurement. Be sure to enter data on the correct line for the technique you use. Repeat several times in succession and average the results.

	Time	Reading 1	Reading 2	Reading 3	Average	
secchi disk						feet or cm
short site tube	e					JTUs
long sight tu	be					cm/meters
turbidimeter						NTUs

Snapshot Day Activity IV - Chlorophyll Sampling

Chlorophyll

The pigment Chlorophyll is what allows plants (and algae) to convert water and carbon dioxide to organic compounds in the presence of light, a process called photosynthesis. There are several types of Chlorophyll to assist plants to capture light at different wavelengths, but all plant cells have chlorophyll "a". This activity will allow us to measure the amount of Chlorophyll "a" in your area of the river.

PLEASE FOLLOW THE PROTOTOL SHEET ON THE NEXT PAGE

120 mls (or ccs) of river water will be collected and filtered through a fine mesh filter to separate any material in the water. Once the water is filtered, examine the filter to evaluate the amount of material that is filtered. This will be an accumulation of not just chlorophyll but any suspended matter that was large enough to be filtered out. Using the color chart included with your kit select the color that best matches your filter and record on this data sheet.

At some sites the filter paper will be removed using tweezers – folded to protect the sample and placed in a vial and put on ice for collection and analysis at Lamont.

If the sample is being collected for Lamont label your vial with the following protocol:

Date – River Site –River Mile_ cc volume Example

100809_PP_25_120cc

Record here -		
TIME	# ON THE COLOR CHART MOST CLOSELY MATCHING SAMPLE	_

Activity IIV - Sediment Sampling

 $\hbox{*Use the Step-By-Step directions sheet provided online on the resources page $$http://www.ldeo.columbia.edu/edu/k12/snapshotday/Resources.html}$

Background Information:

- The sediments in the core represent a period of time. The material at the bottom is older than the material on the top. This is an important principle of geology and much of Earth Science called 'superposition'.
- If material has been accumulating steadily, a sediment core will contain a record of the material transported by the river through time.
- One of the challenges faced by scientists who study sediment cores is determining the length of time represented by the sediments core. You can not tell how many years your core represents by simply looking at it. The amount of time represented by your core will vary depending on the specific place and processes of the river in each area. In sections with high deposition it could represent a very short amount of time (days to a year), while in other areas it could represent a much longer time (10s to 100s of years or longer).
- What you can tell from looking at a core is whether the color changes over the length of the core. You will note the color of the sediments at the very top of the core. If the color is light brown, this is an indication that the surface sediments are oxidized (in contact with oxygen in the water). The oxidized section is the top represents an unconsolidated recent deposition. You will measure this and record it to determine how active the deposition is in your area. Usually, the sediments change to a darker color below the oxidized layer, this is called anoxic (no oxygen) or reducing. It usually means that these sediments have been out of contact with the oxygen in the river water and are older. Often this section will have a sulfur smell noting bacterial decomposition. Extrude your core, then measure and record each section. Complete the core assessment sheet as you observe and describe it. Note anything else that you think is significant. Are there other visible layers? Color changes?
- X-Ray Fluorescence (XRF) Spectrometer—What does this mean? Once you collect your core and describe it you will be sending one to Lamont for X-Ray Fluorescence. This is done with a piece of equipment that can measure lead and other metal concentrations in the sediments. We focus on lead since it can be used as an indicator of time. There is a natural background reading of lead in the river (approx. 20ppm) but human (anthropogenic) influences such as early 20th century industry, leaded gas etc. have caused in increase in that level. Using the XRF we can look at what the readings of various metals are in different areas of the river. This information will be generated at Lamont and the results provided back to the group. The hope is to use this information to roughly constrain (locate the probably range) the age of sediments you collect. A straightforward interpretation of this data is that low levels of lead similar to natural background would indicate sediments that are older than (deposited prior to) approximately 1900, while sediments containing lead a levels elevated above the natural background would indicate sediments that are younger or deposited as part of industrialization.
- Collect a sample for back in the classroom. Collect and bag a sample to take back and analyze in the classroom to look at the history of the sediment in the river. See classroom activity sheet called "The sediments in our river".

DAY in the LIFE PUSH CORE SEDIMENT LOG

GRAB ID#	Site I	Name	DATE		FORM COMPLETED BY:		
					GROUP #		
TIME	LATITUDE		LONGITUDE		WATER DEPTH	LOCATION	
	Yes	No			Descriptors - Please note	additional observations	
H ₂ S smell					H₂S smells of rotten eggs,	suggesting anaerobic bacteria	
Oil							
Oxidized top*					*oxidation (reaction with c lighter colored layer of se	oxygen) creates a distinctly diment.	
					estimate dimensions of ox	cided layer, etc. and draw below	
	Absent	Rare	Common	Abundant	Additional Comments		
Clay					dense feel		
Mud					smooth feel between finge	ers	
Sand					gritty feeling between fing	ers	
Gravel					pea sized pieces of stone		
Pebbles					pieces of stone larger than pea		
Leaves							
Wood							
Shells Oysters (dead/alive?)							
Freshwater mussels (except zebra)							
Zebra mussels							
macroinvertebrates							
Brick							
Coal							
Slag							
Living vegetation:							
Length of Core:					Length of Oxidized core to	op (if present):	
If Bagged - Number On	Core Collec	tion Bag					
				,			
Sketch of your core with	n measurem	nents:					

11 form updated 8/10/12

SEDIMENT SAMPLING – push cores were distributed to a group of our participating stations for this process. If you don't have a corer you can skip this activity. Prior to sampling (i.e., before the bag gets wet), please use a permanent marker and label bag with the following information:

- Date River Site –River Mile
- Example:
- 100809 PP 25 (for Piermont Pier)
- You will be taking a core to examine and describe with your group using the form on the next page. Once the description is complete scoop the pieces into a Ziploc bag and return it to Lamont-Doherty Earth Observatory for X-Ray Fluorescence analysis (this will be picked up with your chlorophyll sample). The core will be homogenized for sampling so do not worry about squishing the sample.

DISCUSSION: Once you have the core for group analysis use the sheet that is in your protocols to look through and analyze it with the group. Discuss how any unusual items might have ended up in the river and the role they play there.

(**Note** if the area you are coring is primarily sand the corer may not work and the sand may fall out when you lift the corer from the water. In this case if you would still like to examine the bottom of the river with your students you might be able to slide a flat piece of something under the base of the corer and still extract a sample to look at. If that isn't possible, consider using a jar to scoop out a section trying to obtain a sample that goes down 3-4 inches. The same activities can be completed with this type of sample. When you bag your sample please note it was not obtained with the corer.)

Snapshot Day Recording Sheet VIII - Chemical Measurements

1. pH A measure o	of the acidity of				tral Range is 7 and average the results.
Time	Reading 1	Reading 2	Reading 3	Average	
How was it	determined? (ch	neck one)			
litmus paper	indicator	solution	meter	pH pen	other
different instit is essentia freshwater p milligrams p One ppm is one second one minute one cent in In saltier par (ppt); one pathe estuary i	truments measured that you mark parts of the river per liter (mg/l), which is the like: in 11.5 days, in two years, \$10,000 arts of the river, yeart per thousand	re salinity as 'down what ins t, the units of m which are equi- you may also s equals 1000 m 30 mg/l (.020	total salts', 'cl strument you u neasurement n valent. ee measuremeng/l. Backgrou 030 ppt). In	nloride' or ever used to measure nay be parts per ents expressed i and level of Cl	salinity, however a conductivity. Therefore the salinity. In million (ppm) or n parts per thousand in the freshwater part of f the open Atlantic Ocean
Repeat sever	ral times in succ	cession and ave	erage the resul	ts. Specify the	units of measurement.
Time	Reading 1	Reading 2	Reading 3	Average	Units
How was it	determined? (ch	neck one)			
drop count to	est kit me	eter refra	ctometer	test strips	hydrometer
` _	conductivity pleas as and then conv		11 1	ınit uS/cm (mic	erosiemens) or mS/cm
2 Diagolaro	d				

3. Dissolved oxygen.


The amount of dissolved oxygen (DO) in a river is one of the most important factors determining its health. Many variables influence DO, including temperature, time of day, presence of plants, and wind conditions. DO measurements are given in mg/l and as percent saturation. At 100% saturation, water of a given temperature cannot hold more DO. If more is added - by wind or turbulence, saturation may temporarily exceed 100%, but in this case oxygen will diffuse from the water into the air. Saturation levels below 100% are not necessarily the result of pollution. At night, when plants aren't producing oxygen through photosynthesis, saturation may fall below 100% as living things use up the available DO.


time	temperature	e in °C	DO (mg/l)	% satu	ration		
		_					
How was it	determined? (check one)	1				
Drop co	unt test kits	amnules	digital	titrator	meter	other	

DETERMINING PERCENT SATURATION THE "QUICK AND EASY" METHOD

Source of chart: http://waterontheweb.org/under/waterquality/oxygen.html

For a quick and easy determination of the percent saturation value for dissolved oxygen at a given temperature, use the saturation chart above. Pair up the mg/l of dissolved oxygen you measured and the temperature of the water in degrees C. Draw a straight line between the water temperature and the mg/l of dissolved oxygen. The percent saturation is the value where the line intercepts the saturation scale. Waterways with a saturation value of 90% or above are considered healthy.





ADDITIONAL CHEMICAL TESTS IF DESIRED

The following tests typically require more complex methods than those described above. There is no requirement to do these, but if you have the equipment and ability, the data would be welcome.

4. Nitrates.

Nitrate (NO₃⁻) is relatively plentiful in freshwater ecosystems but less so in saltwater ecosystems, where it is typically the limiting nutrient.

Repeat several times in succession and average the results.

Time	Reading 1	Reading 2	Reading 3	Average	
					NO_3 (mg/l)

Repeat seve	ral times in suc	cession and ave	erage the results	3.
Time	Reading 1	Reading 2	Reading 3	Average
				$PO_4^{-3} \text{ mg/l}$
acid precipit water is; the water hardneshifts in pH system caus Alkalinity re	a a measure of value	onfuse it with place determines the ter small amound the addition of that absorbs or significant in mg/l of calc	H. pH measure concentration of acid or a f small particles soaks up small of ium carbonate (
Time				
Time	Reading 1	Reading 2	Reading 3	Average
				CaCO ₃ mg/l
The data secessily visible plankton. If together. If you and America together as a possible to the instance - it so that we considered to the please note please see different solutions.	e without magn making repeate you have troubled of classification an shad - look valuering or sunfi- ell males from is possible and the can compare day PULL and TO catches per traj irections on ba	et up for fish ar iffication. This ed collections, re identifying or on possible. You wery similar to osh. Measure the females for movery useful to out a from site to TAL NUMBER p. If your grouck.	and invertebrates sheet can be addrecord data for examining to the regarding of the year one another, as a largest individual st of what you distinguish generate please list of the caugh would like to	such as crabs and crayfish that are apted if you plan to capture and study each haul and then add the catch totals species level, list them at the most r herring - alewife, blue-back herring, do very young sunfish. Group them dual of each species. It will not be catch, but for a few - blue crabs for
				study period
	ipment used:	emenes your m		y periou
	-	esh size)		
eel pot	minnow trap_	dip net	plankton	net other:

FISH SPECIES CAUGHT

Use separate sheet	for each seine OR note	what was caught in	each seine by TIME	noting sein	ne #.
Fish Species:		# of individuals:		of largest	(unit)
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
11					
12					
Macroinvertebrates: 1 2 3 4 5					
Take length of net (5 meters - 50 ft. X 21f inches in a meter = 3 This figure should be If you pull the net in / 39.37 (inches in a m	Per Unit Equivalent (6 0 ft.) X length of pull (6 t X 12 (for inches per fo 20 meters. Then divide e computed for each sein just to close a circle the meter) = Then divide	example 7 yards X 3 = 200t) = total inches. Doyour catch by 320 to the event. If formula is: Net Lengide your catch by this	= 21 ft.) then existed by 39.3 get catch per gth (ft.)X number for y	convert it to 7 inches for meter sein 12 = total cour CPUE.	o r ed. <i>inches</i>
CDLIE	Catch Total				
Seine #Time_	Catch Total	Length of Pull	com	puted	
CPUE Time CPUE	Catch Total	Length of Pull	com	puted	

Hudson River Fish Fauna List

lamprey, silver 58				
a lamprey, sea (fi)	1	lamprey, silver		
			59	chub, creek (ñ)
Section	3	lamprey, sea (ñ)	60	fallfish (ñ)
6 dogfsh, spiny (f) 5 skate, little (f) 8 skate, barndoor (f) 8 sturgeon, shortnose (f) 9 sturgeon, shortnose (f) 10 sturgeon, atlantic (f) 11 gar, longnose 12 bowfin 13 lady fish (fi) 14 bonefish (fi) 15 cel, American (fi) 15 cel, American (fi) 16 worm cel, speckled (fi) 17 cel, conger (fi) 18 herring, bluchack (fi) 19 shad, hickory (fi) 19 shad, hickory (fi) 20 alewife (fi) 21 shad, American (fi) 22 menhaden, Atlantic (fi) 23 herring, atlantic (fi) 24 shad, sizzard 25 herring, round (fi) 26 anchovy, bay (fi) 27 anchovy, bay (fi) 38 stoneroller, central 39 goldfish 30 dace, redside 31 club, lake (fi) 32 carp, grass 31 shiner, satisfin (fi) 33 shiner, spotfin 34 shiner, spotfin 35 shiner, spotfin 36 shiner, spotfin 37 samon, variantic (fi) 38 shiner, spotfin 39 shiner, spotfin 40 shiner, spotfin 41 shiner, spotfin 42 shiner, spotfin 43 shiner, spotfin 45 shiner, spotfin 46 shiner, spotfin 47 pickerel, f(fi) 48 pickerel, f(fi) 49 shiner, spotfin 40 shiner, spotfin 41 dace, pear (fi) 42 chub, hornyhead 43 shiner, spotfin 45 shiner, comeral (fi) 46 shiner, comeral (fi) 47 pickerel, f(fi) 48 shiner, spotfin 49 pollock (fi) 40 shiner, spotfin 41 dace, pear (fi) 42 chub, hornyhead 43 shiner, spotfin 44 shiner, comeral (fi) 45 shiner, comeral (fi) 46 shiner, backethin 47 shiner, backethin 48 shiner, spotfin 49 herring, fike (rewhiting) (fi) 40 shiner, comeral (fi) 41 dace, pear (fi) 42 chub, hornyhead 43 shiner, spotfin 44 shiner, comeral (fi) 45 shiner, spotfin 46 shiner, barkethin 47 shiner, barkethin 48 shiner, spotfin 49 shiner, barkethin 40 shiner, barkethin 41 dace, pear (fi) 42 chub, hornyhead 43 shiner, spotfin 44 shiner, comeral (fi) 45 shiner, barkethin 46 shiner, barkethin 47 shiner, barkethin 48 shiner, spotfin 49 shiner, spotfin 50 dace, anorthern redbelly (fi) 51 dace, anorthern redbelly (fi) 52 dace, finescale (fi) 53 minnow, bustore (fi) 54 dace, east	4	shark (bull shark?) (ñ)	61	sucker, longnose (ñ)
8 skate, barndoor (f) 64 hog sucker, northern (f) 9 sturgeon, shortnose (f) 65 redborse, shorthead 10 sturgeon, Atlantic (f) 67 redborse, shorthead 11 gar, longnose 68 pirapitinga (Characin) 12 bowfin 69 caffsh, white (f) 13 ladyfish (f) 70 builhead, yellow (f) 14 bonefish (f) 71 builhead, yellow (f) 15 eel, American (f) 72 caffsh, channel 16 worm eel, speckled (f) 73 stonceat 16 worm eel, speckled (f) 73 stonceat 17 eel, conger (f) 74 madtom, tadpole (f) 18 herring, blueback (f) 75 madtom, tadpole (f) 19 shad, hickory (f) 76 madtom, tadpole (f) 21 shad, American (f) 78 pike, northern (f) 21 shad, Alantic (f) 79 pickerel, redfin (f) 22 menhaden, Atlantic (f) 79 <	5	dogfish, smooth (ñ)	62	sucker, white (ñ)
8 skafe, barndoor (f) 65 buffalo hybrid (black x smallmouth) 9 sturgeon, Atlantic (fi) 66 redbrose, shorthead 10 sturgeon, Atlantic (fi) 67 weatherfish, Oriental 11 gar, longnose 68 pirapitinga (Characin) 12 bowfin 69 catfish, white (fi) 13 ladyfish (fi) 70 bullhead, brown (fi) 14 bonefish (fi) 71 bullhead, brown (fi) 15 eel, American (fi) 72 carfish, channel 16 worm eel, speckled (fi) 73 stoncat 17 eel, conger (fi) 74 madtom, tadpole (fi) 18 herring, blueback (fi) 75 madtom, brindled 19 shad, hickory (fi) 76 madtom, brindled 20 alewife (fi) 77 pickerel, redfin (fi) 21 shad, American (fi) 78 pick, erel, chain (fi) 22 menhaden, Atlantic (fi) 79 pickerel, chain (fi) mudminnow, exstern (fi) 23 </td <td>6</td> <td>dogfish, spiny (ñ)</td> <td>63</td> <td>chubsucker, creek (ñ)</td>	6	dogfish, spiny (ñ)	63	chubsucker, creek (ñ)
8 skate, barndoor (ñ) 65 buffalo hybrid (black x smallmouth) 9 sturgeon, Atlantic (ñ) 66 reddores, shorthead 10 sturgeon, Atlantic (ñ) 67 weatherfish, Oriental 11 gar, longnose 68 pirapitinga (Characin) 12 bowfin 69 catfish, white (ñ) 13 ladyfish (ñ) 70 bullhead, brown (ñ) 14 bonefish (ñ) 71 bullhead, brown (ñ) 15 eel, American (ñ) 72 carfish, channel 16 worm eel, speckled (ñ) 74 madtom, tadpole (ñ) 17 eel, conger (ñ) 74 madtom, brindled 18 herring, blueback (ñ) 75 madtom, brindled 19 shad, hickory (ñ) 76 madtom, brindled 20 alewife (ñ) 77 pickerel, redfin (ñ) 21 shad, American (ñ) 78 pick, ercl. chain (ñ) 22 menhaden, Atlantic (ñ) 79 pickerel, chain (ñ) 23 herring, loval (ñ) <td< td=""><td>7</td><td>skate, little (ñ)</td><td>64</td><td>hog sucker, northern (ñ)</td></td<>	7	skate, little (ñ)	64	hog sucker, northern (ñ)
			65	buffalo hybrid (black x smallmouth)
11	9	sturgeon, shortnose (ñ)		
11	10	sturgeon, Atlantic (ñ)		
12				
14	12	bowfin		
15				
16				
17	15	eel, American (ñ)		
18	16	worm eel, speckled (ñ)		
19 Shad, hickory (fi) 76 madtom, brindled	17	eel, conger (ñ)		
20	18	herring, blueback (ñ)	75	madtom, margined (ñ)
21	19	shad, hickory (ñ)		
	20	alewife (ñ)		
menhaden, Atlantic (fi) herring, Atlantic (fi) herring, Atlantic (fi) herring, round (fi) anchovy, striped (fi) stouchy, bay (fi) stouchy, bay (fi) dace, pearl (fi) minnow, castern silvery (fi) shiner, common (fi) shiner, common (fi) shiner, common (fi) shiner, comerol (fi) shiner, spotdan (fi) shiner, common (fi) shiner, spotdan (fi) shiner				
24			70	muskellunge, tiger (norlunge)
24			79	pickerei, chain (n)
berring, round (i)	24	shad, gizzard		
anchovy, striped (ñ) anchovy, bay (ñ) anchovy, bay (ñ) stoneroller, central stoneroller, central goldfish acc, redside acc, eastern blacknose acc, bashiner, satinfin (ñ) acc, arp, grass aslmon, chinook acc, redside acc, eastern blacknose acc			91	
Stoneroller, central			02	mudminnow (nybrid) (n)
Stoneroller, central	27	anchovy, bay (ñ)		
29 goldfish 85	28	stoneroller, central		
State	29	goldfish		, , ,
Salmon, chinook Salmon, chinook Salmon, chinook Salmon, satinfin (ñ) Salmon, carp, common Salmon, carp, mirror (var.) Salmon, cutlips (ñ) Salmon, catalnatic (ñ) Salmon, cutlips (ñ) Salmon, cutlip	30	dace, redside	85	trout, rainbow
52 carp, grass 33 shiner, satinfin (ñ) 89 salmon, Atlantic (ñ) 34 shiner, spotfin 90 trout, brook (ñ) 35 carp, common 91 trout, brook (ñ) 36 minnow, cutlips (ñ) 92 trout, brook (ñ) 37 minnow, brassy (ñ) 93 lizardfish, inshore (ñ) 38 minnow, eastern silvery (ñ) 94 trout-perch (ñ) 39 shiner, bridle (ñ) 95 rockling, fourbeard (ñ) 40 shiner, common (n) 96 cod, Atlantic (n) 41 dace, pearl (ñ) 97 hake, silver (whiting) (n) 42 chub, hornyhead 98 tomcod, Atlantic (n) 43 shiner, golden (n) 99 pollock (n) 44 shiner, golden (n) 99 pollock (n) 45 shiner, emerald 101 hake, spotted (n) 46 shiner, blacknose 103 cusk-eel, striped (n) 47 shiner, blacknose 103 cusk-eel, striped (n)		chub, lake (ñ)	87	
Salmon, Atlantic (\hat{n})	32			
State Stat				
Carp, tolinion Carp, mirror (var.) 91		· · · · · · · · · · · · · · · · · · ·		
36 minnow, cutlips (ñ) 92 trout, lake (ñ) 37 minnow, brassy (ñ) 93 lizardfish, inshore (ñ) 38 minnow, eastern silvery (ñ) 94 trout-perch (ñ) 39 shiner, bridle (ñ) 95 rockling, fourbeard (ñ) 40 shiner, common (ñ) 96 cod, Atlantic (ñ) 41 dace, pearl (ñ) 97 hake, silver (whiting) (ñ) 42 chub, hornyhead 98 tomcod, Atlantic (ñ) 43 shiner, golden (ñ) 99 pollock (ñ) 44 shiner, comely (ñ) 100 hake, red (ling) (ñ) 45 shiner, emerald 101 hake, spotted (ñ) 46 shiner, blackchin 102 hake, white (ñ) 47 shiner, blackchose 103 cusk-cel, striped (ñ) 48 shiner, spottail (ñ) 104 toadfish, oyster (ñ) 49 shiner, rosyface 105 goosefish (anglerfish) (ñ) 50 shiner, rosyface 106 needlefish, Atlantic (ñ) 51 dace, nort		•		
minnow, brassy (ñ) 93				,
Shiner, bridle (fi) 95				
Same		, , , , , , , , , , , , , , , , , , , ,		<u> </u>
dace, pearl (n) 97				<i>5</i> ,
chub, hornyhead shiner, golden (ñ) shiner, comely (ñ) shiner, emerald shiner, blackchin shiner, blackchin shiner, spottail (ñ) shiner, sand shiner, sand shiner, rosyface dace, northern redbelly (ñ) dace, finescale (ñ) minnow, bluntnose minnow, bluntnose shiner, blacknose (ñ) shiner, blacknose 103 cusk-eel, striped (ñ) toadfish, oyster (ñ) goosefish (anglerfish) (ñ) needlefish, Atlantic (ñ) houndfish (ñ) blacknose (ñ) minnow, sheepshead minnow, sheepshead minnow, fathead shiner, castern blacknose (ñ) shiner, rosyface lid mummichog (ñ) skillifish, eastern banded (ñ) mummichog (ñ) killifish, striped (ñ)		• • •	· · ·	,
shiner, golden (ñ) shiner, comely (ñ) shiner, emerald shiner, blackchin shiner, blackchin shiner, spottail (ñ) shiner, spottail (ñ) shiner, sand shiner, rosyface dace, northern redbelly (ñ) ace, finescale (ñ) minnow, bluntnose minnow, fathead dace, eastern blacknose (ñ) shiner, golden (ñ) 100				
shiner, comely (ñ) shiner, comely (ñ) shiner, emerald shiner, blackchin shiner, spottail (ñ) shiner, spottail (ñ) shiner, spottail (ñ) shiner, sand shiner, rosyface shiner, rosyface dace, northern redbelly (ñ) adace, finescale (ñ) minnow, bluntnose minnow, fathead dace, eastern blacknose (ñ) dace, longnose (ñ) hake, red (ling) (ñ) loades, spottail (ñ) hake, red (ling) (ñ) hake, red (ling) (ñ) loades, spottail (ñ) hake, red (ling) (ñ) loades, spottail (ñ) lo		•		
shiner, emerald shiner, blackchin shiner, blackchin shiner, blackchin shiner, blacknose shiner, spottail (ñ) shiner, spottail (ñ) shiner, sand shiner, rosyface shiner, sand shiner, spottail (ñ) shiner, spottail (ñ) shiner, sand shiner, spottail (ñ) shiner, spottail (ñ) shiner, sand shiner, spottail (ñ)				<u>.</u>
46 shiner, blackchin 47 shiner, blacknose 48 shiner, spottail (ñ) 49 shiner, sand 50 shiner, rosyface 51 dace, northern redbelly (ñ) 52 dace, finescale (ñ) 53 minnow, bluntnose 54 minnow, fathead 55 dace, eastern blacknose (ñ) 56 dace, longnose (ñ) 57 hake, white (ñ) 58 cusk-eel, striped (ñ) 59 goosefish, oyster (ñ) 50 needlefish, Atlantic (ñ) 50 needlefish, Atlantic (ñ) 51 houndfish (ñ) 52 houndfish (ñ) 53 minnow, sheepshead 54 minnow, fathead 55 dace, eastern blacknose (ñ) 56 dace, longnose (ñ) 57 little November 102 hake, white (ñ) 58 number of (ñ) 59 hake, white (ñ) 50 musk-eel, striped (ñ) 50 meedlefish, Atlantic (ñ) 50 minnow, sheepshead 51 minnow, sheepshead 52 minnow, fathead 53 minnow, fathead 54 minnow, fathead 55 dace, eastern blacknose (ñ) 56 dace, longnose (ñ) 57 little November of (ñ) 58 killifish, spotfin (ñ)				
shiner, blacknose shiner, spottail (ñ) shiner, sand shiner, sand shiner, sand shiner, sand shiner, sand shiner, rosyface shiner, sand shillifish, eastern bandic (ñ) shillifish, shiner, sand shiner, sand shiner, sand shiner, sand shillifish, eastern bandic (ñ) shillifish, striped (ñ) shillifish, spotfin (ñ)				
48 shiner, spottail (ñ) 49 shiner, sand 50 shiner, rosyface 51 dace, northern redbelly (ñ) 52 dace, finescale (ñ) 53 minnow, bluntnose 54 minnow, fathead 55 dace, eastern blacknose (ñ) 56 dace, longnose (ñ) 57 toadfish, oyster (ñ) 58 goosefish (anglerfish) (ñ) 59 needlefish, Atlantic (ñ) 50 needlefish, Atlantic (ñ) 50 needlefish, Atlantic (ñ) 51 houndfish (ñ) 52 minnow, sheepshead 53 minnow, bluntnose 54 minnow, fathead 55 dace, eastern blacknose (ñ) 56 dace, longnose (ñ) 57 lill killifish, spotfin (ñ)				
shiner, sand shiner, rosyface shiner, rosyface 106 needlefish, Atlantic (ñ) houndfish (ñ) dace, northern redbelly (ñ) 107 houndfish (ñ) minnow, sheepshead 108 minnow, sheepshead 109 killifish, eastern banded (ñ) mummichog (ñ) dace, eastern blacknose (ñ) 111 killifish, spotfin (ñ)				
shiner, rosyface dace, northern redbelly (ñ) dace, finescale (ñ) minnow, bluntnose minnow, fathead dace, eastern blacknose (ñ) dace, longnose (ñ) shiner, rosyface 106 needlefish, Atlantic (ñ) houndfish (ñ) minnow, sheepshead killifish, eastern banded (ñ) mummichog (ñ) killifish, striped (ñ) killifish, spotfin (ñ)				
dace, northern redbelly (ñ) dace, finescale (ñ) minnow, sheepshead minnow, sheepshead minnow, fathead dace, eastern blacknose (ñ) dace, longnose (ñ) loo minnow, sheepshead mummichog (ñ) mummichog (ñ) lil killifish, striped (ñ) killifish, spotfin (ñ)				
dace, finescale (ñ) dace, finescale (ñ) minnow, sheepshead log killifish, eastern banded (ñ) mummichog (ñ) dace, eastern blacknose (ñ) dace, longnose (ñ) lil killifish, spotfin (ñ)		· · · · · · · · · · · · · · · · · · ·		
minnow, bluntnose minnow, fathead dace, eastern blacknose (ñ) dace, longnose (ñ) minnow, fathead 110 mummichog (ñ) killifish, striped (ñ) killifish, spotfin (ñ)		· · · · · · · · · · · · · · · · · · ·		
minnow, fathead 55 dace, eastern blacknose (ñ) 56 dace, longnose (ñ) 110 mummichog (ñ) killifish, striped (ñ) killifish, spotfin (ñ)				
dace, eastern blacknose (ñ) dace, longnose (ñ) 111 killifish, striped (ñ) killifish, spotfin (ñ)				
dace, longnose (ñ) 112 killifish, spotfin (ñ)				9
Tiz kinnish, spotim (n)				
113 mosquitofish, western				
		. •	113	mosquitonsn, western

114	114	silvansida huaali	172	.*
16 Silverside, Inland (f)	114	silverside, brook		
		_		1 1 00, 1 1
18		= * * *		
19				
12				
122				
123				
124 pipefish, northern (ñ) 181 butterfly fish, fourcyc (ñ)				
124				
183				
184				
185				
186				
187				
130				
Sculpin, longhorn (R)				
132				
133				
192				
135 bass, white 193 sand lance, American (sand eel) (n) 136 bass, striped (n) 194 sleeper, fat (n) 137 sea bass, black (n) 195 goby, naked (n) 138 gag (grouper) (n) 196 goby, seaboard (n) 139 bass, rock 197 goby, seaboard (n) 140 sunfish, bluespotted (n) 198 goby, highfin (n) 141 sunfish, banded (n) 199 mackerel, Atlantic (n) 142 sunfish, redbreast (n) 200 mackerel, Spanish (n) 143 sunfish, green 201 butterfish (n) 144 pumpkinseed (n) 202 snakehead, northern (n) 145 warmouth 203 flounder, Gulf Stream (n) 146 bluegil 204 flounder, summer (fluke) (n) 147 bass, smallmouth 205 flounder, summer (fluke) (n) 148 bass, largemouth 206 flounder, fourspot (n) 149 crappie, white 206 flounder, winter (n) 150 crappie, black 207 windowpane (n) 151 darter, greenside 208 flounder, winter (n) 152 darter, fantail 209 flounder, winter (n) 153 darter, tessellated 210 tonguefish, northern (n) 154 perch, yellow (n) 211 hogchoker (n) 155 logperch 212 filefish, orange (n) 156 darter, shield 213 filefish, planehead (n) 157 walleye 214 burrfish, striped (n) 158 bigeye, short (n) 215 puffer, smooth (n) 159 bluefish (n) 215 puffer, smooth (n) 160 cobia (n) Class 4 161 sharksucker (n) Class 4 162 jack, crevalle (n) Greera 154 163 scad, round (n) Greera 154 164 moonfish, Atlantic (n) Greera 154 165 lookdown (n) Greera 154 166 permit (n) Greera 154 167 schoolmaster (n) Greera 154 168 snapper, gray (mangrove) (n) Species 217 169 mojarra, spotfin (n) Greera 154 160 mojarra, spotfin (n) Greera 154 161 sharksucker (n) Greera 154 162 mojarra, spotfin (n) Greera 154 163 mojarra, spotfin (n) Greera 154 164 mojarra, spotfin (n) Greera 154 165 mojarra, sp				
136				
137		= '		= / / / / / /
138				
139		-		
140				
141				
142				
143				
144				
145				
146				- · · · · · · · · · · · · · · · · · · ·
147				
bass, largemouth 205				
149		= /		
150				- · · · · · · · · · · · · · · · · · · ·
152	150			
153	151	darter, greenside		
154		_ darter, fantail		
155				
156				_ hogchoker (ñ)
157				
158				_ filefish, planehead (ñ)
bluefish (ñ) cobia (ñ) sharksucker (ñ) idi sharksucker (ñ) idi sacad, round (ñ) idi moonfish, Atlantic (ñ) idi class 4 idi order 26 idi permit (ñ) idi Families 75 idi schoolmaster (ñ) idi Species 217 idi mojarra, spotfin (ñ)		- '		
160			215	_ puffer, smooth (ñ)
sharksucker (ñ) 162			216	_ puffer, northern (ñ)
jack, crevalle (ñ) jack, crevalle (ñ) (ñ) = Native Species (166, or 76%) Taxonomic diversity: Class 4 lookdown (ñ) Order 26 permit (ñ) schoolmaster (ñ) schoolmaster (ñ) snapper, gray (mangrove) (ñ) mojarra, spotfin (ñ) pigfish (ñ) (ñ) = Native Species (166, or 76%) Taxonomic diversity: Class 4 Order 26 Families 75 Genera 154 Species 217 Tom Lake NYSDEC Hudson River Estuary Naturalist Aug		-	217	_ cowfish, scrawled (ñ)
scad, round (ñ) Taxonomic diversity: Class 4 165 lookdown (ñ) Order 26 permit (ñ) schoolmaster (ñ) schoolmaster (ñ) schoolmaster (ñ) mojarra, spotfin (ñ) pigfish (ñ) Taxonomic diversity: Class 4 Order 26 Families 75 Genera 154 Species 217 Tom Lake NYSDEC Hudson River Estuary Naturalist Aug		-		
moonfish, Atlantic (ñ) Class 4 lookdown (ñ) Order 26 permit (ñ) Families 75 schoolmaster (ñ) Genera 154 snapper, gray (mangrove) (ñ) mojarra, spotfin (ñ) pigfish (ñ) Taxbitomic diversity. Class 4 Order 26 Families 75 Genera 154 Species 217 Tom Lake NYSDEC Hudson River Estuary Naturalist Aug			(ñ) = Nati	ive Species (166, or 76%)
lookdown (ñ) Order 26 permit (ñ) Families 75 for schoolmaster (ñ) Genera 154 snapper, gray (mangrove) (ñ) Species 217 mojarra, spotfin (ñ) pigfish (ñ) Tom Lake NYSDEC Hudson River Estuary Naturalist Aug			Taxonon	nic diversity:
permit (ñ) Families 75 schoolmaster (ñ) Genera 154 snapper, gray (mangrove) (ñ) Mojarra, spotfin (ñ) Tom Lake NYSDEC Hudson River Estuary Naturalist Aug pigfish (ñ)			Class	4
167 schoolmaster (ñ) 168 snapper, gray (mangrove) (ñ) 169 mojarra, spotfin (ñ) 170 pigfish (ñ) Genera 154 Species 217 Tom Lake NYSDEC Hudson River Estuary Naturalist Aug		=	Order	26
167 schoolmaster (ñ) Genera 154 168 snapper, gray (mangrove) (ñ) Species 217 169 mojarra, spotfin (ñ) Tom Lake NYSDEC Hudson River Estuary Naturalist Aug 170 pigfish (ñ)			Families	75
169 mojarra, spotfin (ñ) 170 pigfish (ñ) Tom Lake NYSDEC Hudson River Estuary Naturalist Aug		-	Genera	154
169 mojarra, spotfin (ñ) Tom Lake NYSDEC Hudson River Estuary Naturalist Aug 170 pigfish (ñ)			Species	217
170 pigfish (ñ)			-	e NYSDEC Hudson River Estuary Naturalist Aug
171 sheepshead (n)				
	171	_ sheepshead (ñ)		

Snapshot Day Activity X- Other Observations

1. Shipping.

Should you see large ships, tugs, or barges pass your site, note the following information if possible. A loaded ship or barge is full of cargo, and rides lower in the water than a light - empty - vessel. Binoculars are helpful in gathering the information requested here. Note recreational ships if you like in order to note the different types of usage of the river.

Time	Type of ship	Name	Northbound/ Southbound/	cargo

2. Other items of interest.

Feel free to record any other observations. This could include birds seen, items found on the beach, or any other things you find interesting

Snapshot Day Activity XI- Journaling & A Hudson River Almanac Entry

How do we learn about our natural environment? We observe.

Through this activity we are focusing on developing skills of observation that play such an important role in science and Earth systems. Direct observation and careful description helps us compare species, habitats and different geographical regions. Through journaling we hope to observe, record and better understand some of the relationships that exist in the natural world.

Students have a sheet for recording observations during the day. In addition to their personal observations, at the end of the day the space below can be used for drafting a Hudson River Almanac entry from each site, which will be submitted with your data (use additional paper as needed, but keep them to 4-6 sentences please). Sample entry:

September 29 - Dobbs Ferry - Our beach seine was filled with nearly 600 fish-snapper blues, white perch, a vast school of silverside, and several 4"-7" striped bass. Low flying monarchs passed in twos and threes, dipping within inches, brushing against us as they beat into strong southerlies. The students from Irvington were thrilled to be so close to so much loveliness. As they passed, the students called out the tally; they were moving past us at the rate of fifty an hour. Christopher Letts

Hudson River Almanac Entry: