

1 Early Jurassic magnetostratigraphy and paleolatitudes from the Hartford
2 continental rift basin (eastern North America): Testing for polarity bias and abrupt
3 polar wander in association with the Central Atlantic Magmatic Province
4

5 Dennis V. Kent
6

7 Earth and Planetary Sciences, Rutgers University, Piscataway NJ 08854 &
8 Lamont-Doherty Earth Observatory, Palisades NY 10964
9

10 Paul E. Olsen
11

12 Lamont-Doherty Earth Observatory of Columbia University, Palisades NY 10964
13

14 **Abstract**
15

16 To determine whether the ~200 Ma Central Atlantic magmatic province (CAMP)
17 coincides with a normal polarity bias and a purported abrupt change in polar wander at the J1
18 cusp, we collected samples for paleomagnetic study from 80 sites distributed over a ~2500 m-
19 thick section of sedimentary units that are interbedded with and overlie CAMP lavas in the
20 Hartford basin, which together represent the initial 2.4 Myr of the Jurassic according to cycle
21 stratigraphic analysis. Characteristic directions carried by hematite were isolated by thermal
22 demagnetization in 71 sites and define a coherent magnetostratigraphy supported by a positive
23 reversal test and an inter-basin fold test. Despite a pronounced overall normal polarity bias (only
24 three relatively short reverse polarity intervals could be confirmed in the sampled section),
25 normal polarity Chron H24n that encompasses the CAMP extrusive zone is no more than 1.6
26 Myr in duration. Elongation/inclination analysis of the 315 characteristic directions, which have
27 a flattened distribution, produces a result in agreement with a published mean direction for the
28 CAMP volcanic units as well as published results similarly corrected for inclination error from
29 the Newark basin. The three datasets – CAMP volcanics, Newark corrected sediments and
30 Hartford corrected sediments – provide a 201 Ma reference pole for eastern North America at

31 67.0°N 93.8°E $A_{95}=3.2^\circ$. Paleopoles from the Moenave and Wingate Formations from the
32 Colorado Plateau that virtually define the J1 cusp can be brought into agreement with the 201 Ma
33 reference pole with corrections for net clockwise rotation of the Plateau relative to eastern North
34 America and presumed sedimentary inclination error. The corrected data show that apparent
35 polar wander for North America proceeds directly toward higher latitudes over the Late Triassic
36 and Early Jurassic with no obvious change that can be associated with CAMP.

37 Introduction

38

39 The recent recognition of what may be the largest igneous province on Earth, the ~200 Ma
40 Central Atlantic magmatic province (CAMP; *Marzoli et al.*, 1999) (**Fig. 1**), with its close temporal
41 proximity to major biotic turnover at the Triassic-Jurassic boundary [*Olsen*, 1999], adds impetus
42 for seeking confirmation of possibly related phenomena. One is an apparently extended interval of
43 pronounced normal polarity bias that has been found in several data compilations. An early global
44 assessment of paleomagnetic data by [*Irving and Pullaiah*, 1976] suggested there was a poorly
45 defined normal polarity interval in the Triassic, which roughly coincided with the Graham normal
46 interval of *McElhinny and Burek* [1971] and the Newark normal interval of *Pechersky and*
47 *Khramov* [1973]. Although no supporting evidence of an extended normal polarity interval has
48 subsequently been found in magnetostratigraphic data for the Triassic [*Steiner et al.*, 1989; *Ogg*
49 *and Steiner*, 1991; *Gallet et al.*, 1992; *Kent et al.*, 1995; *Muttoni et al.*, 1998; *Szurliés*, 2004], a ~10
50 Myr-long interval of pronounced normal polarity bias and low reversal frequency has been
51 identified in the early Jurassic in several compilations of global paleomagnetic data [*Johnson et*
52 *al.*, 1995; *Algeo*, 1996] and could conceivably be related to perturbation of the geodynamo by
53 ascent of a mantle plume (e.g., *Larson and Olson*, 1991).

54

55 CAMP emplacement also seems to coincide temporally with an abrupt change in North
56 American apparent polar wander (APW) at the so-called J1 cusp [*Gordon et al.*, 1984; *May and*
57 *Butler*, 1986], which was suggested to reflect a major plate reorganization [*Gordon et al.*, 1984] or
58 an episode of true polar wander [*Marcano et al.*, 1999], either of which could have been associated
59 with the emplacement of CAMP. Late Triassic paleopoles from the Newark Supergroup are
60 coherent between basins and do not have cusp-like directions [*Witte et al.*, 1991; *Kent et al.*, 1995;
61 *Kent and Olsen*, 1997] whereas the apparent trend toward the J1 cusp in coeval paleopoles from
62 the southwestern U.S. can be explained by a ~10–15° clockwise rotation of the Colorado Plateau
63 region [*Kent and Witte*, 1993; *Steiner and Lucas*, 2000]. However, early Jurassic paleopoles from
64 the Moenave and Wingate Formations on the Colorado Plateau [*Ekstrand and Butler*, 1989;
65 *Molina-Garza et al.*, 2003] that virtually define the J1 cusp have few reliable counterparts from
66 other parts of North America, where J1 cusp-like directions have thus far been reported only as
67 overprints with uncertain age control from Texas [*Molina-Garza et al.*, 1995] and in some baked

68 sediment sites in contact with CAMP-related igneous intrusions, and thus representing abbreviated
69 recordings of the paleofield with uncertain structural control, in eastern North America [*Kodama et*
70 *al.*, 1994].

71
72 Although there is no obvious evidence of either a normal polarity superchron or the J1
73 cusp in the Late Triassic record from the Newark Supergroup, more data are clearly needed to
74 document the polarity history and APW for North America in the early Jurassic when the CAMP
75 event actually occurred. Paleomagnetic and chronostratigraphic data are already available from
76 the Late Triassic and earliest Jurassic history of the Newark basin and provide some of the best
77 available age constraints on the age and duration of CAMP igneous activity [*Olsen et al.*, 1996b;
78 *Hames et al.*, 2000]. The oldest lavas in the Newark basin immediately postdate (within ~40 kyr
79 by cycle stratigraphy) the Triassic-Jurassic boundary identified on the basis of palynoflora and
80 vertebrate (mainly footprint) evidence coinciding with a small iridium anomaly in a boundary
81 clay layer [*Olsen et al.*, 2002a]. A reverse polarity interval (Chron E23r) occurs just below the
82 Triassic-Jurassic boundary; with an estimated duration of only ~20,000 y, it is the shortest
83 polarity interval amongst the 60 polarity chrons delineated in the astronomical polarity time scale
84 based on the Newark succession [*Kent and Olsen*, 1999]. Despite its brevity, the occurrence of
85 Chron E23r in close proximity to the Triassic-Jurassic boundary has evidently made it a
86 beguiling target for correlation of distant sections, such as St. Audrie's Bay in Britain [*Hounslow et*
87 *al.*, 2004] and the High Atlas of Morocco [*Marzoli et al.*, 2004]. The succeeding normal polarity
88 interval (Chron E24n) already encompasses more than 1000 meters of CAMP lavas and
89 interbedded sedimentary formations [*McIntosh et al.*, 1985; *Witte et al.*, 1991], making it the
90 thickest polarity zone in the Newark basin stratigraphic succession [*Kent and Olsen*, 1999] even
91 though its upper limit has not been found in the overlying Boonton Formation, which apart from
92 becoming conglomeratic close to the border fault, is largely buried by Pleistocene glacial
93 deposits. The full duration of Chron E24n, whose known record already constitutes one of the
94 longer polarity intervals in the Newark geomagnetic polarity time scale (GPTS) [*Kent and Olsen*,
95 1999], has yet to be determined. Correspondingly, the search for J1 cusp directions has not
96 extended over more than about the first ~600 kyr of the Jurassic in the Newark Supergroup
97 record.

98

99 More extensive Jurassic-age deposits are preserved in the nearby Hartford basin, another
100 of the series of exhumed continental rift basins outcropping from Nova Scotia to North Carolina
101 that are filled with continental strata of the Newark Supergroup (**Fig. 2a**). Of all the exposed
102 Mesozoic rift basins in eastern North America, the Hartford basin has the thickest section of
103 continental strata of Early Jurassic age, totaling at least 4500 m (**Fig. 2b, c**). These Jurassic-age
104 strata have long been recognized as containing redbeds and cyclical lacustrine sediments [*Hubert*
105 *et al.*, 1992] but there has been no comprehensive attempt to develop a magnetostratigraphy or to
106 describe the cyclical sequence as a whole. In this paper we focus on the paleomagnetism and
107 cyclostratigraphy of the lower 2500 m of the Jurassic age section, which is the fine grained and
108 cyclical portion that begins with sedimentary units (Shuttle Meadow and East Berlin Formations)
109 that are interbedded with the CAMP lavas and extends into the lower half of the Portland
110 Formation (**Fig. 2c**). We describe and interpret the cyclicity in terms of Milankovitch orbital
111 variations [*Olsen and Kent*, 1996], providing a basis for extending the astronomically calibrated
112 GPTS for the Late Triassic [*Kent and Olsen*, 1999] into the Early Jurassic and a
113 chronostratigraphic context for paleopoles.

114

115 Geologic Framework of Hartford basin

116

117 The Newark, Hartford and related early Mesozoic continental rift basins that are
118 preserved on the margins of the Atlantic–bordering continents formed during the incipient break-
119 up of Pangea in the Triassic (**Fig. 1**). The overall structure of the Hartford basin (**Fig. 2b**) is
120 consistent with a step-faulted half-graben geometry as seen in other Newark Supergroup basins
121 [*Schlische*, 1993]. However, unlike the other exposed rift basins in eastern North America that
122 have their long axes oriented northeast-southwest, the Hartford basin runs nearly north-south
123 with a segmented west-dipping border fault system on its eastern side towards which the basin
124 strata tilt at predominately low to moderate (~10-15°) dips. The border fault system generally
125 parallels the structural fabric of Paleozoic metamorphic basement, suggesting that the border
126 faults may be reactivated structures [*Wise and Robinson*, 1982]. Numerous generally northeast-
127 trending intrabasinal faults with strike-slip and down-to-the-west normal offsets occur especially
128 in the southern and central portions of the basin (e.g., *Davis*, 1898; *Sanders*, 1970). These faults,
129 as well as a series of transverse folds that increase in amplitude and frequency to the east towards

130 the border fault system [*Wheeler, 1937; Schlische, 1995*], complicate the homoclinal geometry of
131 the basin strata. Fission track analyses suggest moderate (2-5 km) burial depths [*Roden and*
132 *Miller, 1991; Roden-Tice and Wintsch, 2002*].

133

134 The large scale lithostratigraphy of the basin-fill is comprised of a tripartite succession
135 (**Fig. 2c**): (1) a lower coarse arkosic unit up to 3000 m-thick named the New Haven Formation;
136 (2) a middle, generally finer grained sequence (the focus of this paper) containing interbedded
137 basalt flows of the CAMP near its base, which together are about 2500 m thick and consist of, in
138 ascending order, the Talcott Basalt, Shuttle Meadow Formation, Holyoke Basalt, East Berlin
139 Formation, Hampden Basalt, and about the lower half of the Portland Formation; and (3) an
140 upper coarse arkosic unit that exceeds 1500 m in thickness comprising the upper Portland
141 Formation (e.g., *Krynine, 1950; Sanders, 1968*). A Jurassic age for the middle and upper
142 succession is based on both palynology [*Cornet et al., 1973; Cornet and Traverse, 1975; Cornet*
143 *and Olsen, 1985*] and vertebrate biostratigraphy [*Lucas and Huber, 2003; Olsen and Galton,*
144 *1977*]. Radioisotopic dates from the CAMP basaltic flows interbedded with these strata have
145 substantial scatter attributed in large part to post-cooling alteration but are not inconsistent with a
146 Jurassic age [*Seidemann, 1989*]. Geochemical and cyclostratigraphic correlation with other
147 basins that have igneous rocks with more secure radioisotopic dates [*Sutter, 1988; Dunning and*
148 *Hodych, 1990; Hames et al., 2000*] support an earliest Jurassic (~200 Ma) age for the basalts
149 [*Olsen et al., 2003*]. Mafic igneous sills (e.g., Barndoor, West Rock, East Rock, Carmel) and
150 dikes (e.g., Buttress, Higganum) intrude the New Haven Formation; some of the dikes evidently
151 served as feeders for the tholeiitic lava flows [*Philpotts and Martello, 1986*] but they do not
152 seem to cut the Portland Formation. Parenthetically, the Deerfield basin is connected to the
153 Hartford basin (**Fig. 2b**) and has an analogous stratigraphic development except that only one
154 basalt unit (Deerfield Basalt, equivalent to the Holyoke Basalt of the Hartford basin [*Luttrell,*
155 *1989; Prevot and McWilliams, 1989; Tollo and Gottfried, 1992*]) is present.

156

157 Published paleomagnetic work in the Hartford basin has focused almost entirely on the
158 CAMP lavas and dikes [*DuBois et al., 1957; Irving and Banks, 1961; De Boer, 1968; Prevot and*
159 *McWilliams, 1989; Smith, 1976; Smith and Noltimier, 1979; see summary by Hozik, 1992*]. A
160 notable exception is the early work of *DuBois [1957]* who reported paleomagnetic results from

161 32 samples of “Triassic rocks from the Connecticut Valley in the State of Connecticut”, whose
162 sampling localities are otherwise not described. Normal and reverse polarity directions were
163 reported by *DuBois* [1957] but, to our knowledge, there has been no follow-up paleomagnetic
164 work on sediments of the Hartford basin until the present study. From the contiguous Deerfield
165 basin, paleomagnetic results from two sedimentary sites (from the Fall River Formation, a unit
166 correlative to the Shuttle Meadow Formation of the Hartford basin) were reported by *McEnroe*
167 *and Brown* [2000].

168

169 Paleomagnetic sampling and measurements

170

171 The Jurassic sedimentary units in the Hartford basin - Shuttle Meadow, East Berlin and
172 Portland formations - were the focus of sampling. Samples were collected from 80 sites in stream
173 and road cuts with a gasoline-powered portable drill and oriented with a magnetic compass. A
174 sampling site typically consisted of 4–6 oriented cores covering several meters of section;
175 sampling at three sites ranged over several tens of meters of section and included 8 to 12 samples
176 each, whereas sampling at 5 sites included only a single oriented hand sample from which up to
177 3 specimens were cut and measured. Bedding attitude was measured at every sampling site for
178 tilt corrections. Outcrops were sporadic due to the low topographic relief and shallow bedding
179 dips; hence, a stratigraphic composite section was assembled from a number of across-strike
180 profiles that were linked by tracing and mapping distinctive beds over the course of numerous
181 field trips to the area. Most of the sampling sites in the Portland Formation were collected in two
182 traverses, the Stony Brook (Connecticut) and Westfield River (Massachusetts) sections;
183 additional stratigraphic coverage was obtained from around Holyoke and South Hadley Falls
184 (Massachusetts) along the Connecticut River, the Chicopee River around Chicopee
185 (Massachusetts) and in the Durham-Portland area in Connecticut. Geologic mapping in the
186 Portland Formation required to link the sampling sites into a composite section resulted in the
187 recognition of new lithostratigraphic members, which were described informally by *Olsen et al.*
188 [2005] and formally in a forthcoming paper [*Olsen and Kent, 2008*] and provide a cycle
189 stratigraphic framework. We believe that our sampling at the 80 sites virtually exhausts available
190 outcrop of the Shuttle Meadow, East Berlin and lower Portland Formations. However, we were
191 able to fill in some gaps for magnetostratigraphy and description of lithostratigraphy for the

192 poorly exposed lowest portion of the Portland Formation by gaining access to some relatively
193 short engineering cores for the Park River project, as described below.

194

195 All magnetic remanence measurements were made on 2G 3-axis DC-SQUID
196 superconducting rock magnetometer housed in a magnetically shielded room (<1000 nT).
197 Thermal demagnetization was performed in a custom-built oven with 3 independent heating
198 zones and a water jacket for reproducible temperature control and housed in high magnetic
199 permeability shields for a low magnetic field environment (<5nT) that is critical for resolving
200 ancient magnetizations that can be masked by spurious magnetizations introduced by lab-
201 induced thermochemical alteration, which was monitored by measuring the magnetic
202 susceptibility with a Bartington instrument after each thermal demagnetization step.

203

204 Progressive thermal demagnetization of NRM reveals a relatively straightforward two-
205 component structure in most samples, comprising a low unblocking temperature (up to 300°C) A
206 component that tends to be aligned about the northerly and down present-day field direction
207 followed by a characteristic magnetization (ChRM) with shallow directions to the north (**Fig. 3a,**
208 **b, e, f, h**) or to the south (**Fig. 3c, d, g**) that converges toward the origin by maximum
209 unblocking temperatures of 685°C. The only systematic departure from this pattern was observed
210 in samples from a site in the East Berlin Formation, which tended to show back-tracking in
211 demagnetization trajectories between about 200° and 400°C (**Fig. 3f**) that could represent a
212 partial reverse polarity overprint embedded between normal polarity A and ChRM components.
213 What is absent in the Hartford samples, however, is a distinct northerly and moderately down
214 component with intermediate unblocking temperatures (~300° to at least 600°C) that was found
215 to be ubiquitous in sedimentary rocks in the Newark and Dan River basins and attributed to a
216 Middle Jurassic remagnetization event [*Witte and Kent, 1991; Kent et al., 1995; Kent and Olsen,*
217 *1997*]. It is possible that the Hartford basin escaped this remagnetization event, perhaps because
218 of its shallower burial and/or lower thermal maturation [*Pratt et al., 1988; Roden and Miller,*
219 *1991*]. Alternatively, the remagnetization event as identified in the Newark and Dan River basins
220 had in fact occurred during CAMP igneous activity and is therefore not expected to be as
221 pronounced or ubiquitous in the Hartford basin rocks we sampled, which are of CAMP age or
222 younger.

223

224 IRM analysis

225

226 The dominant magnetic mineralogy was characterized using isothermal remanent
227 magnetization (IRM) acquisition to 2.5 T using an ASC pulse magnetizer and thermal
228 demagnetization of orthogonal axes IRM by the method of *Lowrie* [1990]. Most of the sampling
229 was done with a preference for red mudstones and siltstones rather than the gray shales and thus
230 a hematite carrier of remanence is expected in most of the sites. IRM acquisition curves (**Fig. 4a**)
231 generally show a gradual approach to saturation to 2.5T although the gray samples (e.g.,
232 JPDA6b, JPDE6b) have remanent coercivities of around 170 mT, which are considerably lower
233 than around 600 mT for the red samples (e.g., JPDM4b, JPDU1b, 160702B05). Thermal
234 demagnetization of IRM (**Fig. 4b-f**) confirms that the dominant and higher coercivity (>200 or
235 >300 mT) component of an orthogonal IRM is invariably associated with a maximum
236 unblocking temperature of about 685°C, which is compatible with hematite. The lower
237 coercivity (<200 or <300 mT) component of the orthogonal IRM shows an inflection in its
238 thermal demagnetization spectrum at around 580°C suggesting the presence of magnetite in the
239 gray samples (**Fig. 4e-f**), whereas the red samples are dominated by hematite (maximum
240 unblocking temperature of about 685°C) in both high and low coercivity fractions.

241

242 Hematite-dominated sampling sites can possess either normal or reverse polarity
243 (northerly or southerly) characteristic directions, for example, sample JPDU1b is from a reverse
244 polarity site (**Fig. 4c**) and samples 160702B05 and JPDM4b are from normal polarity sites (**Fig.**
245 **4a, d**); likewise for sampling sites in gray beds that show contributions from magnetite, for
246 example, sample JPDA6b is from a reverse polarity site and sample JPDE6b is from a normal
247 polarity site (**Fig. 4e, f**). Nevertheless, the characteristic remanence is invariably associated with
248 the hematite phase (unblocking temperature above ~600°C) even in the gray beds that also have
249 a magnetite contribution (**Fig. 3c, h**). We suggest that the characteristic remanence is carried by
250 detrital hematite or an early authigenic product that was acquired during or soon after deposition
251 whereas the sporadic presence of magnetite that is mostly restricted to gray shales, such as the
252 Chicopee fish bed, may reflect its preservation or production in a localized reducing depositional
253 environment.

254

255 Magnetochemical alteration of the samples during thermal demagnetization is common as
256 indicated by changes in magnetic susceptibility, which usually starts to increase noticeably after
257 about the 500°C step. In the gray samples, magnetic susceptibility often continues to increase by
258 more than an order of magnitude by 680°C (**Fig. 4e, f**). The large monotonic susceptibility rise
259 with thermal treatment is associated with the production of magnetite, as revealed by IRM
260 acquisition (**Fig. 5a**) and thermal demagnetization of orthogonal IRM experiments on previously
261 heated samples (**Fig. 5b**), showing that a large IRM phase that approaches saturation by 300 mT
262 and has maximum unblocking temperatures around 575°C becomes the dominant magnetization
263 component after heating. In contrast, red siltstone samples that had only modest susceptibility
264 changes with thermal treatment (e.g., **Fig. 4c**) maintained a predominantly hematite mineralogy
265 characterized by lack of saturation and high (~685°C) maximum unblocking temperatures (**Fig.**
266 **5e, f**). Nevertheless, sediment grain size rather than just color (or initial magnetic mineralogy)
267 seems to be an important determinative factor in the thermal alteration profile. Gray samples that
268 showed large susceptibility changes with heating were typically fine-grained mudstones or
269 shales; however, some mudstones are also reddish and although their magnetizations are
270 dominated by hematite, many of these samples showed large susceptibility increases with initial
271 heating (e.g., sample 160702B05; **Fig. 4a**) that were also associated with the production of
272 magnetite (**Fig. 5c, d**), as in the gray shales. We suspect that the dramatic magnetochemical
273 alteration seen in laboratory heating of the fine-grained gray shales and red mudstones is
274 probably due to the breakdown of clays. Clays are much less abundant in the red siltstones and
275 fine-grained sandstones, making such hematite-bearing rocks less prone to magnetochemical
276 alteration and thereby enhancing their suitability for paleomagnetic study.

277

278 Magnetization directions

279

280 NRM component vectors were estimated using principal component analysis [*Kirschvink*,
281 1980] on demagnetization trajectories typically from 100° to 300°C for the A component and
282 anchored to the origin from 600° to 675°C for the ChRM (except for three sites in fine-grained
283 gray to purplish shales – JPAA and JPAB in the lowermost Portland Formation, and 160702B in
284 the East Berlin Formation – that were analyzed for ChRM between 400° and 600°C due to the

285 onset of erratic directions associated with large susceptibility increases that occurred at higher
286 temperatures). Line-fits with MAD angles greater than 18° were rejected as were data from a few
287 samples that were obviously misoriented when compared with other samples at a site. Out of a
288 total of 398 samples measured and analyzed, over 83% (331 from 78 sites) yielded an acceptable
289 A component direction (median and mean MAD angles of 2.8° and 3.8° , respectively) and nearly
290 80% (315 from 71 sites) yielded acceptable estimates of a ChRM direction (median and mean
291 MAD angles of 3.5° and 5° , respectively).

292

293 The 78 site-mean A component directions from the Shuttle Meadow, East Berlin and
294 Portland Formations are well grouped in geographic coordinates about a mean direction of
295 declination (D) = 353.9° , inclination (I) = 60.2° , with a radius of the circle of 95% confidence
296 (a_{95}) = 1.6° (**Fig. 6**). This is close to a modern field direction (D = 346° I = 68° for present-day
297 field, D = 0° I = 61° for geocentric axial dipole field); moreover, the distribution is significantly
298 more dispersed (precision parameter k decreasing a factor of 2.6; **Table 1**) after bedding tilt
299 corrections, indicative of a negative fold test. The low unblocking temperature A component is
300 thus most probably a recently acquired viscous magnetization.

301

302 The 13 fully oriented sites from the Shuttle Meadow and East Berlin Formations all have
303 northerly and shallow (normal polarity) ChRM directions, whereas in the case of the 58 sites
304 from the Portland Formation, 45 had northerly and shallow (normal polarity) ChRM directions
305 and 13 sites had southerly and shallow (reverse polarity) ChRM directions (**Fig. 7**). The
306 combined tilt-corrected mean normal (58 sites) and reverse (13 sites) directions depart by only
307 6.7° from antipodal, which is less than the critical angle [*McFadden and McElhinny*, 1990],
308 indicating a positive reversal test.

309

310 The directional dispersion for the ChRM site means in the Shuttle Meadow, East Berlin
311 and Portland Formations hardly changes with corrections for bedding tilts because unlike the
312 much steeper A component, the shallow ChRM directions tend to be closer to coaxial with the
313 bedding strikes and thus relatively less sensitive to dip corrections. However, there is sufficient
314 variation in bedding attitudes with respect to available data from correlative sedimentary rock
315 units in the Newark basin – Feltville and Towaca Formations [*Witte and Kent*, 1990] – for an

316 inter-basin fold test. The mean ChRM direction for the 71 tilt-corrected sites from the Shuttle
317 Meadow, East Berlin and Portland Formations ($D = 6.8^\circ$, $I = 22.2^\circ$ $a95 = 3.7^\circ$; **Table 1**)
318 corresponds to a paleopole ($59.0^\circ\text{N } 94.5^\circ\text{E}$) for a nominal locality ($42^\circ\text{N } 72.5^\circ\text{W}$) in the Hartford
319 basin that differs by only an insignificant 3.7° from the paleopole ($55.3^\circ\text{N } 94.5^\circ\text{E } A95 = 5.4^\circ$)
320 for 11 tilt-corrected sites in the Feltville and Towaco Formations from the Newark basin
321 extrusive zone reported by [Witte and Kent, 1990]. In geographic coordinates, however, the
322 mean ChRM direction for the 71 sites from the Hartford basin ($D=2.6^\circ$ $I = 19.8^\circ$ $a95 = 3.6^\circ$;
323 **Table 1**) gives a paleopole ($58.1^\circ\text{N } 102.7^\circ\text{E}$) that differs by a significant 7.8° from the Feltville
324 and Towaco Formations ($56.1^\circ\text{N } 90.0^\circ\text{E } A95 = 5.7^\circ$) calculated without tilt correction [Witte and
325 Kent, 1990]. The inter-basin fold test is thus positive, which is also the case when ChRM
326 directions from only the more strictly equivalent units (Shuttle Meadow and East Berlin
327 Formations versus Feltville and Towaco Formations) are compared. The positive reversal test
328 and positive fold test indicate that the ChRM of the Shuttle Meadow, East Berlin and Portland
329 Formations was acquired early in the history of the rock units and is relatively uncontaminated
330 by secondary components.

331

332 Magnetostratigraphy

333

334 For diagnosing the geomagnetic polarity of the characteristic magnetizations, a virtual
335 geomagnetic pole (VGP) was calculated for each site direction and its latitude was compared to
336 the north paleopole ($59.0^\circ\text{N } 94.5^\circ\text{E}$) corresponding to the overall ChRM mean direction. VGP
337 latitudes close to $+90^\circ$ (or -90°) are interpreted as recording normal (or reverse) polarity of the
338 early Jurassic geomagnetic field and are plotted with respect to stratigraphic level to develop a
339 magnetostratigraphy.

340

341 Previous paleomagnetic work has established that the lava units in the Hartford basin –
342 Talcott, Holyoke and Hampden Basalts – erupted during normal geomagnetic polarity (e.g.,
343 *Prevot and McWilliams* [1989]). This conclusion can now be extended to the entire CAMP
344 extrusive zone of the Hartford basin because the interbedded sedimentary units – Shuttle
345 Meadow and East Berlin Formations – are also characterized by normal polarity (**Fig. 8**). We
346 cannot, of course, exclude the possibility of undetected short reverse polarity intervals but they

347 would have to be less than about 20 m thick, or constitute no more than a few percent of the total
348 stratigraphic thickness of the sedimentary units, at the present sampling density.

349
350 The normal polarity interval in the Hartford extrusive zone encompasses approximately
351 600 m of section and most probably corresponds to the normal polarity interval (Chron E24n) of
352 the homotaxial extrusive zone in the Newark basin, where the three lava units (Orange Mt.,
353 Preakness and Hook Mt. Basalts) as well as the interbedded sedimentary units (Feltville and
354 Towaco Formations), together about 1000 m thick, are also characterized by normal polarity
355 [McIntosh *et al.*, 1985; Prevot and McWilliams, 1989; Witte and Kent, 1990; Kent *et al.*, 1995;
356 Kent and Olsen, 1999] (**Fig. 8**). For convenience and to emphasize this correlation, we refer to
357 the Hartford extrusive zone normal polarity interval as magnetozone H24n. Suitable exposures of
358 the New Haven Formation immediately below the Talcott Basalt could not be found to establish
359 the expected presence of the thin reverse polarity interval corresponding to Chron E23r that
360 occurs in the uppermost Passaic Formation and within a few meters below the Orange Mt. Basalt
361 in the Newark basin [Kent *et al.*, 1995; Kent and Olsen, 1999].

362
363 The outcrop sites from approximately the lower 950 m of the Portland Formation
364 immediately overlying the youngest lavas (Hampden Basalt) are also characterized by normal
365 polarity that can be regarded as an extension of magnetozone H24n (**Fig. 8**). There are, however,
366 several sampling gaps of 100 m or more, mainly because of cover by sediments of Pleistocene
367 Lake Hitchcock in northern Connecticut and Massachusetts. We were able to fill several of these
368 sampling gaps by gaining access to a series of geotechnical cores taken by the Army Corps of
369 Engineers for the construction of the Park River drainage project in and near the city of Hartford
370 (**Fig. 2b**). The cores are presently stored at the Connecticut Department of Environmental
371 Protection Western District Headquarters in Harwinton, Connecticut. A composite section of
372 approximately 400 m thickness above the Hampden Basalt was assembled by projecting bedding
373 attitudes and tracing a series of distinctive beds (informally labeled A to I) in a transect of 14
374 selected drill cores 10 cm-diameter and up to 70 m long (**Fig. 9**). Sample plugs were drilled
375 perpendicular to the core axis (assumed to be vertical) and in the direction of bedding dip, which
376 was assumed to be in the regional direction of east-southeast (120°) and used for azimuthal
377 reorientation of the core segments. We analyzed 57 samples (up to 14 from each core) using the

378 same laboratory measurement and thermal demagnetization techniques as for the outcrop
379 samples. Well-defined, shallow ChRM directions were isolated from 600° to 675°C in 48 of the
380 samples although 9 of these samples had upward inclined A components, suggesting that these
381 particular core segments had been inadvertently turned upside down during handling. VGP
382 latitudes for the ChRM after azimuthal reorientation and correction for bedding tilt of the
383 accepted 39 samples show appreciable scatter (**Fig. 9**), which is not unexpected given the
384 uncertainties in reorienting the samples using core-bedding plane intersections with shallow
385 bedding dips. Nevertheless, the sample VGP latitudes are consistent with normal polarities, with
386 one possible exception: a sample from near marker Bed D, about 100 m above the Hampden
387 Basalt (core FD-12T sample depth 198'), had a negative albeit low VGP latitude, whose
388 significance as an indication of a thin interval of reverse polarity, a polarity excursion, or simply
389 noisy data is unclear and requires confirmation.

390

391 In the outcrop sites, the first unambiguous reverse-polarity magnetizations are
392 encountered near the top of the South Hadley Falls Member, approximately 950 m above the
393 Hampden Basalt, where four closely spaced sites at Cains Pond record high southerly VGP
394 latitudes (**Fig. 8**). These reverse polarity sites cover an interval less than 10 m thick although
395 available bounding constraints allow the reverse polarity interval (magnetozone H24r) to be as
396 much as ~100 m thick. This reverse polarity magnetozone has no counterpart in the Newark
397 basin, where the available section ends in normal polarity Chron E24n (=H24n).

398

399 Another reverse-polarity magnetozone (H25r) of perhaps comparable (~100 m) thickness
400 occurs above an intervening ~300 m interval of normal polarity of magnetozone H25n (**Fig. 8**).
401 Magnetozone H25r, which is in the uppermost Mittinegue Member, is delineated by only two
402 sites, which are, however, located more than 30 km apart along-strike in the Agawam and Stony
403 Brook sections. Within the middle and upper Stony Brook Member, a ~200 m interval of normal
404 polarity (magnetozone H26n) is overlain by the thickest (~200 m) reverse-polarity interval
405 (magnetozone H26r) thus far discovered in the Portland Formation, found in a total of 6 sites
406 from the Stony Brook and Agawam sections. The highest analyzed part of the Portland
407 Formation ends in normal polarity of magnetozone H27n, delineated by two sites in the Stony
408 Brook section about 1800 m above the Hampden Basalt.

409

410 Chronostratigraphic control

411

412 Chronostratigraphic control for the Hartford basin nonmarine stratigraphic succession has
413 traditionally been based on palynology, vertebrate biostratigraphy, and some inconclusive
414 geochronology. The pre-CAMP New Haven Formation is almost entirely fluvial; a Late Triassic
415 (latest Carnian to early Norian) palynoflorule was reported from the lower part of the unit
416 [Cornet, 1977], which also contains reptile fossils of Late Triassic age [Lucas *et al.*, 1998; Olsen
417 *et al.*, 2000]. A major advance was the recognition that a substantial portion of the Hartford
418 section extended into the Jurassic. Palynoflorules in the cyclical lacustrine strata of the Shuttle
419 Meadow Formation and the lower part of the Portland Formation are described as having Liassic
420 (Early Jurassic) affinities [Cornet *et al.*, 1973; Cornet and Traverse, 1975]. The uppermost few
421 meters of the New Haven Formation contains a palynoflorule of typical Early Jurassic aspect
422 [Heilman, 1987; Olsen *et al.*, 2002b], indicating that the Triassic-Jurassic boundary must lie just
423 below the base of the Talcott Basalt, in a homologous position with respect to the CAMP lavas in
424 the Newark basin [Olsen *et al.*, 2002a].

425

426 To date, the Hartford basin has yielded radioisotopic (K-Ar, $^{40}\text{Ar}/^{39}\text{Ar}$) dates that have
427 widely scattered values (150-250+ Ma) attributed to variable argon loss and excess argon
428 [Armstrong and Besancon, 1970; Seidemann *et al.*, 1984; Seidemann, 1988; 1989]. However,
429 paleomagnetic and geochemical data suggest a one-to-one correspondence of the volcanic units
430 in the Hartford basin to those in the Newark basin [Prevot and McWilliams, 1989] where
431 geochronological efforts have been much more successful. The Palisade sill, a traditional target
432 for radioisotopic dating, was probably a feeder for the lower extrusive unit (Orange Mountain
433 Basalt) in the Newark basin [Prevot and McWilliams, 1989]. A U-Pb baddeleyite date of 201 ± 1
434 Ma [Dunning and Hodych, 1990] and a $^{40}\text{Ar}/^{39}\text{Ar}$ biotite date of 202.2 ± 1.3 Ma from a
435 recrystallized sedimentary xenolith [Sutter, 1988] associated with the Palisade sill are consistent
436 with an $^{40}\text{Ar}/^{39}\text{Ar}$ age of 201.0 ± 1.2 Ma on the Orange Mountain Basalt [Hames *et al.*, 2000].
437 Together these dates are compatible with an age somewhere in the range ~200–202 Ma for the
438 Triassic-Jurassic boundary that lies a few meters below the equivalent of the Orange Mountain
439 Basalt in the Jacksonwald area of the Newark basin [Olsen *et al.*, 2002a]. In the Fundy basin of
440 Nova Scotia, the North Mountain Basalt is a few meters above the Triassic-Jurassic boundary

441 level [Fowell and Traverse, 1995; Olsen et al., 2003] and has yielded U-Pb zircon dates of
442 202 ± 1 Ma [Hodych and Dunning, 1992] and 201.27 ± 0.27 [Schoene et al., 2006] that strongly
443 support the geochronology from the Newark basin section. More recently, zircon-bearing tuffs in
444 association with ammonite-bearing strata in Peru yielded an U-Pb age of 201.58 ± 0.28 Ma for the
445 marine Triassic-Jurassic boundary [Schaltegger et al., 2007]. For consistency with our earlier
446 work, we use the rounded-off integer value of 202 Ma for the Triassic-Jurassic boundary that has
447 served as an anchor point in the astronomically-tuned GPTS for the Late Triassic based on
448 coring from the Newark basin [Kent and Olsen, 1999].

449
450 The cyclical lacustrine strata of the Shuttle Meadow, East Berlin and lower Portland
451 Formations provide an opportunity for establishing an astronomical time scale for the Jurassic
452 portion of the Hartford succession [Olsen and Kent, 2008]. The Hartford extrusive zone
453 encompasses about 600 kyr, partitioned about equally between the Shuttle Meadow and East
454 Berlin Formations and assuming that there is comparatively little time represented in the basalt
455 units [Whiteside et al., 2007] (**Fig. 10**). This duration is virtually identical to the cycle
456 stratigraphic estimate for the duration of the Newark extrusive zone (580 kyr; Olsen et al.,
457 1996b), which is also within the resolution of available $^{40}\text{Ar}/^{39}\text{Ar}$ dating of the lowest and highest
458 lava flows in the Newark basin [Hames et al., 2000]. The pervasive and distinctive cyclic
459 lithologic variation that reflects shallower and deeper lakes that we argue are tied to orbitally-
460 controlled precipitation/evaporation regimes continues into the lower Portland Formation. Key
461 features are facies changes at the 20 kyr precession cycle that are modulated by the short (100
462 kyr) and long (400 kyr) eccentricity cycles. Most usefully, the long eccentricity (sometimes
463 referred to as the McLaughlin) cycle often corresponds to mapped lithologic members (e.g., Park
464 River, S. Hadley Falls, and Mittinegue Members) and provides a basis for long-range correlation
465 and a chronological framework. This is similar to the pattern in the Newark basin section [Olsen,
466 1986; Olsen et al., 1996a] where a Triassic numerical time scale based on the fundamental
467 Milankovitch periodicities [Olsen and Kent, 1996; Kent and Olsen, 1999] has been largely
468 confirmed by high-precision geochronology [Wang et al., 1998; Furin et al., 2006]).

469
470 An astronomical time scale for the lower Portland Formation composite section indicates
471 an average sediment accumulation rate of ~ 1000 m/Myr (**Fig. 10**). For comparison, the cyclical

472 Late Triassic part of the Newark basin section accumulated at an average rate of only 160 m/Myr
473 [*Olsen et al.*, 1996a; *Kent and Olsen*, 1999] although the Jurassic sedimentary units that are
474 interbedded with the lavas can also have very high accumulation rates, for example, 26 m for the
475 20 kyr cycle, corresponding to 1300 m/Myr, for the Towaco Formation [*Olsen et al.*, 1996b].
476 Lithologic expressions of 20 kyr precession cyclicity are about 20 m thick in the Portland
477 Formation (somewhat thinner in the Shuttle Meadow and East Berlin Formations); the nominal
478 average paleomagnetic site sampling interval of roughly 30 m thus represents a temporal
479 resolution of one or two precession cycles. Even though the composite section for the Hartford
480 basin has considerable uncertainties in the depth scale because it was assembled from various
481 parts of the basin, only modest tuning (i.e., departures from sediment thickness as a first-order
482 linear proxy of time) was required to account for the cyclicity by Milankovitch climate forcing.
483 In all, the ~2500 m of section from the base of the Talcott Basalt (base of the CAMP extrusive
484 zone) into the lower to middle part of the Portland Formation represents ~6 McLaughlin cycles,
485 or ~2.4 Myr of Early Jurassic time (**Table 2**).

486

487 Geomagnetic polarity time scale

488

489 The Newark astronomically tuned GPTS was anchored to an age of 202 Ma for a level
490 corresponding to the end-Triassic extinction level identified on the basis of palynology and
491 supported by tetrapod footprint evidence [*Olsen et al.*, 1996a; *Kent and Olsen*, 1999; *Olsen et al.*,
492 2002b; *Olsen et al.*, 2003] that occurs about one precession cycle (~20 kyr) before the Orange
493 Mt. Basalt, the first CAMP lava in the Newark basin, and just after the end of Chron E23r, the
494 last geomagnetic polarity reversal in the Triassic [*Kent and Olsen*, 1999] (**Fig. 10**). Normal-
495 polarity Chron E24n, which begins at 202.021 Ma, encompasses the igneous extrusive zone in
496 the Newark basin and is correlative to the normal-polarity interval (Chron H24n) that
497 encompasses the CAMP extrusive zone in the Hartford basin and extends to the uppermost part
498 of the South Hadley Falls Member. This would make Chron H24n equal to nearly 4 McLaughlin
499 cycles, or 1590 kyr in total duration.

500

501 Chron H24r is the first reverse-polarity interval in the Jurassic as recorded in the Hartford
502 section; it is a thin (58 ± 50 m) magnetozone that occurs in the uppermost S. Hadley Falls

503 Member at about 200.45 Ma, or 1550 kyr after the Triassic-Jurassic boundary according to the
504 McLaughlin cyclicity. The overlying Mittinegue Member is mostly normal polarity (Chron
505 H25n) until another thin (51 ± 27 m) reverse polarity magnetozone (Chron H25r) occurs in its
506 youngest part at about 200.02 Ma. The succeeding Stony Brook Member has normal polarity
507 (Chron H26n) in its lower part and a relatively thick (154 ± 15 m) reverse-polarity interval
508 (Chron H26r) in its upper part. The sampled section ends in normal polarity of Chron H27n in
509 what may still be the Stony Brook Member where the cyclic facies character fades; the transition
510 from Chron H26r to Chron H27n occurs at an estimated age of 199.6 Ma and is the youngest
511 polarity reversal delineated thus far in the Portland Formation.

512

513 A plausible correlation can be made between the Hartford continental record and the
514 magnetobiostratigraphy of Hettangian and Sinemurian sediments in the Montcornet core from
515 the Paris basin [*Yang et al.*, 1996], the most detailed available marine record with
516 magnetostratigraphy for the Early Jurassic (**Fig. 11**). The ~30 m section (~1045–1075 m) of the
517 Montcornet core that corresponds to the Hettangian according to biostratigraphy is characterized
518 by predominantly normal polarity punctuated by several thin reverse polarity magnetozones,
519 whereas reverse polarity is more prevalent above about 1045 m in the late Hettangian to early
520 Sinemurian. The Montcornet polarity pattern thus suggests a correlation of Chrons H24n to H26n
521 to the predominantly normal polarities of the Hettangian, and Chron H26r, the longest and
522 youngest reverse polarity interval thus far delineated in the Early Jurassic of the Hartford basin
523 section, to the mostly reverse polarity interval (~1041–1045 m) in sediments designated as late
524 Hettangian to early Sinemurian in the Montcornet core. This general correlation implies that the
525 nearly 600 m thick CAMP extrusive zone of interbedded continental sediments and lavas and the
526 nearly 2000 m thick overlying section of continental sediments of the lower Portland Formation
527 all accumulated during the Hettangian. The ~2.4 Myr year duration estimated from cycle
528 stratigraphy for this part of the Hartford basin section is consistent with its correlation to the
529 Hettangian, which recent U-Pb dating indicates is only 2 to 3 Myr long [*Pálffy and Mundil*, 2006;
530 *Schaltegger et al.*, 2007].

531

532 The ~30 m Hettangian interval of the Montcornet core is also interpreted to have 5 very
533 thin reverse polarity intervals [*Yang et al.*, 1996]. One of the thin reverse intervals in the middle

534 Hettangian and another in the late Hettangian might correspond to the short subchrons H24r and
535 H25r, respectively, in the Hartford section (**Fig. 11**). However, no convincing counterparts have
536 yet been found in the Hartford (or Newark) basin section for any of the 3 thin reverse polarity
537 intervals in the early Hettangian part of the Montcornet core, which ostensibly should correspond
538 in age to the CAMP extrusive zone. Some of the thin polarity intervals in the Montcornet core
539 might be artifacts of inverted or overprinted core segments; alternatively, short polarity intervals
540 remain to be discovered in the CAMP extrusive zone of the Newark and Hartford basins.

541

542 Inclination shallowing

543

544 The mean inclination for the sedimentary units ($22.2\pm 3.7^\circ$ for 71 sites, or $21.1\pm 2.1^\circ$ for
545 315 samples from the Shuttle Meadow, East Berlin, and Portland Formations) is significantly
546 shallower than the mean inclination of $33.9\pm 8^\circ$ for the closely associated CAMP volcanic units
547 (calculated from the mean paleopole of $66.3^\circ\text{N } 97.3^\circ\text{E } A95=5^\circ$ for the three extrusive units in
548 the Newark and Hartford basins [*Prevot and McWilliams, 1989*]). This discrepancy may simply
549 be an artifact of unaveraged paleosecular variation in the CAMP volcanics or more likely is an
550 indication of sedimentary inclination error, which can occur during deposition of hematite-
551 bearing sediments [*Tauxe and Kent, 1984*]. A similar discrepancy was observed in the Newark
552 basin where the elongation/inclination (E/I) technique [*Tauxe and Kent, 2004*] was successfully
553 used to detect and correct inclination error in the characteristic directions from the sedimentary
554 units [*Kent and Tauxe, 2005*].

555

556 The E/I technique was applied to a dataset of 315 sample ChRM directions from the
557 Shuttle Meadow, East Berlin and Portland Formations, which represent 2.4 Myr of the earliest
558 Jurassic. This dataset should be sufficiently large in number of samples and length of record to
559 capture the full range of secular variation yet still short enough to avoid introducing a bias in the
560 directional distribution from polar wander. The distribution of ChRM directions is found to be
561 elongated east-west, *perpendicular to the paleomeridian*, that is, flattened in the paleohorizontal
562 or bedding plane (**Fig. 12a,b**). In contrast, statistical geomagnetic field models using a giant
563 Gaussian process (e.g., CP88.GAD: *Constable and Parker, 1988*; TK03.GAD: *Tauxe and Kent,*
564 2004) predict not only that the mean field inclination, I , is a function of latitude, λ :

565

566 $\tan I = 2 \tan \lambda$ (1)

567

568 but also that secular variation results in a distribution of virtual geomagnetic poles (longitudes
 569 and latitudes) that is essentially circular at any observation site, implying that the distribution of
 570 directions (declinations and inclinations) will be systematically more elongate north-south, *along*
 571 *the paleomeridian*, as the observation site latitude decreases from the pole(s) to the equator. If
 572 the directions were affected by inclination error (either during deposition or imparted by
 573 compaction), the observed inclination, I_o , will be related to the ambient field inclination, I_f , by:

574

575 $\tan I_o = f \tan I_f$ (2)

576

577 where f is the flattening factor (e.g., *King*, 1955). Inclination error affects the distribution of
 578 directions by increasing the east-west elongation while decreasing inclination. If inclination error
 579 is the cause of the shallow bias, the data can be inverted using the inverse of Equation 2
 580 searching for a value of f that yields an E/I combination that is consistent with the field model;
 581 the corrected mean inclination should provide a more accurate estimate of latitude according to
 582 Equation 1. The hypothesis that the statistical properties of the geomagnetic field in remote
 583 epochs were similar to the more recent (0–5 Ma) geomagnetic field was supported by the ability
 584 of the E/I method to produce an internally consistent latitudinal framework in the Late Triassic
 585 from studies made over a broad region [*Kent and Tauxe*, 2005].

586

587 E/I analysis of the Hartford sedimentary ChRM data produces a result consistent with the
 588 geomagnetic field model at a mean flattening factor of 0.54, which is well within the range of f
 589 values determined by E/I analysis of other datasets of similar hematite-bearing sedimentary rocks
 590 [*Kent and Tauxe*, 2005]; the corrected inclination is 35.5° , with bootstrapped 95% confidence
 591 limits of 32° and 39° (**Fig. 12c, d; Table 3**). The corrected inclination is more than 14° steeper
 592 than the uncorrected mean inclination, in keeping with what *Tan et al.* [2007] found with their
 593 magnetic anisotropy correction for the Passaic Formation red beds in the Newark basin. The
 594 corrected Hartford direction is also not significantly different from the mean inclination of
 595 $33.9 \pm 8^\circ$ estimated from the mean pole for the Newark and Hartford CAMP volcanic units

596 [Prevot and McWilliams, 1989], although the volcanics pole represents only a small number of
597 cooling units and thus may not adequately average paleosecular variation.

598

599 Paleomagnetic poles and J1 cusp

600

601 The (north) paleopole corresponding to the corrected Hartford sedimentary ChRM
602 direction is located at 66.6°N 88.2°E A95=2.3°; this is only an insignificant 3.6° away from the
603 Newark and Hartford CAMP volcanics pole at 66.3°N 97.3°E A95=5° [Prevot and McWilliams,
604 1989] and only an insignificant 3.3° from the paleopole for corrected results from latest Triassic
605 and earliest Jurassic sediments in the Martinsville core immediately below the CAMP extrusive
606 zone in the Newark basin (67.8°N 96.1°E A95=2.9°; Kent and Tauxe, 2005) (**Fig. 13**). A
607 comparison of paleopoles of tightly correlated Newark Supergroup strata of Late Triassic age
608 from the widely separated Newark basin and the Dan River basin of North Carolina and Virginia
609 shows no evidence of vertical-axis tectonic rotations [Kent and Olsen, 1997]. Similarly, the
610 positive fold test demonstrated here for data from Early Jurassic strata in the Newark and
611 Hartford basins is further indication that these early Mesozoic rift basins have maintained
612 tectonic coherence with respect to eastern North America. Accordingly, we regard the average of
613 the paleopoles for the CAMP volcanics, Newark corrected sediments and Hartford corrected
614 sediments (67.0°N 93.8°E A95=3.2°, N= 3) as representative of the ~201 Ma pole position for at
615 least eastern North America.

616

617 The paleopole from the Moenave Formation on the Colorado Plateau is usually
618 considered to practically define the J1 cusp (e.g., Molina-Garza *et al.*, 1995). A mean pole
619 position based on modern studies of the Moenave Formation at Vermillion Cliffs in Arizona and
620 Utah [Ekstrand and Butler, 1989] and at Echo Cliffs in Arizona [Molina-Garza *et al.*, 2003], and
621 of its presumed lateral equivalent, the Wingate Formation at Comb Ridge in Utah [Molina-Garza
622 *et al.*, 2003], is located at 59.3°N 59.0°E A95=8.3° (N=3), which is indeed very close to the
623 canonical location of the J1 cusp derived from the paleomagnetic Euler pole model of APW for
624 North America (60.5°N 62.4°E; Gordon *et al.*, 1984) (**Fig. 13**). Available biostratigraphic
625 evidence from palynoflora and tetrapod fossils and footprints suggest that the Moenave and
626 Wingate Formations of the Glen Canyon Group are latest Triassic (Rhaetian) to earliest Jurassic
627 (Hettangian) in age [Molina-Garza *et al.*, 2003]. This is virtually the same time interval

628 encompassed by latest Triassic sediments from the Martinsville core in the Newark basin,
629 earliest Jurassic sediments from the Hartford basin, and the CAMP extrusives in the Newark and
630 Hartford basins. However, the Moenave/Wingate mean pole differs by a significant 17.2° of
631 great circle arc from the coeval 201 Ma mean pole from the Newark/Hartford basins.

632

633 When projected to a nominal sampling location for the Moenave/Wingate on the
634 Colorado Plateau ($36.5^\circ\text{N } 111^\circ\text{W}$), the 201 Ma Newark/Hartford pole predicts a paleomagnetic
635 direction ($D=350.2^\circ I=28.5^\circ$) that is considerably more northwesterly but also much steeper than
636 the Moenave/Wingate direction ($D=5.1^\circ I=12.2^\circ$). The 14.9° difference in declination can be
637 attributed to net clockwise rotation of the Colorado Plateau relative to eastern North America by
638 a comparably large net amount (see summary of Plateau rotation estimates in *Steiner and Lucas*
639 [2000]). Clockwise Plateau rotation, however, cannot also account for the 16.3° shallower
640 direction in the Moenave/Wingate, which is more likely ascribed to inclination error. Flattening
641 factors in the range 0.4 to 0.6 have been found for characteristic magnetizations in other redbed
642 units (e.g., *Kent and Tauxe, 2005*) and if applicable to the Moenave/Wingate magnetizations, a
643 correction using a nominal value of $f\sim 0.5$ would steepen the Moenave/Wingate inclination to
644 23.4° and bring it into good agreement (within the error limits) with the predicted inclination
645 from the Newark/Hartford 201 Ma pole whereas a correction using $f=0.4$ would steepen the
646 Moenave/Wingate inclination to 28.4° and make it agree almost precisely with the predicted
647 inclination from the coeval Newark/Hartford pole.

648

649 The inferred degree of inclination error in the Moenave/Wingate magnetizations
650 obviously needs to be verified by E/I analysis [*Tauxe and Kent, 2004*] or by the anisotropy
651 method [*Tan et al., 2007*]. At this juncture, we find it intriguing that a correction of the
652 Moenave/Wingate mean direction for 13.5° clockwise rotation of the Colorado Plateau [*Kent and*
653 *Witte, 1993*] and for inclination error corresponding to a nominal flattening factor of $f\sim 0.5$ yields
654 a paleopole at $66^\circ\text{N } 86^\circ\text{E}$ that is in excellent agreement with the coeval Newark/Hartford 201 Ma
655 paleopole. The apparent concordance between the corrected mean Moenave/Wingate and coeval
656 Newark/Hartford 201 Ma poles and their appreciable departure from the postulated position of
657 the J1 cusp suggest that this key feature of paleomagnetic Euler pole analysis is largely an
658 artifact of Colorado Plateau rotation and sedimentary inclination error. Instead, the corrected data

659 show that APW for North America proceeds in a more northerly direction to higher latitudes
660 over the Late Triassic and Early Jurassic (**Fig. 13**).

661

662 Discussion

663

664 Chron E24n (=H24n) represents the thickest polarity unit in the Newark Supergroup
665 polarity sequence, encompassing about 1600 m of section in the Hartford basin and a minimum
666 of 1000 m in the Newark basin (**Fig. 8**). Chron E24n began just ~40 kyr prior to the earliest
667 CAMP lavas in the Newark basin [*Kent and Olsen, 1999*], extended over the 600 kyr-long
668 CAMP extrusive zone in both the Newark and Hartford basins [*Olsen et al., 1996b; Whiteside et*
669 *al., 2007*], and apparently ended with the first reverse polarity interval of the Jurassic (Chron
670 H24r) that occurred 950 kyr after the last CAMP lavas in the Hartford basin (**Fig. 10**).
671 Nevertheless, the 1590 kyr duration of Chron H24n is not the longest in the Late Triassic-earliest
672 Jurassic GPTS, being exceeded by three Triassic polarity chrons: 2003 kyr for E11r, 1797 kyr for
673 E16n, and 1618 kyr for E8r [*Kent and Olsen, 1999*]. Moreover, the first 6 polarity intervals of
674 the early Jurassic (H24n to H26r), which range in duration from ~50 kyr to 1590 kyr and
675 represent ~2400 kyr according to cycle stratigraphy (**Table 2**), have an average duration of
676 around 400 kyr, which is shorter than the average duration of 530 kyr for polarity intervals over
677 the preceding 30 Myr of the Late Triassic [*Kent and Olsen, 1999*]. These data do not support the
678 existence of a polarity superchron, or even a marked decrease in geomagnetic reversal frequency
679 in the Early Jurassic. The fact that 80% of the last 2.5 Myr of the Late Triassic and first ~ 2.5
680 Myr of the Early Jurassic had normal geomagnetic polarity is noteworthy but its significance is
681 unclear since there are several other 5 Myr intervals with a strong polarity bias in the Newark
682 GPTS, for example, the interval between 205 and 210 Ma has 80% reverse polarity [*Kent and*
683 *Olsen, 1999*].

684

685 Although the geomagnetic polarity column in the recent geologic time scale [*Gradstein et*
686 *al., 2004*] depicts the earliest Jurassic as having predominantly reverse polarity based on
687 preliminary data from Austria [*Steiner and Ogg, 1988*], latest Triassic and earliest Jurassic time
688 is in fact characterized by predominantly normal geomagnetic polarity based on published
689 magnetostratigraphies from marine sections – St. Audrie's Bay [*Hounslow et al., 2004*] and the

690 Montcornet core from the Paris basin [Yang *et al.*, 1996] – that are consistent with the Hartford
691 data. The St. Audrie’s Bay and especially the Montcornet magnetostratigraphic records are
692 punctuated by a number of relatively thin magnetic zones that have been interpreted as
693 representing short reverse polarity intervals [Yang *et al.*, 1996; Hounslow *et al.*, 2004]. The
694 shortest reverse polarity intervals in the Newark and Hartford polarity sequence are Chrons E23r
695 (~20 kyr) and H24r (at least 10 kyr but perhaps as long as 100 kyr; **Table 2**), which happen to
696 bracket the long normal polarity Chron E24n/H24n that includes CAMP volcanism. One or more
697 levels with anomalous paleomagnetic directions in the Moroccan record of CAMP rocks have
698 been correlated to Chron E23r, suggesting that CAMP volcanism started prior to the end of the
699 Triassic [Knight *et al.*, 2004; Marzoli *et al.*, 2004]. However, the anomalous directions from
700 Morocco taken at face value could just as well reflect one of the short reverse intervals or
701 polarity excursions in the Hettangian of the Montcornet core from the Paris basin [Yang *et al.*,
702 1996] that, if real, remain to be identified in the Newark and Hartford CAMP interval. Additional
703 short polarity intervals may exist in the Hartford (and Newark) basin sections although given the
704 present sampling density, any new polarity intervals would not be expected to be longer than
705 about 20 kyr. In the Cenozoic geomagnetic polarity record, such short features might qualify as
706 reversal excursions or polarity fluctuations, rather than full polarity chrons, and be difficult to
707 use for global correlation (e.g., Krijgsman and Kent, 2004; Lowrie and Kent, 2004). The isolated
708 sample with southerly VGP latitude about 100 m (or ~100 kyr) above the Hampden Basalt (**Fig.**
709 **9**) might be an example of such a reversal excursion that obviously needs to be verified.

710
711 The principal exemplars of the J1 cusp, the paleopoles from the Moenave Formation and
712 laterally correlative Wingate Formation from the Colorado Plateau, can be reconciled to the coeval
713 201 Ma reference paleopole for eastern North America (67.0°N 93.8°E A95=3.2°) based on data
714 from the Newark/Hartford CAMP lavas and sedimentary units corrected for inclination error by
715 net clockwise rotation of the Colorado Plateau and inclination error. The resulting APW path
716 follows a more northerly trend that effectively bypasses the prominent J1 cusp of the
717 paleomagnetic Euler pole model of North American APW [Gordon *et al.*, 1984; May and Butler,
718 1986]. The prevalence of inclination error indicated by comparisons to coeval igneous data [Gilder
719 *et al.*, 2003] and by E/I analysis [Krijgsman and Tauxe, 2004; Tauxe and Kent, 2004; Kent and
720 Tauxe, 2005] and the anisotropy technique [Tan and Kodama, 2002; Tan *et al.*, 2003, 2007] will

721 require a comprehensive re-evaluation of paleomagnetic data from sedimentary rocks used for
722 paleopole and paleolatitudinal studies, including attempts at using paleomagnetic Euler pole
723 analysis for estimating Colorado Plateau rotation [*Bryan and Gordon*, 1990]. In the meantime, we
724 suggest that the APW path for the eastern (stable) part of North America may best be delineated by
725 taking into consideration results from Jurassic igneous units from the White Mountain magma
726 series in New England, including a venerable result from the ~169 Ma Belknap Mountains and
727 Mount Monadnock plutons (85.4°N 354.6°E A95=3.5°; *Opdyke and Wensink* [1966], as
728 recalculated by *Van Fossen and Kent* [1990] and confirmed by them with thermal demagnetization
729 results giving a paleopole that includes the White Mountain batholith at 88.4°N 82.1°E A95=6.1°)
730 and the dual-polarity paleopole for the 169 Ma Moat Volcanics (81.6°N 89.7°E A95=6°; *Van*
731 *Fossen and Kent*, 1990), which would advance an APW path for the Mesozoic that is reminiscent
732 of the high latitude route proposed by *Irving and Irving* [1982]. Parenthetically, virtually all of the
733 other dozen or so listings of igneous results deemed reliable for the Jurassic of North America in a
734 recent paleopole compilation [*Besse and Courtillot*, 2002] have a wide cited age range, from the
735 180 Ma dikes in the Piedmont of North Carolina [*Smith*, 1987] to the 201 Ma Newark Supergroup
736 volcanics [*Prevot and McWilliams*, 1989], but more likely are poorly dated entries for the same
737 short lived (< 1 Myr) CAMP event at around 201 Ma. Even more recent radioisotopic ages for
738 CAMP igneous rocks are spread over a ~10 Myr interval [*Marzoli et al.*, 1999; *Knight et al.*,
739 2004]. Scatter in ages for CAMP (and indirectly, Chron E24n) may also help account for the
740 unverified long interval of low reversal frequency and normal polarity in the Early Jurassic that
741 appears in compilations of paleomagnetic polarity data (e.g., *Johnson et al.*, 1995; *Algeo*, 1996).

742
743 Finally, the new earliest Jurassic (201 Ma) reference paleopole based on the
744 Newark/Hartford data provides accurate and precise paleolatitudinal control, which is fully
745 consistent with Late Triassic data corrected for inclination error from North America and other
746 North Atlantic-bordering continents in a Pangea reconstruction [*Kent and Tauxe*, 2005]. By the
747 Early Jurassic, much of North America had drifted northward into the arid belt, in good
748 agreement with paleoclimate indicators of aridity such as eolian sandstones in the Pomperaug
749 basin of Connecticut (projected paleolatitude ~19°N) in eastern North America [*LeTourneau and*
750 *Huber*, 2006] and the appearance of prominent eolian sandstones in rock units of the Glen
751 Canyon Group (American Southwest) from the Wingate Formation (projected paleolatitude

752 ~15°N) that culminated in deposition of the Navajo Sandstone, one of the largest ergs on Earth
753 [Blakey *et al.*, 1988]. Any global climatic effects of CAMP will need to be evaluated in the
754 context of geographically distinct climate changes reflecting continental drift through latitudinal
755 climate belts.

756

757 Acknowledgements

758

759 We are grateful to the early important contributions of Bill Witte, who participated as a
760 graduate student in the fieldwork and laboratory analyses that initiated this paleomagnetic study
761 of the Hartford basin and for the preliminary data that were included in his PhD thesis at
762 Columbia University. We also thank Peter LeTourneou, Phil Huber, Jessica Whiteside and Luca
763 Lanci for field assistance on several of the many sampling trips to the Hartford basin, Randy
764 Steinen and Margaret Thomas for access to the Park River drainage project geotechnical cores,
765 Gilberto Mello and Cedric Mason for technical support in the paleomagnetic laboratory, Lisa
766 Tauxe for providing an E/I analysis of the data shown in Figure 12, and the JGR editors and the
767 referees (Ken Kodama and Laurie Brown) for constructive reviews. This research was supported
768 by the Earth Sciences Division of the National Science Foundation under several grants, most
769 recently EAR 0446843. Lamont-Doherty Earth Observatory contribution #7122.

770 Table 1. Paleomagnetic directions isolated from Early Jurassic sedimentary rock units from the
 771 Hartford basin.

773 774 775 776 777	Rock unit	N	Geographic				Tilt corrected			
			Dec	Inc	k	a95	Dec	Inc	k	a95
778	A component (100-300°C)									
779 780	SM+EB+PF	78	353.9°	60.2°	101	1.6°	13.4°	62.3°	39	2.6°
781	ChRM (600-675°C)									
782 783	SM	8	356.4	20.4	60	7.2	1.2	20.4	64	7.0
784	EB	5	358.3	21.6	63	9.7	7.7	28.8	41	12.2
785	SM+EB	13	357.1	20.9	66	5.1	3.6	23.6	47	6.1
786	PF									
787	normal	45	5.4	20.2	25	4.3	9.2	22.5	23	4.5
788	reverse	13	178.5	-17.1	12	12.6	181.6	-19.6	12	12.5
789	all	58	3.8	19.6	20	4.3	7.5	21.9	19	4.4
790										
791										
792	SM+EB+PF									
793	normal	58	3.5	20.4	28	3.6	7.9	22.7	26	3.7
794	reverse	13	178.5	-17.1	12	12.6	181.6	-19.6	12	12.5
795	all	71	2.6	19.8	23	3.6	6.8	22.2	21	3.7
796										

797 Rock units are SM, Shuttle Meadow; EB, East Berlin; PF, Portland Formation. N is number of
 798 sites, Dec and Inc are the declination and inclination, k is the best estimate of Fisher's precision
 799 parameter, and a95 is the radius of 95% circle of confidence for the mean direction.

800 Table 2. Astronomically tuned geomagnetic polarity time scale for ~ 5
 801 Myr interval centered on the Triassic-Jurassic boundary set at 202 Ma
 802 based on magnetic and cycle stratigraphy data from the Hartford and
 803 Newark basins.

805	806 Base of unit	807 Depth (m)	808 McLaughlin cycle	809 Age (Ma)
809	H27n	2386.95	66.814	199.687
810	H26r	2232.50	66.443	199.837
811	Stony Brook Mb.	2047.70	66	200.016
812	H26n	2008.60	65.918	200.049
813	H25r	1957.55	65.812	200.092
814	H25n	1578.10	65.019	200.412
815	Mittinegue Mb.	1568.79	65	200.420
816	H24r	1520.00	64.870	200.473
817	S. Hadley Falls Mb.	1194.39	64	200.824
818	Park River Mb.	713.12	63	201.228
819	Smiths Ferry Mb.	379.64	62	201.632
820	Talcott (=Orange Mt) Basalt	0	61.141	201.979
821	Tr/J (Newark basin)	-5.20	61.089	202
822	E24n=H24n	-10.70	61.036	202.021
823	VV (Exeter Mb.)	-12.3	61	202.036
824	E23r	-13.81	60.970	202.048
825	UU (Pine Ridge Mb.)	-59.50	60	202.440
826	TT	-124.7	59	202.844
827	E23n	-152.40	58.550	203.026
828	SS	-186.50	58	203.248
829	E22r	-197.66	57.796	203.330
830	E22n.2n	-230.95	57.215	203.565
831	E22n.1r	-232.56	57.188	203.576
832	RR	-241.30	57	203.652
833	E22n.1n	-288.37	56.284	203.941
834	QQ	-307.00	56	204.056
835	E21r.3r	-333.30	55.506	204.256
836	E21r.2n	-336.13	55.452	204.277
837	E21r.2r	-353.60	55.123	204.410
838	E21r.1n	-359.91	55.004	204.458
839	PP	-360.10	55	204.460

840
 841 Magnetic polarity chrons defined in Newark Basin Coring Project cores
 842 have the prefix E [*Kent and Olsen, 1999*] and those from Hartford basin
 843 outcrop composite section have prefix H (this study); polarity is
 844 designated by suffix n for normal and r for reverse. The base of each chron
 845 is given as the fractional position from the base of the enclosing

846 McLaughlin member cycle, counted up from RaR-8 (informal cycle 1) in
847 Stockton Formation to the Exeter Member (informal cycle 61) in the
848 uppermost Passaic Formation in the Newark basin and continuing to Stony
849 Brook Member (informal cycle 66) of Portland Formation in the Hartford
850 basin. Depth in the Newark basin (negative numbers) is composite
851 stratigraphic thickness scaled downward from base of Orange Mountain
852 Basalt and normalized to Rutgers drill core based on successive core
853 overlap correlations [*Olsen et al.*, 1996a]; depth in the Hartford basin
854 (positive numbers) is measured upward from base of Talcott Basalt (this
855 study; *Olsen and Kent*, 2008]). Ages for the polarity chrons are based on
856 interpolation within McLaughlin cycles, which are assumed to represent
857 the 404-kyr orbital eccentricity modulation of climate precession. The
858 relative chronology is indexed to an estimated age of 202 Ma for the
859 Triassic-Jurassic (Tr/J) boundary event. The depths of unit boundaries,
860 interpolated values of position within a cycle, and ages are quoted with a
861 precision needed for internal consistency.

862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881

Table 3. Results of E/I analysis on ChRM sample directions from the Early Jurassic Shuttle Meadow, East Berlin and lower Portland Formations from the Hartford basin.

Locality	Slat (°N)	Slon (°E)	Age (Ma)	n	Dec (°)	Inc (°)	λ (°N)	f	Inc' (°)	\pm Inc' (°)	λ' (°N)	$\pm \lambda'$ (°)
Hartford basin	42.0	-72.5	201	315	8.0	21.1	10.9	0.54	35.5	32–39	19.2	17.4–22.0

Slat and Slon are the latitude and longitude of the nominal sampling locality; Age is the nominal mean age of the early Jurassic rock units; n is the number of data analyzed; Dec and Inc are the mean declination and mean inclination of the characteristic magnetization data; λ is paleolatitude calculated with dipole formula from the mean inclination; f is the flattening factor determined from E/I analysis; Inc' is the corrected mean inclination and \pm Inc' is its 95% confidence interval; λ' is the corresponding corrected paleolatitude and $\pm \lambda'$ is its 95% confidence interval.

882 Table 4. Paleomagnetic poles from selected Late Triassic and earliest
 883 Jurassic rocks from North America.

886	Locality	Age	<i>f</i>	Plat'	Plon'	A95	Ref.
887		(Ma)		(°N)	(°E)	(°)	
888	<i>Sedimentary results corrected for inclination error</i>						
889	<i>Sedimentary results corrected for inclination error</i>						
890	<i>Sedimentary results corrected for inclination error</i>						
891	Hartford basin						
892	SM+EB+PF (H)	201	0.54	66.6	88.2	2.3	1
893	Newark Basin Coring Project						
894	Martinsville (M)	204	0.66	67.8	96.1	2.9	2,3
895	Weston Canal (W)	207	0.49	66.6	86.5	2.9	2,3
896	Somerset (S)	211	0.49	61.7	95.3	2.0	2,3
897	Rutgers (R)	214	0.66	60.1	97.1	1.4	2,3
898	Titusville (T)	217	0.63	59.9	99.5	1.7	2,3
899	Nursery Road (N)	221	0.40	60.5	101.6	2.5	2,3
900	Princeton (P)	227	0.56	54.2	106.6	2.0	2,3
901	Dan River basin (D)	221	0.59	58.5	99.8	1.1	3,4
902							
903	Colorado Plateau						
904	Moenave+Wingate (mo+wi)	201	1.0*	59.3	59.0	8.3	5,6
905	Moenave+Wingate	201	0.5**	65.3	57.0	8.3	1,5,6
906	plus 13.5° cw rotation (Mo+Wi)	201	0.5**	66.3	85.9	8.3	1,5-7
907							
908	<i>Igneous results</i>						
909							
910	Newark+Hartford lavas (CAMP)	201		66.3	97.3	5.0	8,9
911	Manicouagan (MI)	214		58.8	89.9	5.8	10-12
912							

913 Age is the nominal age of the Late Triassic and early Jurassic rock units; *f* is the
 914 flattening factor determined from E/I analysis (* uncorrected data, ** corrected
 915 data with an assumed flattening factor for Moenave+Wingate); Plat' and Plon'
 916 are the latitude and longitude of the paleopole that corresponds to corrected
 917 directions for the sedimentary results (see also Table 3). Ref. is the literature
 918 source for the age and paleomagnetic data: 1. This study; 2. Kent et al. (1995); 3.
 919 Kent & Tauxe (2005); 4. Kent & Olsen (1997); 5. Ekstrand & Butler 1989; 6.
 920 Molina-Garza et al. (2003); 7. Kent & Witte (1993); 8. Prevot & McWilliams
 921 (1989); 9. Hames et al. (2000); 10. Robertson (1967); 11. Larochelle & Currie
 922 (1967); 12. Hodych & Dunning (1992). 201 Ma reference pole position for stable
 923 North America (Figure 13) is the mean of corrected Hartford basin sediments
 924 (H), corrected Martinsville core sediments (M) and Newark+Hartford lavas
 925 (CAMP) at 67.0°N 93.8°E (A95=3.2°).

926 FIGURE CAPTIONS

927

928 Figure 1. Paleogeographic extent of ~200 Ma Central Atlantic magmatic province (CAMP)
 929 across the central Pangean supercontinent (after *McHone* [2000] and *Whiteside et al.*
 930 [2007]). From south to north: BP, Blake Plateau; C, Culpeper basin; N, Newark basin; H,
 931 Hartford basin; D, Deerfield basin; A, Argana basin; CHA, Central High Atlas basin; F,
 932 Fundy basin.

933

934 Figure 2. a) Early Mesozoic rift basins in eastern North America: 1. Fundy; 2. Hartford; 3.
 935 Newark; 4. Gettysburg, 5. Culpeper; 6. Danville. b) Geologic sketch map of Hartford
 936 basin with sampling sites as open (reverse polarity) and filled (normal and indeterminate
 937 polarity) circles. Army Corps of Engineers Park River drainage project geotechnical
 938 cores are indicated by a series of filled circles with crosses next to label 'PR'. C)
 939 Stratigraphic section of Newark Supergroup in Hartford basin with ties along right
 940 margin of column showing paleomagnetic sampling levels in Shuttle Meadow, East
 941 Berlin and lower Portland Formations.

942

943 Figure 3. Representative vector end-point diagrams (open/filled circles are projections on
 944 vertical/horizontal axes in geographic coordinates with temperatures in centigrade
 945 adjacent to selected points) of thermal demagnetization of NRM of samples from
 946 Portland (a-d, g, h), Shuttle Meadow (e) and East Berlin (f) Formations. These samples
 947 were selected from magnetozones H24n (b, e, f), H24r (d), H25n (h), H25r (c), H26n (a),
 948 and H26r (g) (see Figure 8 for magnetostratigraphy). Insets show changes in magnetic
 949 susceptibility (k/k_0) normalized to peak value after each heating step as a monitor of
 950 magnetochemical alteration during thermal demagnetization.

951

952 Figure 4. IRM acquisition (a) and thermal demagnetization of orthogonal IRM components of
 953 lower (<200 mT or <300 mT) and higher (>200 mT or >300 mT) coercivity (b, c, d, e, f)
 954 with attendant changes in magnetic susceptibility (k) of representative sedimentary rock
 955 samples from the Hartford basin. In this method of IRM acquisition (a), remanent
 956 coercivity is the intersection of the acquisition curve and null IRM. Samples come from

957 red beds (b: East Berlin Formation; c, d: Portland Formation) or from gray shales (e, f:
 958 Portland Formation) and represent sites with either normal polarity (b, d, f) or reverse
 959 polarity (c, e) characteristic directions (see NRM thermal demagnetization data from
 960 some companion specimens in Figure 3).

961

962 Figure 5. Comparison of IRM acquisition (left panels) before and after the samples were
 963 thermally demagnetized to 685°C; in this method, remanent coercivity is the intersection
 964 of the acquisition curve and null normalized IRM. Thermal demagnetization of
 965 orthogonal IRM given after heating to 685°C (right panels) can be compared to initial
 966 thermal demagnetization of IRM of the same samples in Figure 4. Samples are from (a,
 967 b) gray shale from Portland Formation, (c, d) red mudstone from East Berlin Formation,
 968 and (e, f) red siltstone from Portland Formation.

969

970 Figure 6. A-component site-mean directions (filled circles) from Shuttle Meadow, East Berlin,
 971 and lower Portland Formations, before (geographic) and after (bedding) correction for
 972 tectonic bedding tilts, plotted on lower hemisphere of equal area plots; star is geocentric
 973 axial dipole field direction and square is present-day direction. The decrease in precision
 974 parameter (k) after tilt correction is significant and indicates a negative fold test.

975

976 Figure 7. Characteristic (ChRM) site-mean directions from Shuttle Meadow and East Berlin
 977 Formations (triangles) and from lower Portland Formation (circles). Open/filled symbols
 978 are plotted on upper/lower hemisphere of equal area plots before (left, geographic) and
 979 after (right, bedding) correction for tectonic bedding tilts.

980

981 Figure 8. Magnetostratigraphy and lithostratigraphy of Early Jurassic strata in the Hartford basin.
 982 Magnetic polarity chrons are identified next to the polarity column where filled and open
 983 bars denote normal and reverse polarity interpreted from VGP latitudes (values
 984 approaching +90° indicate normal polarity and values approaching -90° indicate reverse
 985 polarity) with respect to overall mean paleopole for outcrop sites (open circles), sediment
 986 samples from Park River cores (small filled circles), and from lava units (bars; *Prevot*
 987 *and McWilliams*, 1989). Letters adjacent to lithology column correspond to key beds used

988 for correlation of Park River cores (see Figure 9). Correlative section from Newark basin
989 is shown at right (lithology after *Olsen et al.* [1996b], polarity column from *Kent and*
990 *Olsen* [1999]) with numerical ages for Orange Mt. and Hook Mt. Basalts shown in
991 lithostratigraphy panel from *Hames et al.* [2000].
992

993 Figure 9. Composite section of the mostly covered interval of Smiths Ferry and Park River
994 Members of lower Portland Formation based on Park River cores taken by Army Corps
995 of Engineers. Projection of bedding dips and tracing of distinctive beds labeled A to I
996 were used to arrange the 14 cores in stratigraphic sequence. VGP latitudes for
997 characteristic magnetizations of indirectly oriented samples were used to interpret
998 polarity (values approaching +90° indicate normal polarity and values approaching -90°
999 indicate reverse polarity; crosses along 0° axis indicate samples that did not provide
1000 interpretable data). See Figure 8 for integration with data from outcrop sections.
1001

1002 Figure 10. Astronomically tuned geomagnetic polarity time scale (GPTS) for the early Jurassic
1003 based on cycle and magnetic polarity stratigraphy of composite section from Hartford
1004 basin. Conversion of stratigraphic thickness to age is based on interpolation within
1005 McLaughlin cycles, which are assumed to represent the 404-kyr orbital eccentricity
1006 variation (modeled precession envelope described by *Whiteside et al.* [2007]), and
1007 indexed to an estimated age of 202 Ma for the Triassic-Jurassic (Tr/J) boundary event.
1008 See Table 2.
1009

1010 Figure 11. Magnetobiostratigraphy of Hettangian and Sinemurian marine sediments from the
1011 lower part of Montcornet core from the Paris basin plotted on a linear depth scale [*Yang*
1012 *et al.*, 1996] and compared to the astronomically-tuned geomagnetic polarity time scale
1013 (GPTS) and lithology column for the early Jurassic from the Hartford basin that was
1014 scaled in time using cycle stratigraphy and a Triassic-Jurassic boundary age of 202 Ma.
1015 Alignments of the Rhaetian-Hettangian (=Triassic-Jurassic) boundary level and the
1016 prominent reverse polarity chron H26r with an interval of predominantly reverse polarity
1017 between ~1041–1045 m in the Montcornet core would suggest that the Hettangian is only
1018 a few million years long. A short duration for the Hettangian is supported by U-Pb dates

1019 from early Sinemurian and Hettangian marine sediments with biostratigraphic control
 1020 [*Pálffy and Mundil, 2006; Schaltegger et al., 2007*].

1021
 1022 Figure 12. Equal area projections of (a) sample ChRM directions after bedding tilt correction
 1023 from East Berlin, Shuttle Meadow, and Portland Formations, and (b) the same
 1024 distribution rotated so that the overall mean direction (Table 1) corresponds to the
 1025 vertical axis to view better the shape of the distribution, which is elongated perpendicular
 1026 to the paleomeridian and indicative of inclination flattening. (c) E/I analysis [*Tauxe and*
 1027 *Kent, 2004*] of the sample ChRM directions from East Berlin, Shuttle Meadow, and lower
 1028 Portland Formations with the trajectory of mean inclination versus elongation of the
 1029 distribution calculated as the data are inverted with values for the flattening factor (f)
 1030 ranging from 0.3 to 1.0. The predicted E/I trend of the TK03.GAD geomagnetic field
 1031 model is shown as dashed line; the E/I of the data consistent with the model is circled and
 1032 corresponds to $f=0.54$. (d) Histogram of 1000 intersections of the kind shown in (c) from
 1033 bootstrapped curves. The mean and 95% confidence bounds of the corrected inclination
 1034 are shown (see **Table 3**).

1035
 1036 Figure 13. Selected Late Triassic and early Jurassic paleopoles from North America. Large
 1037 circles are A95s for igneous poles (CAMP lavas; MI, Manicouagan impact structure) and
 1038 poles corrected for inclination error: P, Princeton; N, Nursery; T, Titusville; R, Rutgers;
 1039 S, Somerset; W, Weston Canal; and M, Martinsville for NBCP cores; D for Dan River
 1040 basin, H for Hartford basin (see **Table 4**), whereas small filled circles with lower case
 1041 letters are paleopoles for same sedimentary units before correction for inclination error.
 1042 Mean paleopole for Moenave and Wingate Formations is shown with circle of confidence
 1043 labeled mo+wi, and as shaded circle of confidence labeled Mo+Wi after correction for
 1044 inclination error (assumed flattening factor, $f=0.5$) and 13.5° net clockwise rotation of
 1045 Colorado Plateau with respect to eastern North America (**Table 4**). Star with circle of
 1046 confidence labeled '201 Ma' is mean paleopole of CAMP lavas and corrected Martinsville
 1047 core (M) and Hartford basin (H) sedimentary directions. J1 is calculated position of the
 1048 ~ 203 Ma cusp joining Permian-Triassic and Jurassic-Cretaceous tracks from a
 1049 paleomagnetic Euler pole model (path shown by line with arrow labeled PEP; *Gordon et*

1050 *al.*, 1984) that now is seen to be biased by inclination error and rotation of the Colorado
1051 Plateau. Heavier line with arrow of time is our preferred empirical APW path for North
1052 America for the Late Triassic and Early Jurassic based on igneous results and
1053 sedimentary data corrected for inclination error from eastern North America.

1054 REFERENCES

1055

- 1056 Algeo, T. J. (1996), Geomagnetic polarity bias patterns through the Phanerozoic, *Journal of*
 1057 *Geophysical Research*, 101, 2785-2814.
- 1058 Armstrong, R. L., and J. Besancon (1970), A Triassic time scale dilemma: K-Ar dating of Upper
 1059 Triassic mafic igneous rocks, eastern U.S.A. and Canada and post-Upper Triassic plutons,
 1060 western Idaho, U.S.A., *Eclogae Geologiae Helveticae*, 63, 15-28.
- 1061 Besse, J., and V. Courtillot (2002), Apparent and true polar wander and the geometry of the
 1062 geomagnetic field over the last 200 Myr, *Journal of Geophysical Research*, 107(B11), EPM
 1063 6/1-6/31, doi:10.1029/2000JB000050.
- 1064 Blakey, R. C., F. Peterson, and G. Kocurek (1988), Synthesis of late Paleozoic and Mesozoic
 1065 eolian deposits of the Western Interior of the United States, *Sedimentary Geology*, 56, 3-125.
- 1066 Bryan, P., and R. G. Gordon (1990), Rotation of the Colorado Plateau: An updated analysis of
 1067 paleomagnetic poles, *Geophysical research Letters*, 17(Geophys. Res. Lett.), 1501-1504.
- 1068 Constable, C. G., and R. L. Parker (1988), Statistics of the geomagnetic secular variation for the
 1069 past 5 m.y., *Journal of Geophysical Research*, 93, 11,569-511,581.
- 1070 Cornet, B., A. Traverse, and N. G. McDonald (1973), Fossil spores, pollen, and fishes from
 1071 Connecticut indicate Early Jurassic age for part of the Newark Group, *Science*, 182, 1243-
 1072 1247.
- 1073 Cornet, B., and A. Traverse (1975), Palynological contributions to the chronology and
 1074 stratigraphy of the Hartford Basin in Connecticut and Massachusetts, *Geoscience and Man*,
 1075 11, 1-33.
- 1076 Cornet, B. (1977), The palynostratigraphy and age of the Newark Supergroup, PhD thesis,
 1077 Pennsylvania State University.
- 1078 Cornet, B., and P. E. Olsen (1985), A summary of the biostratigraphy of the Newark Supergroup
 1079 of eastern North America with comments on Early Mesozoic provinciality, paper presented at
 1080 Simposio Sobre Floras del Triasico Tardio, su Fitogeografia y Paleoecologia: Memoria, III
 1081 Congreso Latinoamericano de Paleontologia, Mexico.
- 1082 Davis, W. M. (1898), The Triassic formation of Connecticut, U.S. Geological Survey Annual
 1083 Report, 18(2), 1-192.
- 1084 De Boer, J. (1968), Paleomagnetic differentiation and correlation of the Late Triassic volcanic
 1085 rocks in the central Appalachians (with special reference to the Connecticut Valley),
 1086 *Geological Society of America Bulletin*, 79(Geol. Soc. America Bull.), 609-626.
- 1087 DuBois, P. M. (1957), Comparison of palaeomagnetic results for selected rocks of Great Britain
 1088 and North America, *Advances in Physics*, 6, 177-186.
- 1089 DuBois, P. M., E. Irving, N. D. Opdyke, S. K. Runcorn, and M. R. Banks (1957), The
 1090 geomagnetic field in Upper Triassic times in the United States, *Nature*, 180, 1186-1187.
- 1091 Dunning, G. R., and J. P. Hodych (1990), U/Pb zircon and baddeleyite ages for the Palisades and
 1092 Gettysburg sills of the northeastern United States: Implications for the age of the
 1093 Triassic/Jurassic boundary, *Geology*, 18, 795-798.
- 1094 Ekstrand, E. J., and R. F. Butler (1989), Paleomagnetism of the Moenave Formation:
 1095 Implications for the Mesozoic North American apparent polar wander path, *Geology*, 17, 245-
 1096 248.
- 1097 Fowell, S. J., and A. Traverse (1995), Palynology and age of the upper Blomidon Formation,
 1098 Fundy basin, Nova Scotia, *Review of Palaeobotany and Palynology*, 86, 211-233.

- 1099 Furin, S., N. Preto, M. Rigo, G. Roghi, P. Gianolla, J. L. Crowley, and S. A. Bowring (2006),
 1100 High-precision U-Pb zircon age from the Triassic of Italy: Implications for the Triassic time
 1101 scale and the Carnian origin of calcareous nannoplankton and dinosaurs, *Geology*, 34(12),
 1102 1009-1012.
- 1103 Gallet, Y., J. Besse, L. Krystyn, J. Marcoux, and H. Theveniaut (1992), Magnetostratigraphy of
 1104 the Late Triassic Bolucektasi Tepe section (southwestern Turkey): Implications for changes in
 1105 magnetic reversal frequency, *Physics of the Earth and Planetary Interiors*, 73, 85-108.
- 1106 Gilder, S., Y. Che, J. Cogné, X. Tan, V. Courtillot, D. Sun, and Y. Li (2003), Paleomagnetism of
 1107 Upper Jurassic to Lower Cretaceous volcanic and sedimentary rocks from the western Tarim
 1108 Basin and implications for inclination shallowing and absolute dating of the M-O (ISEA?)
 1109 chron, *Earth and Planetary Science Letters*, 206, 587-600.
- 1110 Gordon, R. G., A. Cox, and S. O'Hare (1984), Paleomagnetic Euler poles and the apparent polar
 1111 wander and absolute motion of North America since the Carboniferous, *Tectonics*, 3, 499-
 1112 537.
- 1113 Gradstein, F. M., J. G. Ogg, and A. G. Smith (Eds.) (2004), *A Geologic Time Scale 2004*, 589
 1114 pp., Cambridge University Press, Cambridge.
- 1115 Hames, W. E., P. R. Renne, and C. Ruppel (2000), New evidence for geologically-instantaneous
 1116 emplacement of earliest Jurassic Central Atlantic magmatic province basalts on the North
 1117 American margin, *Geology*, 28, 859-862.
- 1118 Heilman, J. J. (1987), The catastrophic day in the Early Jurassic, *Journal of Science Education*,
 1119 25, 8-25.
- 1120 Hodych, J. P., and G. R. Dunning (1992), Did the Manicouagan impact trigger end-of-Triassic
 1121 mass extinction?, *Geology*, 20(Geology), 51-54.
- 1122 Hounslow, M. W., P. E. Posen, and G. Warrington (2004), Magnetostratigraphy and
 1123 biostratigraphy of the Upper Triassic and lowermost Jurassic succession, St. Audrie's Bay,
 1124 UK, *Palaeoceanography, Palaeoclimatology, Palaeoecology*, 213, 331-358.
- 1125 Hozik, M. J. (1992), Paleomagnetism of igneous rocks in the Culpeper, Newark, and
 1126 Hartford/Deerfield basins, in *Eastern North American Mesozoic magmatism*, edited by J. H.
 1127 Puffer and P. C. Ragland, pp. 279-308, Geological Society of America, Boulder.
- 1128 Hubert, J. F., P. E. Feshbach-Meriney, and M. A. Smith (1992), The Triassic-Jurassic Hartford
 1129 Rift Basin, Connecticut and Massachusetts: Evolution, sandstone diagenesis, and hydrocarbon
 1130 history, *American Association of Petroleum Geologists Bulletin*, 76, 1710-1734.
- 1131 Irving, E., and M. R. Banks (1961), Paleomagnetic results from the Upper Triassic lavas of
 1132 Massachusetts, *Journal of Geophysical Research*, 66, 1935-1939.
- 1133 Irving, E., and G. Pullaiah (1976), Reversals of the geomagnetic field, magnetostratigraphy, and
 1134 relative magnitude of paleosecular variation in the Phanerozoic, *Earth Science Reviews*, 12,
 1135 35-64.
- 1136 Irving, E., and G. A. Irving (1982), Apparent polar wander paths Carboniferous through
 1137 Cenozoic and the assembly of Gondwana, *Geophysical Surveys*, 5, 141-188.
- 1138 Johnson, H. P., D. Van Patten, M. A. Tivey, and W. W. Sager (1995), Geomagnetic polarity
 1139 reversal rate for the Phanerozoic, *Geophysical research Letters*, 22, 231-234.
- 1140 Kent, D. V., and W. K. Witte (1993), Slow apparent polar wander for North America in the Late
 1141 Triassic and large Colorado Plateau Rotation, *Tectonics*, 12, 291-300.
- 1142 Kent, D. V., P. E. Olsen, and W. K. Witte (1995), Late Triassic-earliest Jurassic geomagnetic
 1143 polarity sequence and paleolatitudes from drill cores in the Newark rift basin, eastern North
 1144 America, *Journal of Geophysical Research*, 100, 14,965-914,998.

- 1145 Kent, D. V., and P. E. Olsen (1997), Paleomagnetism of Upper Triassic continental sedimentary
1146 rocks from the Dan River-Danville rift basin (eastern North America), *Geological Society of*
1147 *America Bulletin*, 109, 366-377.
- 1148 Kent, D. V., and P. E. Olsen (1999), Astronomically tuned geomagnetic polarity time scale for
1149 the Late Triassic, *Journal of Geophysical Research*, 104, 12,831-812,841.
- 1150 Kent, D. V., and L. Tauxe (2005), Corrected Late Triassic latitudes for continents adjacent to the
1151 North Atlantic, *Science*, 307, 240–244.
- 1152 King, R. F. (1955), The remanent magnetism of artificially deposited sediment, *Monthly Notices*
1153 *Royal Astronomical Society, Geophysics Supplement*, 7, 115-134.
- 1154 Kirschvink, J. L. (1980), The least-squares line and plane and the analysis of palaeomagnetic
1155 data, *Geophysical Journal of the Royal astronomical Society*, 62, 699-718.
- 1156 Knight, K. B., S. Nomade, P. R. Renne, A. Marzoli, H. Bertrand, and N. Youbi (2004), The
1157 Central Atlantic Magmatic Province at the Triassic-Jurassic boundary: paleomagnetic and
1158 $^{40}\text{Ar}/^{39}\text{Ar}$ evidence from Morocco for brief, episodic volcanism, *Earth and Planetary Science*
1159 *Letters*, 228, 143–160.
- 1160 Kodama, K. P., M. T. Cioppa, E. Sherwood, and A. C. Warnock (1994), Paleomagnetism of
1161 baked sedimentary rocks in the Newark and Culpeper basins: Evidence for the J1 cusp and
1162 significant Late Triassic apparent polar wander from the Mesozoic basins of North America,
1163 *Tectonics*, 13, 917-928.
- 1164 Krijgsman, W., and D. V. Kent (2004), Non-uniform occurrence of short-term polarity
1165 fluctuations in the geomagnetic field? New results from Middle to Late Miocene sediments of
1166 the North Atlantic (DSDP Site 608), in *Timescales of the Paleomagnetic Field*, edited by J. E.
1167 T. Channell, D. V. Kent, W. Lowrie and J. Meert, pp. 161–174, American Geophysical Union,
1168 Washington, D.C.
- 1169 Krijgsman, W., and L. Tauxe (2004), Shallow bias in Mediterranean paleomagnetic directions
1170 caused by inclination error, *Earth and Planetary Science Letters*, 222, 685–695.
- 1171 Krynine, P. D. (1950), Petrology, stratigraphy and origin of the Triassic sedimentary rocks of
1172 Connecticut, *Connecticut Geology and Natural History Survey Bulletin*, 73, 247 p.
- 1173 Larson, R. L., and P. Olson (1991), Mantle plumes control magnetic reversal frequency, *Earth*
1174 *and Planetary Science Letters*, 107, 437-447.
- 1175 LeTourneau, P. M., and P. Huber (2006), Early Jurassic eolian dune field, Pomperaug basin,
1176 Connecticut and related synrift deposits: Stratigraphic framework and paleoclimatic context,
1177 *Sedimentary Geology*, 187, 63–81.
- 1178 Lowrie, W. (1990), Identification of ferromagnetic minerals in a rock by coercivity and
1179 unblocking temperature properties, *Geophysical research Letters*, 17, 159-162.
- 1180 Lowrie, W., and D. V. Kent (2004), Geomagnetic polarity timescales and reversal frequency
1181 regimes, in *Timescales of the Paleomagnetic Field*, edited by J. E. T. Channell, D. V. Kent,
1182 W. Lowrie and J. Meert, pp. 117–129, American Geophysical Union, Washington, D.C.
- 1183 Lucas, S. G., A. B. Heckert, and P. Huber (1998), *Aetosaurus* (Archosauromorpha) from the
1184 Upper Triassic of the Newark Supergroup, eastern United States, and its biochronological
1185 significance, *Palaeontology*, 41, 1215–1230.
- 1186 Lucas, S. G., and P. Huber (2003), Vertebrate biostratigraphy and biochronology of the
1187 nonmarine Late Triassic, in *The Great Rift Valleys of Pangea in Eastern North America*,
1188 Volume 2: Sedimentology, Stratigraphy, and Paleontology, edited by P. M. LeTourneau and
1189 P. E. Olsen, pp. 143–191, Columbia University Press, New York.

- 1190 Luttrell, G. W. (1989), Stratigraphic nomenclature of the Newark Supergroup of Eastern North
1191 America, US Geological Survey Bulletin 1572, 136 pp.
- 1192 Marcano, M. C., R. Van der Voo, and C. Mac Niocaill (1999), True polar wander during the
1193 Permo-Triassic, *Journal of Geodynamics*, 28, 75-95.
- 1194 Marzoli, A., P. R. Renne, E. M. Piccirillo, M. Ernesto, G. Gellieni, and A. De Min (1999),
1195 Extensive 200-million-year-old continental flood basalts of the Central Atlantic Magmatic
1196 Province, *Science*, 284, 616-618.
- 1197 Marzoli, A., H. Bertrand, K. B. Knight, S. Cirilli, N. Buratti, C. V. rati, S. Nomade, P. R. Renne,
1198 N. Youbi, R. Martini, K. Allenbach, R. Neuwerth, C. Rapaille, L. Zaninetti, and G. Bellieni
1199 (2004), Synchrony of the Central Atlantic magmatic province and the Triassic-Jurassic
1200 boundary climatic and biotic crisis, *Geology*, 32, 973-976.
- 1201 May, S. R., and R. F. Butler (1986), North American Jurassic apparent polar wander:
1202 implications for plate motion, paleogeography and Cordilleran tectonics, *Journal of*
1203 *Geophysical Research*, 91, 11519-11544.
- 1204 McElhinny, M. W., and P. J. Burek (1971), Mesozoic paleomagnetic stratigraphy, *Nature*, 232,
1205 98-102.
- 1206 McEnroe, S. A., and L. L. Brown (2000), Palaeomagnetism, rock magnetism and geochemistry
1207 of Jurassic dykes and correlative redbeds, Massachusetts, USA, *Geophysical Journal*
1208 *International*, 143, 22-38.
- 1209 McFadden, P. L., and M. W. McElhinny (1990), Classification of the reversal test in
1210 palaeomagnetism, *Geophysical Journal International*, 103, 725-729.
- 1211 McHone, J. G. (2000), Non-plume magmatism and rifting during the opening of the central
1212 Atlantic Ocean, *Tectonophysics*, 316(3-4), 287-296.
- 1213 McIntosh, W. C., R. B. Hargraves, and C. L. West (1985), Paleomagnetism and oxide
1214 mineralogy of upper Triassic to lower Jurassic red beds and basalts in the Newark Basin,
1215 *Geological Society of America Bulletin*, 96, 463-480.
- 1216 Molina-Garza, R. S., J. W. Geissman, and R. Van der Voo (1995), Paleomagnetism of the
1217 Dockum Group (Upper Triassic), northwest Texas: Further evidence for the J1-cusp in the
1218 North America apparent polar wander path and implications for rate of Triassic apparent polar
1219 wander and Colorado plateau rotation, *Tectonics*, 14, 979-993.
- 1220 Molina-Garza, R. S., J. W. Geissman, and S. Lucas (2003), Paleomagnetism and
1221 magnetostratigraphy of the lower Glen Canyon and upper Chinle Groups, Jurassic-Triassic of
1222 northern Arizona and northeast Utah, *Journal of Geophysical Research*, 108(B4),
1223 doi:10.1029/2002JB001909.
- 1224 Muttoni, G., D. V. Kent, S. Meco, M. Balini, A. Nicora, R. Rettori, M. Gaetani, and L. Krystyn
1225 (1998), Towards a better definition of the Middle Triassic magnetostratigraphy and
1226 biostratigraphy in the Tethyan realm, *Earth and Planetary Science Letters*, 164, 285-302.
- 1227 Ogg, J. G., and M. B. Steiner (1991), Early Triassic magnetic polarity time scale - integration of
1228 magnetostratigraphy, ammonite zonation and sequence stratigraphy from stratotype sections
1229 (Canadian Arctic Archipelago), *Earth and Planetary Science Letters*, 107, 69-89.
- 1230 Olsen, P. E., and P. M. Galton (1977), Triassic-Jurassic extinctions: Are they real?, *Science*, 197,
1231 983-986.
- 1232 Olsen, P. E. (1986), A 40-million-year lake record of early Mesozoic orbital climatic forcing,
1233 *Science*, 234(Science), 842-848.
- 1234 Olsen, P. E., and D. V. Kent (1996), Milankovitch climate forcing in the tropics of Pangea
1235 during the Late Triassic, *Paleogeography, Paleoclimatology, Paleocology*, 122, 1-26.

- 1236 Olsen, P. E., and D. V. Kent (2008), Cyclostratigraphy and geochronology of the Early Jurassic
1237 lacustrine strata of the Hartford Basin (Newark Supergroup, Connecticut and Massachusetts,
1238 USA): Implications for the Jurassic time scale, *Earth and Planetary Science Letters*,
1239 submitted.
- 1240 Olsen, P. E., R. W. Schlichte, and M. S. Fedosh (1996b), 580 Ky duration of the Early Jurassic
1241 flood basalt event in eastern North America estimated using Milankovitch cyclostratigraphy,
1242 in *The Continental Jurassic*, edited by M. Morales, pp. 11-22, Museum of Northern Arizona.
- 1243 Olsen, P. E. (1999), Giant lava flows, mass extinctions, and mantle plumes, *Science*, 284, 604-
1244 605.
- 1245 Olsen, P. E., H.-D. Sues, and M. A. Norell (2000), First record of *Erpetosuchus* (Reptilia:
1246 Archosauria) from the Late Triassic of North America, *Journal of Vertebrate Paleontology*,
1247 20(4), 633–636.
- 1248 Olsen, P. E., D. V. Kent, H.-D. Sues, C. Koeberl, H. Huber, A. Montanari, E. C. Rainforth, S. J.
1249 Fowell, M. J. Szajna, and B. W. Hartline (2002a), Ascent of dinosaurs linked to an iridium
1250 anomaly at the Triassic-Jurassic boundary, *Science*, 296, 1305-1307.
- 1251 Olsen, P. E., C. Koeberl, H. Huber, A. Montanari, S. J. Fowell, M. Et Touhami, and D. V. Kent
1252 (2002b), Continental Triassic-Jurassic boundary in central Pangea: recent progress and
1253 discussion of an Ir anomaly, *Geological Society of America Special Paper*, 356, 505-522.
- 1254 Olsen, P. E., D. V. Kent, M. Et-Touhami, and J. Puffer (2003), Cyclo-, magneto-, and bio-
1255 stratigraphic constraints on the duration of the CAMP event and its relationship to the
1256 Triassic-Jurassic Boundary, in *The Central Atlantic Magnetic Province: Insights from*
1257 *Fragments of Pangea*, edited by W. E. Hames, J. G. McHone, P. R. Renne and C. Ruppel, pp.
1258 7-32 (DOI: 10.1029/1136GM1002), American Geophysical Union, Washington, DC.
- 1259 Olsen, P. E., J. H. Whiteside, P. M. LeTourneau, and P. Huber (2005), Jurassic cyclostratigraphy
1260 and paleontology of the Hartford basin, in *97th New England Intercollegiate Geological*
1261 *Conference*, edited by B. J. Skinner and A. R. Philpotts, pp. A4-1 – A4-51, Department of
1262 Geology and Geophysics, Yale University, New Haven, Connecticut.
- 1263 Opdyke, N. D., and H. Wensink (1966), Paleomagnetism of rocks from the White Mountain
1264 Plutonic-Volcanic series in New Hampshire and Vermont, *Journal of Geophysical Research*,
1265 71, 3045-3051.
- 1266 Pálffy, J., and R. Mundil (2006), The age of the Triassic/Jurassic boundary: new data and their
1267 implications for the extinction and recovery, *Volumina Jurassica*, IV, 294.
- 1268 Pechersky, D. M., and A. N. Khramov (1973), Mesozoic palaeomagnetism scale of the USSR,
1269 *Nature*, 244, 499-501.
- 1270 Philpotts, A. R., and A. Martello (1986), Diabase feeder dikes for the Mesozoic basalts in
1271 southern New England, *American Journal of Science*, 286(Amer. J. Sci.), 105-126.
- 1272 Pratt, L. M., C. A. Shaw, and R. C. Burruss (1988), Thermal histories of the Hartford and
1273 Newark basins inferred from maturation indices of organic matter, in *Studies of the Early*
1274 *Mesozoic Basins of the Eastern United States*, edited by A. J. Froelich and G. R. J. Robinson,
1275 pp. 58–62, US Geological Survey, Denver.
- 1276 Prevot, M., and M. McWilliams (1989), Paleomagnetic correlation of the Newark Supergroup
1277 volcanics, *Geology*, 17, 1007-1010.
- 1278 Roden, M. K., and D. S. Miller (1991), Tectonothermal history of Hartford, Deerfield, Newark
1279 and Taylorsville Basins, eastern United States, using fission-track analysis, *Schweizerische*
1280 *Mineralogische und Petrographische Mitteilungen*, 71, 187-203.

- 1281 Roden-Tice, M. K., and R. P. Wintsch (2002), Early Cretaceous normal faulting in southern New
1282 England: Evidence from apatite and zircon fission-track ages, *Journal of Geology*, 110, 159-
1283 178.
- 1284 Sanders, J. E. (1968), Stratigraphy and primary sedimentary structures of fine-grained, well-
1285 bedded strata, inferred lake deposits, Upper Triassic, central and southern Connecticut, in *Late*
1286 *Paleozoic and Mesozoic Continental Sedimentation, Northeastern North America*, edited by
1287 G. D. Klein, pp. 265-305, Geological Society of America, Boulder, CO.
- 1288 Sanders, J. E. (1970), Stratigraphy and structure of the Triassic strata of the Gaillard graben,
1289 south-central Connecticut, Connecticut Geological and Natural History Survey Guidebook, 3,
1290 15.
- 1291 Schaltegger, U., J. Guex, A. Bartolini, B. Schoene, and M. Ovtcharov (2007), Precise U-Pb age
1292 constraints for end-Triassic mass extinction, its correlation to volcanism and Hettangian post-
1293 extinction recovery, *Earth and Planetary Science Letters*, in press (doi:
1294 10.1016/j.epsl.2007.11.031).
- 1295 Schlische, R. W. (1993), Anatomy and evolution of the Triassic-Jurassic continental rift system,
1296 eastern North America., *Tectonics*, 12, 1026-1042.
- 1297 Schlische, R. W. (1995), Geometry and origin of fault-related folds in extensional settings,
1298 *American Association of Petroleum Geologists Bulletin*, 79, 1661-1678.
- 1299 Schoene, B., J. L. Crowley, D. J. Condon, M. D. Schmitz, and S. A. Bowring (2006),
1300 Reassessing the uranium decay constants for geochronology using IDOTIMS U-Pb data,
1301 *Geochimica et cosmochimica Acta*, 70, 426-445.
- 1302 Seidemann, D. E., W. D. Masterson, M. P. Dowling, and K. K. Turekian (1984), K-Ar dates and
1303 $^{40}\text{Ar}/^{39}\text{Ar}$ age spectra for Mesozoic basalt flows of the Hartford Basin, Connecticut, and the
1304 Newark Basin, New Jersey, *Geological Society of America Bulletin*, 95, 594-598.
- 1305 Seidemann, D. E. (1988), The hydrothermal addition of excess ^{40}Ar to the lava flows from the
1306 Early Jurassic in the Hartford Basin (northeastern U.S.A.): Implications for the time scale,
1307 *Chemical Geology (Isotope Geoscience Section)*, 72, 37-45.
- 1308 Seidemann, D. E. (1989), Age of the Triassic/Jurassic boundary: A view from the Hartford basin,
1309 *American Journal of Science*, 289, 553-562.
- 1310 Smith, T. E. (1976), Paleomagnetic study of lower Mesozoic diabase dikes and sills of
1311 Connecticut and Maryland, *Canadian Journal of Earth Sciences*, 13, 697-609.
- 1312 Smith, T. E., and H. C. Noltimier (1979), Paleomagnetism of the Newark Trend igneous rocks of
1313 the north central Appalachians and the opening of the central Atlantic Ocean, *American*
1314 *Journal of Science*, 279, 778-807.
- 1315 Smith, W. A. (1987), Paleomagnetic results from a crosscutting system of northwest and north-
1316 south trending diabase dikes in the North Carolina Piedmont, *Tectonophysics*, 136, 137-150.
- 1317 Steiner, M., J. G. Ogg, Z. Zhang, and S. Sun (1989), The Late Permian/Early Triassic magnetic
1318 polarity time scale and plate motions of South China, *Journal of Geophysical Research*, 94,
1319 7343-7363.
- 1320 Steiner, M. B., and J. G. Ogg (1988), Early and Middle Jurassic magnetic polarity time scale,
1321 2nd International Symposium on Jurassic Stratigraphy, 1097-1111.
- 1322 Steiner, M. B., and S. G. Lucas (2000), Paleomagnetism of the Late Triassic Petrified Forest
1323 Formation, Chinle Group, western United States: Further evidence of "large" rotation of the
1324 Colorado Plateau, *Journal of Geophysical Research*, 105, 791-725,808.
- 1325 Sutter, J. F. (1988), Innovative approaches to the dating of igneous events in the early Mesozoic
1326 basins of the Eastern United States, *U.S. Geological Survey Bulletin*, 1776, 194-200.

- 1327 Szurlies, M. (2004), Magnetostratigraphy: the key to a global correlation of the classic Germanic
1328 Trias - case study Volpriehausen Formation (Middle Buntsandstein) Central Germany, *Earth*
1329 *and Planetary Science Letters*, 227, 395–410.
- 1330 Tan, C., and K. P. Kodama (2002), Magnetic anisotropy and paleomagnetic inclination
1331 shallowing in red beds: Evidence from the Mississippian Mauch Chunk Formation,
1332 Pennsylvania, *Journal of Geophysical Research*, 107(B11), doi:10.1029/2001JB001636.
- 1333 Tan, X., K. P. Kodama, H. Chen, D. Fang, D. Sun, and Y. Li (2003), Paleomagnetism and
1334 magnetic anisotropy of Cretaceous red beds from the Tarim basin, northwest China: Evidence
1335 for a rock magnetic cause of anomalously shallow paleomagnetic inclinations from central
1336 Asia, *Journal of Geophysical Research*, 108(B2), doi:10.1029/2001JB001608.
- 1337 Tan, X., K. P. Kodama, S. Gilder, and V. Courtillot (2007), Rock magnetic evidence for
1338 inclination shallowing in the Passaic Formation red beds from the Newark basin and a
1339 systematic bias of the Late Triassic apparent polar wander path for North America, *Earth and*
1340 *Planetary Science Letters*, 254(3-4), 345-357.
- 1341 Tauxe, L., and D. V. Kent (1984), Properties of a detrital remanence carried by haematite from
1342 study of modern river deposits and laboratory redeposition experiments, *Geophysical Journal*
1343 *of the Royal astronomical Society*, 77, 543-561.
- 1344 Tauxe, L., and D. V. Kent (2004), A simplified statistical model for the geomagnetic field and
1345 the detection of shallow bias in paleomagnetic inclinations: Was the ancient magnetic field
1346 dipolar?, in *Timescales of the Paleomagnetic Field*, edited by J. E. T. Channell, D. V. Kent,
1347 W. Lowrie and J. Meert, pp. 101–115, AGU, Washington, D.C.
- 1348 Tollo, R. P., and D. Gottfried (1992), Petrochemistry of Jurassic basalt from eight cores, Newark
1349 basin, New Jersey: Implications for the volcanic petrogenesis of the Newark Supergroup, in
1350 *Eastern North American Mesozoic magmatism*, edited by J. H. Puffer and P. C. Ragland, pp.
1351 233–259, Geological Society of America, Boulder.
- 1352 Van Fossen, M. C., and D. V. Kent (1990), High-latitude paleomagnetic poles from Middle
1353 Jurassic plutons and Moat Volcanics in New England and the controversy regarding Jurassic
1354 apparent polar wander for North America, *Journal of Geophysical Research*, 95, 17,503-
1355 517,516.
- 1356 Wang, Z. S., E. T. Rasbury, G. N. Hanson, and W. J. Meyers (1998), Using the U-Pb system of
1357 calcretes to date the time of sedimentation of clastic sedimentary rocks, *Geochimica et*
1358 *cosmochimica Acta*, 62, 2823-2835.
- 1359 Wheeler, G. (1937), The west wall of the New England Triassic lowland, *State Geological and*
1360 *Natural History Survey of Connecticut, Bulletin*, 58, 73 p.
- 1361 Whiteside, J. H., P. E. Olsen, D. V. Kent, S. J. Fowell, and M. Et-Touhami (2007), Synchrony
1362 between the Central Atlantic magmatic province and the Triassic-Jurassic mass-extinction
1363 event?, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 244(1-4), 345-367.
- 1364 Wise, D. U., and P. Robinson (1982), Tectonics of the Mesozoic Connecticut Valley graben,
1365 Geological Society of America, *Abstracts with Programs*, 14, 96.
- 1366 Witte, W. K., and D. V. Kent (1990), The paleomagnetism of red beds and basalts of the
1367 Hettangian Extrusive Zone, Newark Basin, New Jersey, *Journal of Geophysical Research*, 95,
1368 17,533-517,545.
- 1369 Witte, W. K., and D. V. Kent (1991), Tectonic implications of a remagnetization event in the
1370 Newark Basin, *Journal of Geophysical Research*, 96, 19,569-519,582.



1371 Witte, W. K., D. V. Kent, and P. E. Olsen (1991), Magnetostratigraphy and paleomagnetic poles
1372 from Late Triassic-earliest Jurassic strata of the Newark Basin, Geological Society of America
1373 Bulletin, 103, 1648-1662.



1374 Yang, Z., M.-G. Moreau, H. Bucher, J.-L. Dommergues, and A. Trouiller (1996), Hettangian and
1375 Sinemurian magnetostratigraphy from Paris Basin, Journal of Geophysical Research, 101,
1376 8025-8042.









1377







1378







Central Atlantic Magmatic Province















Shuttle Meadow, East Berlin &
Portland Formations


N = 78 sites

ChRM




Shuttle Meadow, East Berlin &
Portland Formations


HARTFORD BASIN



NEWARK BASIN




HARTFORD BASIN



Paris Basin - Montcornet core

Yang et al. (1996)

U-Pb dates
(marine strata)



*(Palfy & Mundil, 2006;
Schaltegger et al., 2007)*

E/I analysis: East Berlin, Shuttle Meadow & Portland Formations

