

## Direct measurements of CO<sub>2</sub> flux in the Greenland Sea

Siv K. Lauvset,<sup>1,2,3</sup> Wade R. McGillis,<sup>4,5</sup> Ludovic Bariteau,<sup>6</sup> C. W. Fairall,<sup>7</sup>  
Truls Johannessen,<sup>1,2,3</sup> Are Olsen,<sup>2,3</sup> and Christopher J. Zappa<sup>4,5</sup>

Received 8 April 2011; revised 12 May 2011; accepted 16 May 2011; published 22 June 2011.

[1] During summer 2006 eddy correlation CO<sub>2</sub> fluxes were measured in the Greenland Sea using a novel system set-up with two shrouded LICOR-7500 detectors. One detector was used exclusively to determine, and allow the removal of, the bias on CO<sub>2</sub> fluxes due to sensor motion. A recently published correction method for the CO<sub>2</sub>-H<sub>2</sub>O cross-correlation was applied to the data set. We show that even with shrouded sensors the data require significant correction due to this cross-correlation. This correction adjusts the average CO<sub>2</sub> flux by an order of magnitude from  $-6.7 \times 10^{-2} \text{ mol m}^{-2} \text{ day}^{-1}$  to  $-0.61 \times 10^{-2} \text{ mol m}^{-2} \text{ day}^{-1}$ , making the corrected fluxes comparable to those calculated using established parameterizations for transfer velocity. **Citation:** Lauvset, S. K., W. R. McGillis, L. Bariteau, C. W. Fairall, T. Johannessen, A. Olsen, and C. J. Zappa (2011), Direct measurements of CO<sub>2</sub> flux in the Greenland Sea, *Geophys. Res. Lett.*, 38, L12603, doi:10.1029/2011GL047722.

### 1. Introduction

[2] Because the atmospheric CO<sub>2</sub> concentration is rising due to the burning of fossil fuels, land use changes, and cement production it is important to accurately quantify the size of the total ocean carbon sink and its variations with time. For this we need to know the air-sea CO<sub>2</sub> flux. Because it is difficult to measure the air-sea CO<sub>2</sub> flux (F<sub>CO<sub>2</sub></sub>) directly, global estimates mostly rely on calculations of the form

$$F_{\text{CO}_2} = kS\Delta f\text{CO}_2 \quad (1)$$

where  $\Delta f\text{CO}_2$  is the difference between the fugacity of CO<sub>2</sub> in the sea ( $f\text{CO}_2^{\text{sea}}$ ) and in the air ( $f\text{CO}_2^{\text{air}}$ ).  $S$  is the gas solubility, and  $k$  is an estimate of the gas transfer velocity usually parameterized as a function of wind speed at a mean height of 10 m above the water surface for neutral atmospheric stability ( $U_{10N}$ ). We define  $F_{\text{CO}_2}$  to be negative into the ocean. The most widely used parameterizations of  $k$  have been derived using tracer release experiments [*Ho et al.*, 2006; *Liss and Merlivat*, 1986; *Nightingale et al.*, 2000],

wind-wave tank experiments [*Liss and Merlivat*, 1986], and radiocarbon invasion rates [*Naegler et al.*, 2006; *Sweeney et al.*, 2007; *Wanninkhof*, 1992]. Yet, none of these capture the complete range of processes relevant to air-sea gas exchange, nor are they consistent. To resolve these issues we need direct measurements of the air-sea CO<sub>2</sub> flux.

[3] Direct measurements of the  $F_{\text{CO}_2}$  can be obtained using the eddy correlation (EC) method [*McGillis et al.*, 2001a, 2004; *Wanninkhof and McGillis*, 1999], but several difficulties with the EC method still exist after a decade of significant technical advances. First among these is the considerable greater magnitude of observed EC  $F_{\text{CO}_2}$  compared to bulk parameterization or tracer derived fluxes [*Broecker et al.*, 1986; *Kondo and Tsukamoto*, 2007; *Prytherch et al.*, 2010], resulting in few published data sets of  $F_{\text{CO}_2}$  from EC experiments. Recent research suggests that the measurements are too high due to a strong contamination of the CO<sub>2</sub> signal by water vapor, in addition to sensor motion contamination and dilution effects.

[4] In this paper we will present the first data set of EC  $F_{\text{CO}_2}$  measured in the Greenland Sea featuring unique environmental conditions, and using a novel instrument set-up. We also use this data set to test the suitability of the PKT correction method [*Prytherch et al.*, 2010] for data sets measured in such environmental conditions and with this instrument set-up.

### 2. Experiment and Methods

[5] The data were obtained on the Greenland Sea cruise 58GS20060721 [*Olsen and Omar*, 2007], carried out onboard the research vessel *G.O. Sars* between July 21 and August 3, 2006. The cruise started in Akureyri, Iceland and ended in Tromsø, Norway. The flux measurement system was set up on a mast installed at the bow of the ship ~14.5 m above the sea surface. Two LICOR-7500 open path non-dispersive infrared (NDIR) detectors, a 3D Gill Sonic anemometer, a Motionpak, and a compass were collocated on top of the ship mast. The NDIR detectors were mounted ~1 m from the sonic anemometer and motion system, and both were shrouded in rigid plastic housings. The shrouds prevent loss of data due to severe weather conditions and icing, leading to a more robust data set. The instrument set-up is schematically shown in Figure 1. Air entered the first sensor, hereafter referred to as ‘sample’, at 570 l min<sup>-1</sup> and passed through a mixing chamber connected to a high volume pump; the intake of the second sensor, hereafter referred to as ‘null’, is taken from the mixing chamber using a second flow path at 200 ml min<sup>-1</sup>. The sensors were mounted next to each other with the long axes aligned vertically so that they experienced the same motion, and the rigid fit in the shroud prevented flexing of the support structures of the detectors thus reducing the motion artifact. On subsequent

<sup>1</sup>Geophysical Institute, University of Bergen, Bergen, Norway.

<sup>2</sup>Uni Bjerkes Centre, Bergen, Norway.


<sup>3</sup>Bjerkes Centre for Climate Research, Bergen, Norway.

<sup>4</sup>Department of Earth and Environmental Engineering, Columbia University, New York, New York, USA.

<sup>5</sup>Lamont Doherty Earth Observatory, Columbia University, Palisades, New York, USA.

<sup>6</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Boulder, Colorado, USA.

<sup>7</sup>Earth System Research Laboratory, NOAA, Boulder, Colorado, USA.



**Figure 1.** Schematic of the instrument set-up onboard R/V *G.O. Sars* July 21 – August 3, 2006.

deployments it has been found that the mixing chamber acts like a low pass filter with a time constant of 300 s, and by using Equation 11 of Horst [1997] we calculate that using this set up the null sensor measurement fluctuations are reduced by ~97%. The remaining signal is thus essentially due to the motion alone. Motion contamination was first described by Fairall *et al.* [2000] and previously correction methods for close-path systems based on covariance with a calibration gas [McGillis *et al.*, 2001b] and a second identical null sensor with a sealed input [McGillis *et al.*, 2004] have been used, and correlation with measured ship motion variables [Yelland *et al.*, 2009; Miller *et al.*, 2010] for open-path systems which need less correction. We assume that the motion artifact is the same for both sensors and subtract this from the sample signal on a point-by-point basis.

[6] The ship was equipped with an underway fCO<sub>2</sub> system [Pierrot *et al.*, 2009] used to measure the fCO<sub>2</sub> in both the surface ocean and the atmosphere, the sea surface temperature (SST) and the sea surface salinity (SSS). This system is calibrated every 3–4 hours using three referenced standard gases obtained from the National Oceanographic and Atmospheric Administration Earth System Research Laboratory (NOAA/ESRL). Navigation data were retrieved from the ship’s measurement system.


[7] Collected data were processed in 10-minute blocks and fluxes were obtained by correlating the motion-corrected vertical velocity with the fast fluctuations of interest (CO<sub>2</sub>, temperature, H<sub>2</sub>O). The short averaging time does not present a problem since the portions of fluxes not resolved at low frequency represent only a small contribution to the total covariance. For details concerning how the high frequency wind speed measurements were corrected for the ship’s movement see Edson *et al.* [1998]. Only data with relative wind vectors ±90° to bow-on were used. Data

blocks where the standard deviation of ship speed and ship heading were greater than 0.2 m s<sup>-1</sup> and 5° respectively were removed, as were data with large motion corrections. The flow tilt calculated from the sonic anemometer was also used to roughly account for flow distortion effects [Fairall *et al.*, 1997]. Because of problems with the NDIR sensors and the ship compass during the first half of the cruise, the results of this study are mostly based on the last half of the cruise, east of ~7°W. Out of the total 1896 10-minute averages, 652 passed all quality controls.


[8] Both latent heat and CO<sub>2</sub> fluxes were computed from the sample NDIR sensor, while the sensible heat flux was computed from vertical velocity–sonic temperature covariance. The humidity contribution to sonic temperature was removed using the bulk latent heat flux using Equation 8 of Schotanus *et al.* [1983]. We use the bulk algorithm because the sonic anemometer produce more reliable timeseries than fast humidity sensors and because using the bulk algorithm eliminates any effect of spatial separation of the humidity sensor and the sonic anemometer. The effects of humidity and temperature on the CO<sub>2</sub> measurements were removed prior to calculating the flux by converting the measured molar densities into mixing ratios using the gas law and the high frequency temperature measurements and air pressure (which is of minor importance). The shroud on the sample sensor acts like a low pass filter with a time constant of 0.3 s which using Equation 11 of Horst [1997] gives a signal attenuation of ~5%. This error in the temperature dilution correction due to our set up is within acceptable boundaries. The shrouded set-up is designed to reduce the sensor contamination from rain and heavy sea spray, but the intake is not filtered. As a consequence a larger than expected CO<sub>2</sub>–H<sub>2</sub>O cross-correlation, which is most likely due to hygroscopic particles on the optical surfaces, was observed also after correcting for the dilution effect. Additional correction was therefore made using the PKT method [Prytherch *et al.*, 2010]. Consistent with subsequent deployments we see an increase in the magnitude of the correction with time, suggesting that contamination is the reason behind the cross-correlation.

### 3. Results and Discussion

[9] The average EC F<sub>CO<sub>2</sub></sub> calculated from the pre-PKT data is  $-6.7 \times 10^{-2} \text{ mol m}^{-2} \text{ day}^{-1}$ , with a standard deviation of  $0.27 \text{ mol m}^{-2} \text{ day}^{-1}$ . The raw CO<sub>2</sub> flux has considerable scatter, but there is a statistically significant (>95%) negative correlation with latent heat flux (F<sub>H<sub>2</sub>O</sub>, Figure 2a) which indicates that the PKT correction is warranted. The average post-PKT F<sub>CO<sub>2</sub></sub> is  $-0.61 \times 10^{-2} \text{ mol m}^{-2} \text{ day}^{-1}$ , with a standard deviation of  $0.11 \text{ mol m}^{-2} \text{ day}^{-1}$ . This is comparable to the CO<sub>2</sub> flux calculated using Wanninkhof’s [1992] k-U<sub>10N</sub> parameterization ( $-0.56 \times 10^{-2} \text{ mol m}^{-2} \text{ day}^{-1}$ ). The post-PKT F<sub>CO<sub>2</sub></sub> still have considerable scatter, especially when the F<sub>H<sub>2</sub>O</sub> is low, but there is no longer a negative correlation (Figure 2b). The post-PKT F<sub>CO<sub>2</sub></sub> is small (Figure 2b), but this is not unexpected given the calm wind conditions (Figure 3). The “flux” measured by the null sensor is on average  $-8.8 \times 10^{-4} \text{ mol m}^{-2} \text{ day}^{-1}$  with a standard deviation of  $0.018 \text{ mol m}^{-2} \text{ day}^{-1}$  (Figure 2c). Removing the null “flux” removes scatter from the sample flux data reducing the standard deviation from 0.23 to 0.11. The difference between flux data before and after removing the null “flux”



**Figure 2.** Eddy correlation CO<sub>2</sub> flux as a function of latent heat flux before and after the PKT correction. Two outliers are not shown on plot a ( $1.7$  and  $-6.3$  mol m<sup>-2</sup> day<sup>-1</sup>) and one on plot b ( $-4.8$  mol m<sup>-2</sup> day<sup>-1</sup>) to avoid the variability appearing smaller. (a) CO<sub>2</sub> flux from the sample LICOR before PKT correction, (b) CO<sub>2</sub> flux from the sample LICOR after PKT correction, also shown is a map of the cruise track covered between July 21, 2006 and August 3, 2006 where the red dot indicates 7°W (c) “flux” from the null LICOR. Note that this subplot has a different scale on the y-axis.




**Figure 3.** (a) The temporal CO<sub>2</sub> flux with the zero line indicated in grey, (b) the undersaturation ( $\Delta f\text{CO}_2$ ), (c) the sonic wind speed, (d) the surface ocean temperature, and (e) the air temperature during the cruise.

was analysed using a one-way ANOVA test and found to be statistically significant with  $>90\%$  confidence. This shows that even under very calm ocean conditions having a null sensor to remove the bias from motion is valuable.

[10] The  $\Delta f\text{CO}_2$  was large throughout the cruise but there are two distinct regimes: the cold, fresh Polar waters in the west with on average  $-119.5 \pm 13$  ( $1\sigma$ )  $\mu\text{atm}$ ; and the warmer and more saline Atlantic waters in the east with on average  $-76.5 \pm 12$  ( $1\sigma$ )  $\mu\text{atm}$  (Figure 3b). This implies a large negative flux of CO<sub>2</sub> into the sea. The wind speed during the cruise ranged from  $0.5$  m s<sup>-1</sup> to  $10.8$  m s<sup>-1</sup> with a mean of  $4.5 \pm 1.9$  ( $1\sigma$ ) m s<sup>-1</sup> (Figure 3c). 90% of all wind speed recorded were less than  $7$  m s<sup>-1</sup> and 15% less than  $2.5$  m s<sup>-1</sup> so we have a quite large data set of  $F_{\text{CO}_2}$  at very low wind speeds. No previously published EC experiment has reported significant amounts of data at wind speeds less than  $2.5$  m s<sup>-1</sup> so the Greenland Sea experiment is in this respect unique.

[11] Transfer velocity ( $k$ ) was calculated using equation (1) and bin averaged in  $2$  m s<sup>-1</sup>  $U_{10N}$  intervals (Figure 4). The pre-PKT fluxes yield a very strong non-linear relationship in  $k$ - $U_{10N}$ , while the corrected fluxes imply a  $k$ - $U_{10N}$  relationship in the same range as established parameterizations. There are, however, significantly higher  $k$  at very low wind speeds than has been reported previously, and the two first bin averages in Figure 4 are significantly higher than Wanninkhof’s [1992] line (grey). The second is biased high due to three outliers in this data subset, while the first have very low  $F_{\text{H}_2\text{O}}$  which could lead to the PKT correction not



**Figure 4.** (top) The  $k$  bin-averaged in  $2$  m s<sup>-1</sup> wind speed intervals plotted against  $U_{10N}$ . The number of data points in each bin is 23, 313, 168, 40, 25, and 7 respectively. See the legend for further details. The error bars show the standard error of the mean. (bottom) Close-up of the post-PKT  $k$ . The thin black lines show the 95% confidence interval (estimated as plus or minus two times the standard error of the mean).

working optimally [Prytherch *et al.*, 2010]. There is no significant diurnal cycle in the  $F_{\text{CO}_2}$  and the buoyancy fluxes so diurnal processes, like nighttime mixed layer deepening and surface renewal, cannot explain the elevated  $k$  at very low wind speeds like it could in the equatorial Pacific [McGillis *et al.*, 2004]. The dramatic increase in the PKT correction with wind speed, which is also seen in unpublished flux data from the Southern Ocean Gas Exchange Experiment, is likely due to there being only a few data points at winds greater than  $8 \text{ m s}^{-1}$ , which all have significantly higher  $F_{\text{H}_2\text{O}}$  than the data at lower winds. This thus leads to the humidity dependent PKT correction being larger. The variability in the corrected data is quite large, and our data set is too small and the wind speed range too narrow to either confirm previous or derive a new  $k\text{-}U_{10\text{N}}$  relationship.

#### 4. Conclusions

[12] Application of the PKT correction method to observations of EC CO<sub>2</sub> flux from the Greenland Sea lowers EC flux data by an order of magnitude, thus making the corrected direct fluxes comparable to bulk fluxes determined using established  $k\text{-}U_{10\text{N}}$  parameterizations, at least within the narrow range of wind speeds we encountered at this cruise. However, the combined high variability and limited range in wind speed means that the results from this data set cannot confirm established parameterizations. However, given the magnitude of the correction needed for these data despite using shrouded, and thus somewhat weatherproofed, sensors, it is clear that we need a more dedicated effort to understand the mechanisms causing the large CO<sub>2</sub>-H<sub>2</sub>O cross-correlation. Presented in this study are data at wind speeds less than  $2.5 \text{ m s}^{-1}$ , and at the overall low wind speeds experienced during this cruise the flux of CO<sub>2</sub> is small and the variability is large. This despite the large  $\Delta f\text{CO}_2$  which suggests that the potential for carbon uptake is very large, but apparently not utilized in the summer due to low wind speeds. Measurements in fall and winter are obviously needed in order to get a robust estimate of the size of the Greenland Sea carbon sink.

[13] **Acknowledgments.** This analysis has been funded by the Norwegian Research Council through the project CARBON-HEAT (185093/S30). The 2006 Greenland Sea research cruise onboard R/V *G.O. Sars* was funded by the CARBOOCEAN EU IP (5111176-2). We would like to thank the captain and crew onboard *G.O. Sars* for their help and cooperation during the cruise, as well as all others who helped setting up and taking down the system. This is publication A331 from the Bjerknes Centre for Climate Research.

[14] The Editor thanks Brent Else and an anonymous reviewer for their assistance in evaluating this paper.

#### References

- Broecker, W. S., J. R. Ledwell, T. Takahashi, R. Weiss, L. Merlivat, L. Memery, T. H. Peng, B. Jahne, and K. O. Munnich (1986), Isotopic versus micrometeorological ocean CO<sub>2</sub> fluxes: A serious conflict, *J. Geophys. Res.*, *91*(C9), 10,517–10,527, doi:10.1029/JC091iC09p10517.
- Edson, J. B., A. A. Hinton, K. E. Prada, J. E. Hare, and C. W. Fairall (1998), Direct covariance flux estimates from mobile platforms at sea, *J. Atmos. Oceanic Technol.*, *15*(2), 547–562, doi:10.1175/1520-0426(1998)015<0547:DCFEFM>2.0.CO;2.
- Fairall, C. W., A. B. White, J. B. Edson, and J. E. Hare (1997), Integrated shipboard measurements of the marine boundary layer, *J. Atmos. Oceanic Technol.*, *14*(3), 338–359, doi:10.1175/1520-0426(1997)014<0338:ISMOTM>2.0.CO;2.
- Fairall, C. W., J. E. Hare, J. B. Edson, and W. R. McGillis (2000), Parameterization and micrometeorological measurement of air-sea gas transfer, *Boundary Layer Meteorol.*, *96*(1–2), 63–106, doi:10.1023/A:1002662826020.
- Ho, D. T., C. S. Law, M. J. Smith, P. Schlosser, M. Harvey, and P. Hill (2006), Measurements of air-sea gas exchange at high wind speeds in the Southern Ocean: Implications for global parameterizations, *Geophys. Res. Lett.*, *33*, L16611, doi:10.1029/2006GL026817.
- Horst, T. W. (1997), A simple formula for attenuation of eddy fluxes measured with first-order-response scalar sensors, *Boundary Layer Meteorol.*, *82*(2), 219–233, doi:10.1023/A:1000229130034.
- Kondo, F., and O. Tsukamoto (2007), Air-sea CO<sub>2</sub> flux by eddy covariance technique in the equatorial Indian Ocean, *J. Oceanogr.*, *63*(3), 449–456, doi:10.1007/s10872-007-0040-7.
- Liss, P. S., and L. Merlivat (1986), Air-sea gas exchange rates: Introduction and synthesis, in *The Role of Air-Sea Exchange in Geochemical Cycling*, edited by P. Buat-Ménard, pp. 113–127, D. Reidel, Dordrecht, Netherlands.
- McGillis, W. R., J. B. Edson, J. E. Hare, and C. W. Fairall (2001a), Direct covariance air-sea CO<sub>2</sub> fluxes, *J. Geophys. Res.*, *106*(C8), 16,729–16,745, doi:10.1029/2000JC000506.
- McGillis, W. R., J. B. Edson, J. D. Ware, J. W. H. Dacey, J. E. Hare, C. W. Fairall, and R. Wanninkhof (2001b), Carbon dioxide flux techniques performed during GasEx-98, *Mar. Chem.*, *75*, 267–280, doi:10.1016/S0304-4203(01)00042-1.
- McGillis, W. R., *et al.* (2004), Air-sea CO<sub>2</sub> exchange in the equatorial Pacific, *J. Geophys. Res.*, *109*, C08S02, doi:10.1029/2003JC002256.
- Miller, S. D., C. Marandino, and E. S. Saltzman (2010), Ship-based measurement of air-sea CO<sub>2</sub> exchange by eddy covariance, *J. Geophys. Res.*, *115*, D02304, doi:10.1029/2009JD012193.
- Naegler, T., P. Ciais, K. Rodgers, and I. Levin (2006), Excess radiocarbon constraints on air-sea gas exchange and the uptake of CO<sub>2</sub> by the oceans, *Geophys. Res. Lett.*, *33*, L11802, doi:10.1029/2005GL025408.
- Nightingale, P. D., G. Malin, C. S. Law, A. J. Watson, P. S. Liss, M. I. Liddicoat, J. Boutin, and R. C. Upstill-Goddard (2000), In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cycles*, *14*(1), 373–387, doi:10.1029/1999GB900091.
- Olsen, A., and A. Omar (2007), G.O.Sars Repeat Section 75N\_2006 cruise data, [http://cdiac.ornl.gov/oceans/RepeatSections/clivar\\_75N.html](http://cdiac.ornl.gov/oceans/RepeatSections/clivar_75N.html), Carbon Dioxide Inf. Anal. Cent., Oak Ridge Natl. Lab., U.S. Dep. of Energy, Oak Ridge, Tenn.
- Pierrot, D., C. Neill, K. Sullivan, R. Castle, R. Wanninkhof, H. Luger, T. Johannessen, A. Olsen, R. A. Feely, and C. E. Cosca (2009), Recommendations for autonomous underway pCO<sub>2</sub> measuring systems and data-reduction routines, *Deep Sea Res., Part II*, *56*(8–10), 512–522, doi:10.1016/j.dsr2.2008.12.005.
- Prytherch, J., M. J. Yelland, R. W. Pascal, B. I. Moat, I. Skjelvan, and C. C. Neill (2010), Direct measurements of the CO<sub>2</sub> flux over the ocean: Development of a novel method, *Geophys. Res. Lett.*, *37*, L03607, doi:10.1029/2009GL041482.
- Schotanus, P., F. T. M. Nieuwstadt, and H. A. R. Debruin (1983), Temperature-measurement with a sonic anemometer and its application to heat and moisture fluxes, *Boundary Layer Meteorol.*, *26*(1), 81–93, doi:10.1007/BF00164332.
- Sweeney, C., E. Gloor, A. R. Jacobson, R. M. Key, G. McKinley, J. L. Sarmiento, and R. Wanninkhof (2007), Constraining global air-sea gas exchange for CO<sub>2</sub> with recent bomb <sup>14</sup>C measurements, *Global Biogeochem. Cycles*, *21*, GB2015, doi:10.1029/2006GB002784.
- Wanninkhof, R. (1992), Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, *97*(C5), 7373–7382, doi:10.1029/92JC00188.
- Wanninkhof, R., and W. R. McGillis (1999), A cubic relationship between air-sea CO<sub>2</sub> exchange and wind speed, *Geophys. Res. Lett.*, *26*(13), 1889–1892, doi:10.1029/1999GL900363.
- Yelland, M. J., R. W. Pascal, P. K. Taylor, and B. I. Moat (2009), Auto-Flux: An autonomous system for the direct measurement of the air-sea fluxes of CO<sub>2</sub>, heat and momentum, *J. Oper. Oceanogr.*, *2*(1), 15–23.
- L. Bariteau, Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, 325 Broadway, Boulder, CO 80309, USA.
- C. W. Fairall, Earth System Research Laboratory, NOAA, 325 Broadway, Boulder, CO 80305, USA.
- T. Johannessen and S. K. Lauvset, Geophysical Institute, University of Bergen, Allégaten 70, N-5007 Bergen, Norway. (siv.lauvset@gfi.uib.no)
- W. R. McGillis and C. J. Zappa, Lamont Doherty Earth Observatory, Columbia University, 61 Rte. 9W, Palisades, NY 10964, USA.
- A. Olsen, Uni Bjerknes Centre, Allégaten 55, N-5007 Bergen, Norway.