

United States  
Department of  
Agriculture

Forest Service



**Southern  
Research Station**

e-General Technical  
Report SRS-101

# Proceedings

## 15<sup>th</sup> Central Hardwood Forest Conference

**Knoxville, TN**

**February 27–March 1, 2006**



This document is being published in electronic format only (Web and CD). Any corrections or additions will be posted to the Web site ([www.srs.fs.usda.gov/pubs](http://www.srs.fs.usda.gov/pubs)).

Cover photo by: David J. Moorhead, University of Georgia,  
[www.forestryimages.org](http://www.forestryimages.org).

---

The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the U.S. Department of Agriculture Forest Service or the University of Tennessee of any product or service to the exclusion of others that may be suitable. Statements by contributors from outside the U.S. Department of Agriculture Forest Service or the University of Tennessee may not necessarily reflect the policy of the Department or the University.

---

---

All articles were received in digital format and were edited for uniform type and style; each author is responsible for the accuracy and content of his or her own paper. Statements of contributors from outside the U.S. Department of Agriculture may not necessarily reflect the policy of the Department.

---



Remarks about pesticides appear in some technical papers contained in these proceedings. Publication of these statements does not constitute endorsement or recommendation of them by the conference sponsors, nor does it imply that uses discussed have been registered. Use of most pesticides is regulated by State and Federal Law. Applicable regulations must be obtained from the appropriate regulatory agencies.

**CAUTION:** Pesticides can be injurious to humans, domestic animals, desirable plants, and fish and other wildlife—if they are not handled and applied properly. Use all pesticides selectively and carefully. Follow recommended practices given on the label for use and disposal of pesticides and pesticide containers.

---

April 2007

Southern Research Station  
200 W.T. Weaver Blvd.  
Asheville, NC 28804



# **Proceedings**

# **15<sup>th</sup> Central Hardwood**

# **Forest Conference**

Proceedings of a Conference held at the  
University of Tennessee, Knoxville, TN  
February 27 – March 1, 2006

**Edited by:**

David S. Buckley  
Wayne K. Clatterbuck

**Sponsored by:**

University of Tennessee, Department of Forestry,  
Wildlife and Fisheries, Knoxville, TN,  
and U.S. Department of Agriculture Forest Service,  
Southern Research Station, Asheville, NC

**Program Committee:**

David S. Buckley  
Stacy L. Clark  
Wayne K. Clatterbuck  
Callie J. Schweitzer

**Published by:**

U.S. Department of Agriculture Forest Service  
Southern Research Station  
Asheville, NC

## **FOREWORD**

The Central Hardwood Forest Conference is a series of biennial meetings that have been hosted by universities and research stations of the U.S. Department of Agriculture Forest Service in the central hardwood forest region in the Eastern United States. The objective of the conference is to bring together forest managers and scientists to discuss research and issues concerning the ecology and management of forests in the central hardwood region. This, the 15th Conference, included presentations pertaining to forest health and protection; ecology and forest dynamics; natural and artificial regeneration; forest products; wildlife; site classification; management and forest resources; mensuration and models; soil and water; agroforestry; and fire. The conference consisted of 86 oral presentations and 30 poster presentations resulting in the papers and abstracts published here.

## **STEERING COMMITTEE**

David S. Buckley	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN
Stacy L. Clark	USDA Forest Service, Southern Research Station, Normal, AL
Wayne K. Clatterbuck	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN
Callie J. Schweitzer	USDA Forest Service, Southern Research Station, Normal, AL

## **CONCURRENT SESSION LEADERS**

David Buckley	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN
Daniel Cassidy	University of Georgia, Athens, GA
Stacy Clark	USDA Forest Service, Southern Research Station, Normal, AL
Wayne Clatterbuck	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN
Jennifer Franklin	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN
Sam Jackson	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN
Karen Kuers	University of the South, Forestry and Geology Dept., Sewanee, TN
Chris Oswalt	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN
David Mercker	University of Tennessee, Forestry, Wildlife and Fisheries, Jackson, TN
Callie Schweitzer	USDA Forest Service, Southern Research Station, Normal, AL
Jeff Stringer	University of Kentucky, Department of Forestry, Lexington, KY
Adam Taylor	University of Tennessee, Forestry, Wildlife and Fisheries, Knoxville, TN

The Steering Committee of the 15th Central Hardwood Forest Conference wishes to thank Carol Whitlock and other members of the USDA Forest Service Southern Research Station communications unit for technical editing and publishing of these proceedings.

David S. Buckley coordinated the peer review process and Wayne K. Clatterbuck coordinated the meeting facilities and the registration.

## REVIEW PROCEDURES

Manuscripts for oral presentations were assigned to one of the editors and peer-reviewed by at least two professionals unless otherwise indicated. Reviews were returned to authors to revise their manuscripts. Revised manuscripts were then submitted to the Southern Research Station, Forest Service U.S. Department of Agriculture for final editing and publishing. Authors are responsible for the accuracy and content of their papers.

Mary A. Arthur, David S. Buckley, Stacy L. Clark, Wayne K. Clatterbuck, Luben Dimov, Jordan M. Marshall, Christopher M. Oswalt, Callie J. Schweitzer, and Jeffrey W. Stringer comprised the conference review team and provided reviews of abstracts and multiple manuscripts.

The Steering Committee is grateful to the following professionals for providing peer reviews of manuscripts for the proceedings:

Mary Beth Adams	Burnell Fischer	William B. Kurtz	Ronald A. Rathfon
Scott Bowe	Ross Fitzhugh	David R. Larsen	Stephen R. Shifley
Don C. Bragg	W. Mark Ford	Andrew J. Lister	Robert L. Smith
Brian K. Brashaw	Jennifer A. Franklin	Brian R. Lockhart	Martin A. Spetich
Patrick H. Brose	Lee E. Frelich	Robert P. Long	Philip Steele
Quang V. Cao	Emile S. Gardiner	William Luppold	Ron Stephens
Daniel Cassidy	Kurt W. Gottschalk	William MacDonald	Matthew A. Struckhoff
Neil Clark	Peter Gould	Brian C. McCarthy	Daniel K. Struve
Dean W. Coble	Arthur E. Gover	W. Henry McNab	Daniel J. Twedt
Mark Cowell	Bill Healy	Gary W. Miller	Daniel R. Unger
David R. Coyle	Eric Heitzman	Randall S. Morin	J.W. Van Sambeek
Michael Currier	John D. Hodges	Charles P. Nicholson	Siqun Wang
Jeffrey O. Dawson	Shelby G. Jones	David Ostermeier	Christopher R. Webster
Kenneth G. Day	John Kabrick	George R. Parker	Jake F. Weltzin
Michael Demchik	Richard Keim	James B. Pickens	Harry V. Wiant, Jr.
John Edwards	Patrick D. Keyser	Felix Ponder, Jr.	Daniel A. Yaussy
Pamela J. Edwards	Karen Kuers	KaDonna Randolph	James Zaczek
Andrew Ezell			

The 15th Central Hardwood Forest Conference logo and website were developed by Jordan M. Marshall.

The Steering Committee thanks the following people for their contribution to the 15th Central Hardwood Forest Conference: Jordan M. Marshall (Web site), Brien Ostby and Chris Oswalt (logistics), and Mirian Wright (registration and budget), all with the Department of Forestry, Wildlife and Fisheries, at the University of Tennessee. Their work was essential to our effort.

# CONTENTS

<b>PLENARY PAPERS</b> .....	<b>1</b>
<b>The Resilience of Upland-Oak Forest Canopy Trees to Chronic and Acute Precipitation Manipulations</b> .....	<b>3</b>
<i>Paul J. Hanson, Timothy J. Tschaplinski, Stan D. Wullschleger, Donald E. Todd, Jr., and Robert M. Augé</i>	
<b>Carbon Dioxide Fluxes in a Central Hardwoods Oak-Hickory Forest Ecosystem</b> .....	<b>13</b>
<i>Stephen G. Pallardy, Lianhong Gu, Paul J. Hanson, Tilden P. Myers, Stan D. Wullschleger, Bai Yang, Jeffery S. Riggs, Kevin P. Hosman, and Mark Heuer</i>	
<b>MENSURATION AND MODELS</b> .....	<b>21</b>
<b>The Sine Method as a More Accurate Height Predictor for Hardwoods</b> .....	<b>23</b>
<i>Don C. Bragg</i>	
<b>A Diameter Distribution Approach to Estimating Average Stand Dominant Height in Appalachian Hardwoods</b> .....	<b>33</b>
<i>John R. Brooks</i>	
<b>Development of Interim Oak Assessment Guidelines for the SILVAH Decision-Support System</b> .....	<b>37</b>
<i>Patrick H. Brose</i>	
<b>Digital Photo Monitoring for Tree Crown Foliage Change Evaluation</b> .....	<b>46</b>
<i>Neil Clark and Sang-Mook Lee</i>	
<b>Stocking Equations for Regeneration in Mixed Oak Stands</b> .....	<b>55</b>
<i>Songlin Fei, Kim C. Steiner, and James C. Finley</i>	
<b>A Form of Two-Phase Sampling Utilizing Regression Analysis</b> .....	<b>60</b>
<i>Michael A. Fiery and John R. Brooks</i>	
<b>Evaluation of LANDSAT Imagery for Detecting Ice Storm Damage in Upland Forests of Eastern Kentucky</b> .....	<b>69</b>
<i>W. Henry McNab, Tracy Roof, Jeffrey F. Lewis, and David L. Loftis</i>	
<b>Mesavage and Girard Form Class Taper Functions Derived from Profile Equations</b> .....	<b>77</b>
<i>Thomas G. Matney and Emily B. Schultz</i>	
<b>Predicting the Cover-Up of Dead Branches Using a Simple Single Regressor Equation</b> .....	<b>86</b>
<i>Christopher M. Oswald, Wayne K. Clatterbuck, and E.C. Burkhardt</i>	
<b>SOIL AND WATER</b> .....	<b>95</b>
<b>Long-Term Stream Chemistry Monitoring on the Fernow Experimental Forest: Implications for Sustainable Management of Hardwood Forests</b> .....	<b>97</b>
<i>Mary Beth Adams and James N. Kochenderfer</i>	
<b>Ecosystem Restoration Treatments Affect Soil Physical and Chemical Properties in Appalachian Mixed Oak Forests</b> .....	<b>107</b>
<i>Ralph E.J. Boerner, Jennifer A. Brinkman, and Daniel A. Yaussy</i>	

<b>Reclamation of Skid Roads with Fiber Mats and Native Vegetation: Effects on Erosion .....</b>	<b>116</b>
<i>Shawn T. Grushecky, David W. McGill, William Grafton, John Edwards, and Lisa Tager</i>	
<b>Impact of Alternative Harvesting Technologies on Thinning Entry and Optimal Rotation Age for Eastern Hardwoods.....</b>	<b>122</b>
<i>Chris B. LeDoux</i>	
<b>Nitrogen Dynamics Post-Harvest: The Role of Woody Residues.....</b>	<b>129</b>
<i>Kathryn Piatek</i>	
<b>Biomass Removal and Its Effect on Productivity of an Artificially Regenerated Forest Stand in the Missouri Ozarks .....</b>	<b>135</b>
<i>Felix Ponder, Jr.</i>	
<b>Attributes of Down Woody Materials in Hardwood Forests of the Eastern United States.....</b>	<b>144</b>
<i>Christopher W. Woodall, Sonja N. Oswald, and Randall S. Morin</i>	
<b>FOREST HEALTH AND PROTECTION .....</b>	<b>155</b>
<b>Impact of the Hemlock Woolly Adelgid on Radial Growth of Eastern Hemlock in Pennsylvania .....</b>	<b>157</b>
<i>Donald D. Davis, Matthew S. Fromm, and Matthew D. Davis</i>	
<b>Testing the Efficacy of Triclopyr and Imazapyr Using Two Application Methods for Controlling Tree-Of-Heaven Along a West Virginia Highway .....</b>	<b>163</b>
<i>William E. Eck and David W. McGill</i>	
<b>Abundance of <i>Armillaria</i> within Old-Growth Eastern Hemlock Stands in South-Central Pennsylvania.....</b>	<b>169</b>
<i>Matthew. S. Fromm and Donald D. Davis</i>	
<b>Rotation Length Based on a Time Series Analysis of Timber Degradate Caused by Oak Borers.....</b>	<b>176</b>
<i>Richard P. Guyette, Rose-Marie Muzika, and Aaron Stevenson</i>	
<b>Red Oak Decline and Mortality by Ecological Land Type in the Missouri Ozarks .....</b>	<b>181</b>
<i>John M. Kabrick, Zhaofei Fan, and Stephen R. Shifley</i>	
<b>Herbicide Treatments for Controlling Invasive Bush Honeysuckle in a Mature Hardwood Forest in West-Central Indiana .....</b>	<b>187</b>
<i>Ron Rathfon and Keith Ruble</i>	
<b>Cottonwood Leaf Beetle Control with Imidacloprid Soaked Cuttings.....</b>	<b>198</b>
<i>Terry L. Robison and Randall J. Rousseau</i>	
<b>Down Deadwood Dynamics on a Severely Impacted Oak Decline Site .....</b>	<b>206</b>
<i>Martin A. Spetich</i>	
<b>Defoliation and Oak Mortality in Southern New England .....</b>	<b>214</b>
<i>Jeffrey S. Ward</i>	
<b>FIRE.....</b>	<b>223</b>
<b>Survival of Striped Maple Following Spring Prescribed Fires in Pennsylvania.....</b>	<b>225</b>
<i>Patrick H. Brose, Gary W. Miller, and Kurt W. Gottschalk</i>	

<b>Fuels Consumption and Nitrogen Loss Following Prescribed Fire: A Comparison of Prescription Types in the Southern Appalachians.....</b>	<b>231</b>
<i>Barton D. Clinton and James M. Vose</i>	
<b>Initial Effects of Prescribed Burning and Thinning on Plant Communities in the Southeast Missouri Ozarks .....</b>	<b>241</b>
<i>E.R. McMurry, Rose-Marie Muzika, E.F. Loewenstein, K.W. Grabner, and G.W. Hartman</i>	
<b>Effects of Landscape Position and Season of Burn on Fire Temperature in Southern Ohio's Mixed Oak Forests .....</b>	<b>250</b>
<i>Doug J. Schwemlein and Roger A. Williams</i>	
<b>Forest Fuels and Landscape-Level Fire Risk Assessment of the Ozark Highlands, Missouri.....</b>	<b>258</b>
<i>Michael C. Stambaugh, Richard P. Guyette, and Daniel C. Dey</i>	
<b>MANAGEMENT AND FOREST RESOURCES.....</b>	<b>267</b>
<b>The Impact of Thinning and Fertilization Treatments on Sugar Concentration, Volume, and Total Sugar of Silver Maple Sap.....</b>	<b>269</b>
<i>M.L. Crum, J.J. Zaczek, J.E. Preece, S.G. Baer, and J.K. Buchheit</i>	
<b>Alternative Silvicultural Practices in Appalachian Forest Ecosystems: Implications for Species Diversity, Ecosystem Resilience, and Commercial Timber Production .....</b>	<b>276</b>
<i>Thomas R. Fox, Carola A. Haas, David W. Smith, David L. Loftis, Shepard M. Zedaker, Robert H. Jones, and A.L. Hammett</i>	
<b>Forest Certification and Nonindustrial Private Forest Landowners: Who Will Consider Certifying and Why? .....</b>	<b>281</b>
<i>David C. Mercker and Donald G. Hodges</i>	
<b>The Importance and Distribution of Hickory Across Virginia .....</b>	<b>286</b>
<i>Anita K. Rose and James F. Rosson, Jr.</i>	
<b>The Past, Present, and Future of Indiana's Oak Forests .....</b>	<b>295</b>
<i>Stephen R. Shifley and Christopher W. Woodall</i>	
<b>SITE CLASSIFICATION.....</b>	<b>305</b>
<b>Ecosystem Classification and Succession in the Central Till Plain of Indiana .....</b>	<b>307</b>
<i>Benjamin J. Dolan and George R. Parker</i>	
<b>Forest Conservation in the Cumberland Plateau and Mountains: Assessing Distribution and Structure of Landform Forest Associations.....</b>	<b>317</b>
<i>Daniel L. Druckenbrod, Virginia H. Dale, and Lisa M. Olsen</i>	
<b>Influence of Landform and Soil Characteristics on Canopy and Ground-Flora Composition and Structure of First- and Second-Order Headwater Riparian Forests in Unglaciaded Ohio .....</b>	<b>325</b>
<i>Kathryn L. Holmes, P. Charles Goebel, and David M. Hix</i>	
<b>A Multi-Criteria GIS Analysis for Ranking Potential Restoration Areas in the Fragmented Kaskaskia River Watershed Bottomland Hardwood Forest .....</b>	<b>337</b>
<i>Jean C. Mangun, Michael D. Gaskins, Andrew D. Carver, Karl W.J. Williard, and James J. Zaczek</i>	
<b>Aborted Yellow-Poplar Geographic Seed Source Test Serves to Verify Productivity on Cumberland Plateau Undulating Sandstone Uplands: 42-Year Results .....</b>	<b>345</b>
<i>Glendon W. Smalley and Elliot D. Olgivie</i>	



<b>WILDLIFE</b> .....	<b>353</b>
<b>A Method of Quantifying Forest Vertical Structure for the Purpose of Evaluating Bat Habitat</b> .....	<b>355</b>
<i>Marne M. Avina, Roger A. Williams, and Stanley D. Gehrt</i>	
<b>Developing Management Guidelines for Cerulean Warbler Breeding Habitat</b> .....	<b>364</b>
<i>Paul B. Hamel and Kenneth V. Rosenberg</i>	
<b>Ruffed Grouse (<i>Bonasa Umbellus</i>) Use of Stands Harvested Via Alternative Regeneration Methods in the Southern Appalachians</b> .....	<b>375</b>
<i>Benjamin C. Jones and Craig A. Harper</i>	
<b>Effects of Long-Term Prescribed Fire on Small Mammal Population Dynamics and Movement in an Oak Barrens Community in Tennessee – Preliminary Results</b> .....	<b>383</b>
<i>Rebecca L. Stratton and Wayne K. Clatterbuck</i>	
<b>Forest Management to Improve Breeding Habitat for Priority Songbirds in Upland Oak-Hickory Forests</b> .....	<b>388</b>
<i>Benjamin S. Thatcher, David A. Buehler, Patrick D. Martin, and Robert M. Wheat</i>	
<b>Kentucky Hunter Perceptions of Harvest Regulations and Their Effects on White-Tailed Deer Populations</b> .....	<b>400</b>
<i>Kara W. Throgmorton, Jean C. Mangun, and Andrew D. Carver</i>	
<b>AGROFORESTRY</b> .....	<b>407</b>
<b>Establishment of Upland and Bottomland Agroforestry Plantations in Tennessee and Mississippi</b> .....	<b>409</b>
<i>David M. Casey, Scott E. Schlarbaum, John T. Ammons, Fred L. Allen, Donald G. Hodges, William G. Minser, III, Arnold M. Saxton, Jason S. Maxedon, Chad Pope, and Chris R. Graves</i>	
<b>A Preliminary Economic Analysis of Silvopasture in Missouri’s Ozark Forests</b> .....	<b>418</b>
<i>Larry D. Godsey, W.D. “Dusty” Walter, John P. Dwyer, H.E. “Gene” Garrett</i>	
<b>Epicormic Response When Converting Hardwood Forests to a Silvopasture</b> .....	<b>425</b>
<i>W.D. “Dusty” Walter, Daniel C. Dey, and John P. Dwyer</i>	
<b>FOREST PRODUCTS</b> .....	<b>433</b>
<b>Assessing Veneer Log Quality Attributes</b> .....	<b>435</b>
<i>Delton Alderman, David Brinberg, and R.O. Goodykoontz</i>	
<b>Current Trends in the U.S. Wood Flooring Industry</b> .....	<b>443</b>
<i>Brian H. Bond, Matt Bumgardner, and Omar Espinoza</i>	
<b>The Occurrence of Log Ellipticality in Hardwoods and Its Impact on Lumber Value and Volume Recovery</b> .....	<b>451</b>
<i>Brian Bond, Janice K. Wiedenbeck, and Roncs Ese-etame</i>	
<b>An Assessment of Hardwood Lumber Markets in China</b> .....	<b>460</b>
<i>Scott A. Bowe, Matthew S. Bumgardner, and Xiping Wang</i>	
<b>Expanding Forest Management to Include Management of Nontimber Forest Resources</b> .....	<b>470</b>
<i>James L. Chamberlain</i>	

<b>Product Recovery from Tree Grade 1 Northern Red Oak on Menominee Tribal Lands.....</b>	<b>478</b>
<i>John P. Dwyer and Daniel C. Dey</i>	
<b>Can Smaller Diameter Hardwood Logs be Profitably Sawed into Lumber? .....</b>	<b>485</b>
<i>Matthew S. Scholl, Janice K. Wiedenbeck, and Paul R. Blankenhorn</i>	
<b>Using External High-Resolution Log Scanning to Determine Internal Defect Characteristics .....</b>	<b>497</b>
<i>Ed Thomas, Liya Thomas, Clifford Shaffer, and Lamine Mili</i>	
<b>Hardwood Log Merchandising and Bucking Practices in West Virginia .....</b>	<b>506</b>
<i>J. Wang, S. Grushecky, Y. Li, and J. McNeel</i>	
<b>Acoustic Assessment of Stress Level and Potential Wood Quality of Logs Affected by Oak Decline.....</b>	<b>513</b>
<i>Xiping Wang, Henry E. Stelzer, Jan Wiedenbeck, and Robert J. Ross</i>	
<b>NATURAL REGENERATION.....</b>	<b>525</b>
<b>Twenty-Two Year Changes in Regeneration Potential in an Old-Growth <i>Quercus</i> Forest on the Mid-Cumberland Plateau, Tennessee .....</b>	<b>527</b>
<i>Stacy L. Clark, Scott J. Torreano, David L. Loftis, and Luben D. Dimov</i>	
<b>Natural Oak Regeneration Following Clearcutting on the Hoosier National Forest.....</b>	<b>536</b>
<i>Robert C. Morrissey, John R. Seifert, Douglass F. Jacobs, John A. Kershaw, Jr., and Marcus F. Selig</i>	
<b>Oak Regeneration Response to Moderate and Heavy Traffic under Mechanical Harvesting in an Oak-Hickory Forest on the Cumberland Plateau.....</b>	<b>547</b>
<i>Callie Jo Schweitzer</i>	
<b>Fifteen Years of Stump Sprout Development for Five Oak Species in Southern Indiana.....</b>	<b>553</b>
<i>Dale R. Weigel and Daniel C. Dey</i>	
<b>ARTIFICIAL REGENERATION.....</b>	<b>561</b>
<b>A Comparison of the 36-Year Performance of Artificial and Natural Oak Regeneration in the Ridge and Valley Province of Eastern Tennessee .....</b>	<b>563</b>
<i>Samuel W. Jackson</i>	
<b>Evaluating the Flood Tolerance of Bottomland Hardwood Artificial Reproduction.....</b>	<b>572</b>
<i>John M. Kabrick, Daniel C. Dey, and Jonathan R. Motsinger</i>	
<b>Twenty-Four Years of Growth of Naturally Regenerated Hardwoods, Planted Yellow- Poplar, and Planted Pine in Plots with and without Competition Control on an Upland Hardwood Site on the Cumberland Plateau Near Sewanee, TN .....</b>	<b>581</b>
<i>Karen Kuers</i>	
<b>Deployment of High-Quality Oak Seedlings from Local Seed Sources Along Elevational Gradients in West Tennessee Bottomlands.....</b>	<b>591</b>
<i>Jason S. Maxedon, Scott E. Schlarbaum, and Donald G. Hodges</i>	
<b>Nitrate Reductase Activity in 1+0 <i>Juglans Nigra</i> Seedlings with N Fertilization.....</b>	<b>598</b>
<i>M.A. Nicodemus, K.F. Salifu, and D.F. Jacobs</i>	
<b>Fifteen-Year Performance of Five Oak Species in Plantation Culture.....</b>	<b>605</b>
<i>Randall J. Rousseau and Terry L. Robison</i>	

<b>Deer Browsing Patterns in a Recently Afforested Bottomland.....</b>	<b>612</b>
<i>Kenneth J. Ruzicka, John W. Groninger, and James J. Zaczek</i>	
<b>Exponential Nutrient Loading and Retranslocation Response of <i>Quercus Rubra</i> Seedlings.....</b>	<b>618</b>
<i>K. Francis Salifu, Douglass F. Jacobs, and Z. Birge</i>	
<b>ECOLOGY AND FOREST DYNAMICS .....</b>	<b>627</b>
<b>Genotypic Variation in Flood Tolerance of Black Walnut and Three Southern Bottomland Oaks .....</b>	<b>629</b>
<i>Mark V. Coggeshall, J.W. Van Sambeek, and Scott E. Schlarbaum</i>	
<b>Effects of Shade on the Growth of Natural and Artificially Established White Oak (<i>Quercus Alba</i> L.) Regeneration.....</b>	<b>638</b>
<i>Dylan Dillaway and Jeff Stringer</i>	
<b>Overstory and Regeneration Structure and Relationships in Mixed Stands on the Southern Cumberland Plateau .....</b>	<b>644</b>
<i>Luben D. Dimov and Callie Jo Schweitzer</i>	
<b>Culm Production and Morphology of Fresh and Stored Rhizomes from Field-Planted and Wild Giant Cane.....</b>	<b>652</b>
<i>John L. Hartleb and James J. Zaczek</i>	
<b>Changes in Tree Species Importance Following Harvesting Disturbance in North Mississippi Between 1967 and 1994 .....</b>	<b>658</b>
<i>Andrew J. Hartsell and James F. Rosson, Jr.</i>	
<b>A High Resolution Laser-Based Technique for Quantifying the Elemental Composition of Wood: Applications in Forest Fire Ecological Response.....</b>	<b>668</b>
<i>Madhavi Z. Martin, Nicole Labbé, Stan D. Wullschleger, Nicolas André, and Timothy G. Rials</i>	
<b>Plant Composition in Oak Savanna and Woodland Restoration at Prairie Fork Conservation Area in Missouri.....</b>	<b>674</b>
<i>Nadia E. Navarrete-Tindall, J.W. Van Sambeek, Jamie Coe, and Warren Taylor</i>	
<b>Dynamics of a Bottomland Hardwood-Pine Stand in Greene County, Tennessee.....</b>	<b>686</b>
<i>Matthew G. Olson and P. Daniel Cassidy</i>	
<b>Natural History from Dendrochronology: Maximum Ages and Canopy Persistence of Rarely Studied Hardwood Species.....</b>	<b>695</b>
<i>Neil Pederson, Anthony W. D'Amato, and David A. Orwig</i>	
<b>Effects of Lime, Fertilizer, and Herbicide on Herbaceous Species Diversity and Abundance following Red Oak Shelterwood Harvest.....</b>	<b>702</b>
<i>William E. Sharpe and Chad R. Voorhees</i>	
<b>Plant Communities Associated with Multi-Aged Clearcuts in the Missouri Ozarks .....</b>	<b>709</b>
<i>Irene M. Unger, Rose-Marie Muzika, and Nevin Aspinwall</i>	
<b>Land-Use History and Resulting Forest Succession in the Illinois Ozark Hills .....</b>	<b>719</b>
<i>Saskia L. van de Gevel and Charles M. Ruffner</i>	
<b>Developing a Field Facility for Evaluating Flood Tolerance of Hardwood Seedlings and Understory Ground Covers.....</b>	<b>727</b>
<i>J.W. Van Sambeek, Robert L. McGraw, John M. Kabrick, Mark V. Coggeshall, Irene M. Unger, and Daniel C. Dey</i>	

<b>POSTERS.....</b>	<b>735</b>
<b>Spatial Allocation of West Virginia Timber Product Output Data .....</b>	<b>737</b>
<i>John P. Brown</i>	
<b>A Case Study Assessment of Small-Diameter Utilization in the Upper Midwest .....</b>	<b>738</b>
<i>Matthew S. Bumgardner and Scott A. Bowe</i>	
<b>The Role of the WVU Extension Service in Forestry Education and Technical Assistance for Private Forestland Owners .....</b>	<b>739</b>
<i>Larry G. Campbell, David W. McGill, Chad Pierskalla, and Kevin Saunders</i>	
<b>Stem Diameter and Horizontal Crown Area Correlations for Hardwood Tree Seedlings Planted on Reclaimed Strip-Mined Lands in Eastern Kentucky .....</b>	<b>740</b>
<i>Lucas R. Cecil and Jeffrey Stringer</i>	
<b>Individual-Tree, Outside-Bark, Merchantable Green Weight Equations and Scaling Factors for Sawtimber-Sized Northern Red Oak, White Oak, and Sweetgum in Northwest Arkansas.....</b>	<b>741</b>
<i>Paul F. Doruska, Jonathan I. Hartley, Matthew B. Hurd, David W. Patterson, and Don C. Bragg</i>	
<b>Impact of Channelization and Dam Construction on Kaskaskia River Morphology.....</b>	<b>742</b>
<i>Xizhen Du and Karl W. J. Williard</i>	
<b>Soil Amendment Effects on Oak Seedlings and Woody Competitors.....</b>	<b>743</b>
<i>Jennifer Franklin and Richard Evans</i>	
<b>Relating Land-Use Practices to Sediment Loads in West Virginia's Upper Elk River Watershed .....</b>	<b>744</b>
<i>Jennifer B. Fulton, J. Todd Petty, Steven E. Harouff, Kyle J. Hartman, David W. McGill, and Shawn T. Grushecky</i>	
<b>Composition and Structure of an Old-Growth White Oak Forest in Transition .....</b>	<b>745</b>
<i>P. Charles Goebel, D.M. Hix, Kathryn L. Holmes, Marie E. Semko-Duncan, and C.E. Dygert</i>	
<b>Increased Use of Low-Quality Wood in the Upland Hardwood Region of North America: Can We Utilize More Oak in Oriented Strand Board? .....</b>	<b>746</b>
<i>Jody D. Gray, Joseph F. McNeel, and John R. Noffsinger</i>	
<b>Variation among Years for Mast Production by Oaks in Missouri.....</b>	<b>748</b>
<i>David P. Gwaze</i>	
<b>The Encyclopedia of Southern Appalachian Forest Ecosystems (ESAFE) .....</b>	<b>750</b>
<i>William Hubbard, Daniel Cassidy, and H. Michael Rauscher</i>	
<b>Effects of <i>Microstegium Vimineum</i>, an Invasive C<sub>4</sub> Grass, on Hardwood Regeneration.....</b>	<b>751</b>
<i>Rochelle R. Jacques and Brian C. McCarthy</i>	
<b>Effects of Controlled Burning and Shelterwood Thinning on Oak Mast Production in Two Southeastern Ohio Forests .....</b>	<b>752</b>
<i>Jeffrey A. Lombardo and Brian C. McCarthy</i>	
<b>Evaluation and Collection of Superior Black Cherry Trees in the Allegheny National Forest.....</b>	<b>753</b>
<i>James McKenna and Keith Woeste</i>	
<b>Evaluation of Tree Species Composition as a Tool for Classifying Moisture Regimes in Oak Forests of Eastern Kentucky.....</b>	<b>754</b>
<i>W. Henry McNab, David L. Loftis, Mary A. Arthur, and Jessi E. Lyons</i>	

<b>Carbon Sequestration and Enhanced Wildlife Habitat Resulting from Bottomland Hardwood Afforestation Activities in the Lower Mississippi Alluvial Valley.....</b>	<b>755</b>
<i>Richard P. Maiers, Andrew J. Londo, Donald L. Grebner, Jeanne C. Jones, Changyou Sun, Michael S. Cox, Jarod H. Fogarty, and Janet C. Dewey</i>	
<b>Survey of West Virginia Forestry Consultants: Services Provided and Fees Charged to the Private Forest Land Owners in the State of West Virginia.....</b>	<b>757</b>
<i>Dheeraj Nelli, David W. McGill, Kathryn G. Arano, and Shawn T. Grushecky</i>	
<b>Fire History of a Southern Illinois Bottomland Forest.....</b>	<b>758</b>
<i>John L. Nelson, Charles M. Ruffner, and John W. Groninger</i>	
<b>The Successional Status of Two Table Mountain Pine (<i>Pinus pungens</i>) Stands in the Southern Appalachians, Tennessee.....</b>	<b>759</b>
<i>Christopher M. Oswald, Wayne K. Clatterbuck, and Brian T. Hemel</i>	
<b>Response of the Non-Native Invasive Grass, <i>Microstegium Vimineum</i> (Trin.) A. Camus, to Three Levels of Canopy Disturbance.....</b>	<b>760</b>
<i>Christopher M. Oswald, Sonja N. Oswald, and Wayne K. Clatterbuck</i>	
<b>Correlations between Tree Crown Condition and Shade Tolerance, Crown Form, and Light Availability.....</b>	<b>761</b>
<i>KaDonna C. Randolph</i>	
<b>Evaluating the Distribution and Shade-Tolerance of Hay-Scented Fern Across a Light Gradient.....</b>	<b>762</b>
<i>Alejandro A. Royo and Walter P. Carson</i>	
<b>Natural Resource Interpretive Programs: An Evaluation.....</b>	<b>764</b>
<i>A.J. Stegmann and B.E. Cutter</i>	
<b>Use of Native Seed Mixtures to Improve Erosion Control and Wildlife Habitat on Log Landings following Timber Harvest in the Upper Elk Watershed of West Virginia.....</b>	<b>765</b>
<i>Lisa R. Tager, Shawn Grushecky, David W. McGill, William Grafton, and John Edwards</i>	
<b>Influence of Iron Industry Charcoal Production on Forest Composition and Structure on a Western Highland Rim Forest, Tennessee.....</b>	<b>766</b>
<i>Saskia L. van de Gevel, Justin L. Hart, David F. Mann, and Wayne K. Clatterbuck</i>	
<b>Value Loss Rate for Hardwood Trees Uprooted in a Severe Windstorm on the Allegheny Plateau.....</b>	<b>767</b>
<i>Janice K. Wiedenbeck and Susan Stout</i>	
<b>Utilization Options for Decadent Eastern Hemlock Timber.....</b>	<b>768</b>
<i>Matthew F. Winn and Philip A. Araman</i>	
<b>Groundwater Nitrogen and Phosphorus Dynamics in Giant Cane and Deciduous Forest Riparian Buffers.....</b>	<b>769</b>
<i>Chad M. Yocum, Karl W.J. Williard, Sara G. Baer, and James J. Zaczek</i>	
<b>Survival and Growth of Northern Red Oak Planting Stock Types through 17 Years after Planting.....</b>	<b>770</b>
<i>James J. Zaczek, Kim C. Steiner, and Tim Phelps</i>	

# NATURAL HISTORY FROM DENDROCHRONOLOGY: MAXIMUM AGES AND CANOPY PERSISTENCE OF RARELY STUDIED HARDWOOD SPECIES

Neil Pederson, Anthony W. D'Amato, and David A. Orwig<sup>1</sup>

**Abstract**—Tree-ring research has made significant contributions to our understanding of environmental change and forest stand dynamics. Its application to understanding natural history, however, has been limited. Biodiversity of the central hardwood forest offers many opportunities for tree-ring based, natural history research. Recent tree-ring research examining several rarely studied hardwood species has yielded ages well beyond maximum expectations. For example, a sampling of 20 *Magnolia acuminata* trees in one population included two individuals 315 and 348 years, respectively, which are nearly two centuries more than the average life expectancy reported for this species. Also, research in recently discovered old-growth stands in western Massachusetts has illustrated the common occurrence of *Betula lenta* in *Tsuga canadensis* dominated old-growth forests with individuals frequently living beyond 320 years in these systems. These studies illustrate that tree-ring research can expand our knowledge of the natural history of central hardwood species.

---

## INTRODUCTION

The science of dendrochronology (tree-ring analysis) has enhanced our understanding of environmental change, succession, and forest stand dynamics. In the eastern U.S., this type of research has been substantial and rich (i.e., Lorimer 1980, Foster 1988, Canham 1990, Runkle 1990, Stahle and Chaney 1994, Nowacki and Abrams 1997, Orwig and others 2001, Shumway and others 2001, Lafon and Speer 2002). In recent years, however, dendrochronology has been less frequently applied towards the understanding of natural history (cf. Dayton 2003). This type of natural history information may be more important now than ever before (Dayton, 2003; Schmidly 2005) as species and ecosystems are threatened by invasive species, land-use (development/urban sprawl), forest fragmentation and future climate change. The term 'natural history', unfortunately, is a rather nebulous term (Schmidly, 2005). In this paper we will focus on the application of dendrochronology towards determining a species' longevity and its persistence in the forest. We posit the idea, however, that dendrochronology can reveal many aspects of a species' life-history traits, which is an important part of its natural history.

Tree-ring studies in old-growth forests are valuable sources of information regarding the natural history of central hardwood species (cf. Rentch and others 2003a,b), as well as the natural disturbance dynamics, and development patterns for forest types throughout the region. This paper will highlight specific examples of new information concerning the natural history of four species. This information is derived from two recent dendrochronological studies in old-growth forests (Pederson 2005, D'Amato and others in review). The purpose of these studies was to identify the climatic sensitivity and growth history of several species in the eastern U.S. (Pederson 2005) and to estimate the amount of old-growth forest in Massachusetts (D'Amato and others in review). And yet, the ease at which new maximum ages were found highlights the fact that there are significant gaps in the natural history of many species characteristics of the central hardwood forest. Our hope is that this paper will stimulate additional research that will enrich our knowledge for many eastern U.S. species.

The first portion of this paper will focus on the longevity of four rarely studied temperate hardwood species: black birch (*Betula lenta* L.), cucumbertree (*Magnolia acuminata* L.), red maple (*Acer rubrum* L.), and shagbark hickory (*Carya ovata* (Mill.) K. Koch). New data and a review of maximum ages found

---

<sup>1</sup> Neil Pederson, Assistant Professor, Department of Biological Sciences, Moore 235, Eastern Kentucky University, Richmond, KY 40475; Anthony W. D'Amato, Graduate Research Assistant, University of Massachusetts, Holdsworth Natural Resource Center, Amherst, MA 01003-9285; and David A. Orwig, Forest Ecologist, Harvard Forest, Petersham, MA 01366.

in primary literature sources will be compared to illustrate how the ages found in our research are well beyond common maximum age expectations. There is currently no information available for any of these species in the International Tree-Ring Databank, a storehouse of dendrochronological information on hundreds of species worldwide (ITRDB 2005).

The second portion of this paper will focus on the natural history (recruitment, longevity and persistence) of black birch within eastern hemlock-mixed hardwood dominated old-growth forests in Massachusetts. Black birch is commonly reported in association with eastern hemlock (*Tsuga canadensis* (L.) Carr.) in old-growth forest ecosystems throughout the central hardwood and New England regions (e.g., Hough and Forbes 1943, Foster 1988); however, little is known about the natural history and dynamics of this species in old-growth forests. Data from eleven old-growth eastern hemlock-mixed hardwood forests in western Massachusetts are used to illustrate the surprising persistence of black birch as well as its overlooked importance in the structure and dynamics of old-growth eastern hemlock forests.

## METHODS

Increment cores were collected and processed using standard tree-ring analysis techniques (Cook and Kairiukstis 1990). Generally, a minimum of twelve trees in a stand was cored, with one to two cores removed from each tree depending on the goals of the study (Pederson 2005, D'Amato and others in review). Twenty *M. acuminata* and twenty-one *C. ovata* trees representing the perceived range of age classes from a stand in George Washington National Forest in the Blue Ridge Mountains of central Virginia were selected for coring (Pederson 2005). Twenty *A. rubrum* trees in the eastern Catskills, NY equal to or greater than 10 cm dbh within two designated stands were randomly selected for coring (Charles Canham, 65 Sharon Turnpike, P.O. Box AB, Millbrook NY 12545-0129 & Paul Sheppard, 105-C1 West Stadium, Tucson, AZ 85721 USA personal communication). *B. lenta* outside of Massachusetts were sampled over two periods. In 1974, twelve individuals were sampled at the Mohonk Preserve in New Paltz, NY (Dr. Edward Cook, unpublished data). A second collection of 17 individuals was made at the Preserve in a different stand in 2002 (Neil Pederson, unpublished data). Finally, all trees equal to or greater than 10 cm dbh falling within 3-5 400 m<sup>2</sup> plots were sampled in the study examining old-growth *B. lenta* in Massachusetts (D'Amato and others in review).

Cores were glued to wooden core mounts and, in most cases, progressively sanded up to 600-grit sandpaper. *B. lenta*, *M. acuminata* and *A. rubrum* samples were often sanded using 2400 or 3200 grit sandpaper to ensure ring boundaries were visually distinct in these species with diffuse porous ring structures. Samples of *B. lenta*, *M. acuminata* and *A. rubrum* were not stained or enhanced in any way beyond sanding. Ages presented here are derived from crossdated samples. Finally, ages presented here are minimum ages; no extrapolations have been made for the number of missing rings to the center of the tree or the time it took each tree to reach coring height. Therefore, the ages presented for these trees are certainly less than their absolute age.

## RESULTS AND DISCUSSION

### Maximum Ages for *A. rubrum*, *B. lenta*, *M. acuminata*, and *C. ovata*

Maximum ages and comparisons for each species are summarized in table 1. Maximum age in a sample of 40 *A. rubrum* trees is 300 years. The next four oldest trees from this collection were 212, 132, 129 and 128 years old. Interestingly, this maximum age is 150 years greater than the maximum age listed for this species in Loehle (1988) and 10 years older than the oldest reported in the early Pennsylvania study by Hough and Forbes (1943). The oldest *Betula lenta* tree in a sample of 29 trees is 361 years, while the next four oldest trees were 318, 257, 169 and 166 years old. Similar to the *A. rubrum* finding, the oldest *B. lenta* in the population was far greater than the maximum age listed for this species in the current USDA Silvics manual (Burns and Honkala 1990) and Hough and Forbes (1943) (table 1). Further discussion of the age structure of *B. lenta* will be presented in the next section. The oldest *C. ovata* in a sample of 20 trees is 354 years old with the next four oldest trees in the population being 257, 255, 254 and 251 years old. The oldest individual tree is 54 years greater than the maximum age for this species listed in Loehle

**Table 1—Comparison of maximum ages from the *Silvics of North America* (Burns and Honkala 1990), Hough and Forbes (1943), and recently acquired data**

Species	Silvics manual <sup>a</sup>	Hough and Forbes	New data	Difference <sup>b</sup>
	----- years -----			
<i>Acer rubrum</i>	150	290	300 <sup>c</sup>	150
<i>Betula lenta</i>	265	265	361 <sup>d</sup>	96
<i>Carya ovata</i>	300	n/a	354 <sup>e</sup>	54
<i>Magnolia acuminata</i>	150	310	348 <sup>e</sup>	198

<sup>a</sup> If no age is given in the Silvics manual, ages are substituted from Loehle (1988) or Hough and Forbes (1943).

<sup>b</sup> Difference between new maximum age data and Silvics manual, or if no age is given, Loehle (1988) or Hough and Forbes (1943).

<sup>c</sup> Catskill Mountains, NY.

<sup>d</sup> New Paltz, NY.

<sup>e</sup> George Washington National Forest, VA.

(1988). Finally, the oldest *M. acuminata* in a sample of 20 trees is 348 years old with the next four oldest trees in the population obtaining ages of 318, 215, 177 and 174 years old. The oldest individual tree is 198 years greater than the maximum age for this species listed in Burns and Honkala (1990) and 38 years greater than in Hough and Forbes (1943).

The frequency at which greater ages are found in our studies compared to the Silvics Manual of North America (Burns and Honkala 1990) or Loehle's (1988) list of known maximum longevity suggests that there is much yet to learn about the basic natural history of central hardwood and eastern US forests. Further, with the exception of *A. rubrum*, the proximity of the next oldest individuals in the recent studies either to the new or previous maximum age suggests that longevity in these species may be even greater than those reported here. This hypothesis is built upon the premise that maximum ages for each species has a normal distribution; a 'common' maximum age. It could be, however, that a species' maximum age could be significantly larger than what is reported if maximum age has an asymmetrical distribution or a long tail (*sensu* Clark and others 1998). Such a distribution could explain the 800+ year maximum age reported for eastern hemlock (*T. canadensis*) (Burns and Honkala 1990), despite the fact that a collection of more than 1000 *T. canadensis* collected from across its range has not yielded an individual greater than 600 years old (ITRDB 2005).

The similarity of the new maximum ages presented here and those ages reported in the classic study of the high plateau region of Pennsylvania by Hough and Forbes (1943) indicates that these maximum ages may represent a common maximum age. It should be noted, however, that the ages of Hough and Forbes (1943) are ring counts of stumps in the field and could be significantly off due to measurement error. Likewise, a lack of dating control might explain the extreme maximum age reported for *T. canadensis*. Crossdated samples of *A. rubrum* indicate the possibility of up to 12 missing rings per tree in extreme cases (Pederson 2005). Similarly, work on *M. Acuminata* and *B. lenta* indicates that these species can have a significant number (greater than 10) of false and missing rings (N. Pederson, unpublished data). Nonetheless, the near agreement of the Pennsylvania ages and recent studies suggest that the work of Hough and Forbes (1943) provides an excellent early estimate of the maximum ages for many of these species.



### Black Birch in Old-Growth Eastern Hemlock Forests

Overall, black birch was a minor component of the old-growth stands investigated in Massachusetts, making up less than 15 percent of the overstory species composition (table 2). Within these stands, black birch was most commonly found in the intermediate and codominant crown classes and attained diameters smaller than the associated hemlock (table 2, mean hemlock diameter = 33.2 cm). Maximum ages of black birch individuals within these stands exceeded those previously reported for this species (Hough and Forbes 1943), including several individuals between 320-332 years old (table 2). In addition, only two of the eleven stands investigated did not contain at least one black birch individual greater than 210 years old (table 2).

Black birch recruitment generally occurred in episodic peaks with other species within these stands (fig. 1). These episodic recruitment patterns indicate that successful birch recruitment occurred predominantly during large disturbance events, such as the hurricane of 1893 (fig. 1). These findings are consistent with other studies that have also demonstrated the importance of moderate disturbance events in facilitating the establishment of black birch (Ward and Stephens 1996). While these recruitment events lead to an increase in the amount of black birch, the age data collected from these sites suggests that several older trees may also have become established from smaller unknown events.

Although other studies have previously reported the presence of *B. lenta* in old-growth forests (Hough and Forbes 1943, Morey 1936, Foster 1988, Orwig and others 2001), this species has traditionally been thought of as an associate of younger forest ecosystems (e.g., Stephens and Waggoner 1970, Trimble 1970). The findings from our research indicate that black birch commonly plays a prominent role in the structure and dynamics of old-growth hemlock stands attaining ages well beyond previous expectations. In the populations examined in this study, most *B. lenta* were less than 150 years old; however, 58 percent of those individuals greater than 180 years old have lived beyond 250 years illustrating the ability of this species to consistently persist within the canopy of these old-growth stands. Interestingly, *B. lenta* is often the species replacing eastern hemlock in stands infested with hemlock woolly adelgid in southern New

**Table 2—Summary table for attributes of black birch populations in 11 old-growth mixed eastern hemlock stands in western Massachusetts**

Study area	Importance value <sup>a</sup>	Age <sup>b</sup>	Diameter cm
Mt. Everett	9.4	65 (77)	18.2
Grinder Brook	5.7	175 (218)	27.3
Bash Bish Falls	2.8	172 (211)	28.3
Cold River A1	11.0	127 (251)	28.9
Cold River A2	9.0	169 (326)	30.3
Cold River B	9.0	146 (238)	32.0
Cold River D	8.6	165 (261)	21.4
Manning Brook	7.2	87 (158)	24.0
Wheeler Brook	10.4	203 (284)	27.5
Black Brook	12.5	182 (328)	30.3
Todd Mountain	6.9	163 (332)	28.4
Average	8.4	150	27.0

<sup>a</sup> Importance value = (relative basal area + relative density)/2.

<sup>b</sup> Average age with maximum found at site in parentheses.




Figure 1—Example age structures from two eastern hemlock-mixed hardwood dominated old-growth forests, (A) Cold River A2 and (B) Todd Mountain, in western Massachusetts. N = 74 (13 *B. lenta*) and 88 (9 *B. lenta*) for (A) and (B), respectively.

England (Orwig and Foster 1998). In light of our findings, it is likely that *B. lenta* may persist in these affected stands longer than previously expected.

## CONCLUSIONS

Data presented here suggests that rarely studied trees species can live much longer than previously thought. Our results also show how dendrochronology is an excellent tool to deepen our knowledge of the natural history of central hardwood tree species. Besides maximum age, tree-ring analysis can reveal other facets of natural history such as how long a species can remain suppressed in the understory, how they respond to disturbance events or how long they can sustain high levels of productivity. Not only will this dendrochronology-derived information help expand our knowledge of natural history, it will provide

important data for simulation models to produce realistic estimations of tree longevity. Future studies of other central hardwood species will be critical for future efforts aimed at modeling long-term forest dynamics, as well as for predicting the population and system responses to the effects of environmental changes and novel disturbances such as the hemlock woolly adelgid, on future forest structure and composition (Dayton 2003; Schmidly, 2005).

The ‘accidental’ discovery of new maximum ages for four species typical of central hardwood forests and the surprising persistence of *B. lenta* suggests the lack of natural history knowledge for many eastern U.S. tree species. We hope this presentation emphasizes the need for more research. Such research would be an excellent avenue for inspiring motivated undergraduate and graduate-level students to incorporate the field of natural history into their studies.

## ACKNOWLEDGMENTS

N. Pederson greatly acknowledges support of the Department of Energy’s Global Change Education Program for support during his dissertation research. Many thanks to Drs. Charles Canham and Paul Sheppard for collecting and sharing the *A. rubrum* samples and Ed Cook for collecting and sharing the New Paltz *B. lenta* samples. This is Lamont-Doherty Earth Observatory Contribution Number 6869.

## LITERATURE CITED

- Burns, R.M. and B.H. Honkala, (eds.). 1990. Silvics of North America: Volume 2, Hardwoods. USDA Forest Service Agricultural Handbook 654. 877 p.
- Canham, C.D. 1990. Suppression and release during canopy recruitment in *Fagus grandifolia*. Bulletin of the Torrey Botanical Club 117: 1-7.
- Clark J.S., C. Fastie, G. Hurtt, and others. 1998. Reid’s Paradox of rapid plant migration. BioScience 48:13–24.
- Cook, E.R. and L.A. Kairiukstis (eds). 1990. Methods of Dendrochronology. Kluwer Academic Publications, Hingham, MA: 408 p.
- D’Amato, A.W., D.A. Orwig, and D.R. Foster. In review. New estimates of Massachusetts old-growth: a useful tool for forest reserve planning. Submitted to The Northeastern Naturalist.
- Dayton, P.K. 2003. The importance of the natural sciences to conservation. American Naturalist 162: 1-13.
- Foster, D.R. 1988. Disturbance history, community organization and vegetation dynamics of the old-growth Pisgah Forest, southwestern New Hampshire, U.S.A. Journal of Ecology 76: 105-134.
- Hough, A.F., and R.D. Forbes. 1943. The ecology and silvics of forests in the high plateaus of Pennsylvania. Ecological Monographs 13:301-320.
- ITRDB – International Tree-Ring Data Bank. 2005. Contributors of the International Tree-Ring Data Bank, IGBP PAGES/World Data Center for Paleoclimatology, NOAA/NCDC Paleoclimatology Program, Boulder, Colorado, USA. Web document: <http://www.ncdc.noaa.gov/paleo/treering.html>.
- Lafon, C.W. and J.H. Speer. 2002. Using dendrochronology to identify major ice storm events in oak forests of southwestern Virginia. Climate Research 20: 41-54.
- Loehle, C. 1988. Tree life history strategies: the role of defenses. Canadian Journal of Forest Research 18: 209-222.
- Lorimer, C.G. 1980. Age structure and disturbance history of a southern Appalachian virgin forest. Ecology 61: 1169-1184.
- Morey, H.F. 1936. Age-size relationships of Heart’s Content, a virgin forest in northwestern Pennsylvania. Ecology 17: 251-257.
- Nowacki, G.J. and M.D. Abrams. 1997. Radial-growth averaging criteria for reconstructing disturbance histories from presettlement-origin oaks. Ecological Monographs 67: 225-249.
- Orwig, D.A. and D.R. Foster. 1998. Forest response to the introduced hemlock woolly adelgid in southern New England, USA. Journal of the Torrey Botanical Society 125: 59-72.

- Orwig, D.A., C.V. Cogbill, D.R. Foster, and J.F. O'Keefe. 2001. Variations in old-growth structure and definitions: Development and dynamics of forests on Wachusett Mountain, Massachusetts. *Ecological Applications* 11:437-452.
- Pederson, N. 2005. Climatic Sensitivity and Growth of Southern Temperate Trees in the Eastern US: Implications for the Carbon Cycle. Ph.D. Thesis. Columbia University, NY.
- Rentch, J.S., M.A. Fajvan, and R.R. Hicks, Jr. 2003a. Spatial and temporal disturbance characteristics of oak-dominated old-growth stands in the central hardwood forest region. *Forest Science* 49: 778-789.
- Rentch, J.S., M.A. Fajvan, and R.R. Hicks, Jr. 2003b. Oak establishment and canopy accession strategies in five old-growth stands in the central hardwood forest region. *Forest Ecology and Management* 184: 285-297.
- Runkle, J.R. 1990. Gap dynamics in an Ohio *Acer-Fagus* forest and speculations on the geography of disturbance. *Canadian Journal of Forest Research* 20: 632-641.
- Schmidly, D.J. 2005. What it means to be a naturalist and the future of natural history at American universities. *Journal of Mammalogy* 86(3): 449-456.
- Shumway, D.L., M.D. Abrams, and C.M. Ruffner. A 400 year history of fire and oak recruitment in an old-growth forest in western Maryland, U.S.A. *Canadian Journal of Forest Research* 31: 1437-1443.
- Stahle, D.W. and P.L. Chaney, 1994. A predictive model for the location of ancient forests. *Natural Areas Journal* 14: 151-158.
- Stephens, G.R., and P.E. Waggoner. 1970. The forests anticipated from 40 years of natural transitions in mixed hardwoods. Connecticut Agricultural Experiment Station, Bulletin 707. New Haven. 58 p.
- Trimble, G.R., Jr. 1970. 20 years of intensive even-aged management: effect on growth, yield, and species composition in two hardwood stands in West Virginia. USDA Forest Service, Research Paper NE-154. Northeastern Forest Experiment Station, Broomall, PA. 12 p.
- Ward, J.S., and Stephens, G.R. 1996. Influence of crown class on survival and development of *Betula lenta* in Connecticut, U.S.A. *Canadian Journal of Forest Research* 26: 277-288.