

Turner-Peterson, C. E., Olsen, P. E., Nuccio, V. F., 1985, Modes of uranium occurrence in the Newark Basin, New Jersey and Pennsylvania: U.S. Geological Survey Circular 946, p. 120-124.

24. MODES OF URANIUM OCCURRENCE IN BLACK MUDSTONES IN THE NEWARK BASIN, NEW JERSEY AND PENNSYLVANIA

Christine E. Turner-Peterson, P.E. Olsen, and Vito F. Nuccio

INTRODUCTION

The Newark basin in eastern Pennsylvania, central New Jersey, and southern New York is an early Mesozoic rift-related basin that formed in Late Triassic time and continued to receive sediment through Early Jurassic time (see fig. 24.1). Anomalous concentrations of uranium occur in sandstone of the Stockton Formation (Turner-Peterson, 1980) and mudstone of the Stockton and Lockatong Formations (fig. 24.2).

Mudstone-hosted uranium deposits, discussed in this report, occur as thin (less than 0.5 m) but laterally continuous uraniferous zones in certain types of black mudstone. The usual grade in these mineralized mudstones is 0.01–0.02 percent uranium oxide as U_3O_8 (Turner-Peterson, 1980); locally higher grades (as high as 0.29 percent) are known. These uranium values are higher than values for other uranium occurrences in black shale reported in the literature. Black-shale uranium occurrences in the Devonian and Mississippian Chattanooga Shale and the Alum Shale in Sweden average 0.006 and 0.03 percent, respectively (Davidson, 1961; Chase, 1979). In all the black-shale occurrences, uranium was probably fixed at or near the sediment-water interface, representing a nearly syngenetic concentration of uranium within reduced lake-bottom sediments.

NEW RESULTS

As a result of recent work, uranium mineralization in the Lockatong black mudstones can now be tied to portions of lacustrine cycles defined originally by Van

Houten (1964) and refined by Olsen (1980a, 1984). Highest uranium contents occur in division 2 (of the detrital short cycle of Olsen, 1984), which is characterized by fine, calcareous laminated black siltstone (fig. 24.3). High uranium content correlates with high organic carbon and high sulfide sulfur contents. The uranium is commonly, but not always, associated with conchostracans, coprolites, ostracodes, and fish fossils.

One of the uranium anomalies has been located at precisely the same stratigraphic interval in the Lockatong at three widely spaced localities spanning a lateral distance of approximately 60 km. The widespread lateral extent of these black-shale uranium occurrences in the Lockatong Formation suggests syndepositional mineralization. This feature may also be useful for stratigraphic correlation.

A newly discovered anomaly in the Stockton Formation is of particular interest because the uranium occurs in a shaley black mudstone containing lacustrine fossils that resembles many of the uranium-bearing black shales of the Lockatong. Abundant conchostracans occur in the mineralized interval, indicating a lacustrine environment for the mineralization. Organic-carbon contents range from 0.05 to 0.66 percent, and uranium achieves a maximum grade of 0.29 percent uranium oxide, which is ore grade (table 24.1).

DISCUSSION

Because uranium occurrences are commonly associated with organic material in both sandstone and shale, a discussion of the types of organic matter that occur in the Newark basin is of interest. In general,




FIGURE 24.1.—Generalized geologic map of Newark basin (modified from Glaeser, 1966). Units dip to the north and northwest toward the northern border fault, and the section is repeated by faulting near the Delaware River. The stratigraphic units shown are the Stockton (Trs), Lockatong (Trl), and Passaic (JTTrp) Formations and the Jurassic basalts and interbedded sedimentary rocks, shown here as Jurassic undifferentiated (Ju). Location 1 shows site of a newly discovered uranium anomaly in a gray mudstone of the Stockton Formation (see table 24.1). Cross section A-A' shown in figure 24.2.




FIGURE 24.2.—Stratigraphic section of the Newark Supergroup along the Delaware River from Stockton, N.J., northward (modified from Van Houten, 1969). Section measured along A-A' (fig. 24.1).

organic material that is characterized by numerous oxygen-bearing functional groups has a greater affinity for metals than do other types of organic material (Schnitzer and Khan, 1978). Humic substances, in particular, are known to be effective concentrators of uranium (Szalay, 1958), whereas hydrocarbons generally

are not (Andreyev and Chumachenko, 1964). New data indicate that many of the Jurassic lacustrine beds in the Newark basin have organic characteristics and thermal maturity making them suitable as hydrocarbon source beds (Olsen, 1984; Hatcher and Romankiw, chapter 11, this volume; Pratt and others, chapter 13,

this volume), but humic substance contents in these samples are low, and thus these beds are not favorable for uranium mineralization. Lacustrine beds in the Triassic Lockatong Formation also contain only small amounts of terrestrial plant material and thus are inferred not to contain much humic material. In fact, visual examination of the kerogen in typical black mudstone of division 2 in the Lockatong Formation shows that detrital plant debris makes up only a minor fraction of the total organic material present (Olsen, 1984). However, the presence of thin coaly beds and detrital plant fragments in other parts of the basin indicates that humic-substance-producing organic material (terrestrial plant debris) was available and present in the lake basin. Visual inspection of kerogen from the Stockton anomaly site during vitrinite reflectance studies showed that type 3 (terrestrial plant material deriv-

ative) is abundant in the samples enriched in uranium (table 24.1). Even though the organic matter in the Stockton anomaly is apparently rich in humic substances favorable for fixing uranium, it is unclear at this point whether the organic matter played an active role in the fixing of uranium. Fixation of uranium could occur by complexing with humic matter or by precipitation in response to reducing conditions near the sediment-water interface. The mechanism is of particular concern as the Lockatong uranium occurrences are in black mudstones that apparently do not contain large quantities of terrestrial plant debris, which contribute humic substances. The extremely high grade (0.29 percent uranium oxide) of the Stockton black mudstone, however, may reflect the influence of an unusual abundance of terrestrial plant material in this particular unit.




FIGURE 24.3.—Typical detrital cycle in the Lockatong Formation. Note that total organic carbon (TOC) is highest in division 2. Most uranium anomalies in this formation are associated with the lower part of division 2, which commonly contains coprolites, conchostracans, ostracodes, and fish remains. TOC values are from Olsen (1984). Fossil content is intended to be schematic and is summarized from Olsen (1984). Uranium and sulfur contents are also schematic.

TABLE 24.1.—Vertical section (120 cm) across zone of uraniferous mudstone in the Stockton Formation. Units generally 2.5–15 cm thick; sample 6 is from a 2.5-cm-thick unit. See figure 24.1 for location

[U₃O₈ in percent; determined by delayed neutron activation analysis. Organic carbon in percent; determined by difference (total organic carbon minus carbonate carbon). Vitrinite reflectance in percent; —, not determined]

Sample number	Lithology	U ₃ O ₈	Organic carbon	Vitrinite reflectance
13	Dark red siltstone	<0.01	0.02	—
12	Medium gray mudstone	<.01	.03	—
11	Medium gray mudstone	<.01	.02	—
10	Medium gray mudstone	<.01	.02	—
9	Medium gray sandstone	<.01	.03	—
8	Dark gray siltstone	<.01	<.01	—
7	Dark gray siltstone	<.01	.08*	—
6	Medium gray claystone	.29	.10**	—
5	Black mudstone	.04	.66**	0.68
4	Light gray claystone	.01	.05	—
3	Black mudstone	.01	.33	1.15
2	Light gray claystone	<.01	.10	—
1	Black mudstone	<.01	.23	.66

*Contains plant fragments. **Contains conchostracans.

CONCLUSIONS

Some Triassic black lacustrine mudstones of the Newark basin are rich in uranium, with contents commonly in the range of 0.01–0.02 weight percent uranium oxide. High uranium concentrations in these black mudstones correlate with high organic carbon and high total sulfur contents. This uranium enrichment is interpreted to be syndepositional and possibly related to precipitation in response to reducing conditions at the sediment-water interface. The occurrence of uranium enrichment at the same stratigraphic position in the Lockatong Formation at three widely separated localities supports this model.

A newly discovered uranium anomaly in the Stockton Formation is of particular interest because of the high uranium content (0.29 percent U₃O₈). High uranium concentrations occur in gray to black lacustrine mudstones with locally high contents of organic

matter, similar to the associations observed elsewhere. This site is distinctive in that the mineralized horizon contains a high proportion of terrestrial organic matter.

REFERENCES

- Andreyev, P.F., and Chumachenko, A.P., 1964, Reduction of uranium by natural organic substances: *Geochemical International*, v. 1, p. 3–7.
- Chase, C.K., 1979, The Chattanooga and other Devonian shales — the United States' largest uranium source; *Symposium Papers, Synthetic Fuels from Oil Shale*: Chicago, Illinois, Institute of Gas Technology, p. 547–578.
- Davidson, C.F., 1961, The kolm deposits of Sweden: *Mining Magazine*, v. 105, p. 201–207.
- Glaeser, J.D., 1966, Provenance, dispersal, and depositional environments of Triassic sediments in the Newark-Gettysburg basin: *Pennsylvania Geological Survey Bulletin G43*, 168 p.
- Olsen, P.E., 1980a, Fossil great lakes of the Newark Supergroup in New Jersey, in Manspeizer, Warren, ed., *Field studies in New Jersey geology and guide to field trips*, 52nd Annual Meeting, New York State Geological Association: Newark, N.J., Rutgers University Press, p. 352–398.
- , 1980b, The latest Triassic and Early Jurassic formations of the Newark basin (eastern North America, Newark Supergroup): *Stratigraphy, structure, and correlation*: New Jersey Academy of Science Bulletin 25, p. 25–51.
- , 1984, Comparative paleolimnology of the Newark Supergroup — A study of ecosystem evolution: New Haven, Conn., Yale University, unpub. Ph.D. dissertation, 726 p.
- Szalay, A., 1958, The significance of humus in the geochemical enrichment of uranium: *Proceedings of the Second International Conference on the Peaceful Uses of Atomic Energy*, Geneva, v. 2, p. 182–186.
- Schnitzer, M., and Khan, S.U., 1978, *Soil organic matter*: New York, Elsevier, 319 p.
- Turner-Peterson, C.E., 1980, Sedimentology and uranium mineralization in the Triassic-Jurassic Newark basin, in Turner-Peterson, C.E., ed., *Uranium in sedimentary rocks — Application of the facies concept to exploration*: Short Course Notes, Rocky Mountain Section of Society of Economic Paleontologists and Mineralogists, p. 149–175.
- Van Houten, F.B., 1964, Cyclic lacustrine sedimentation, Upper Triassic Lockatong Formation, central New Jersey and adjacent Pennsylvania: *Kansas Geological Survey Bulletin 169*, p. 497–531.