

**NEWARK SUPERGROUP, A REVISION  
OF THE NEWARK GROUP IN EASTERN NORTH AMERICA**By A. J. Froelich<sup>1</sup> and P. E. Olsen<sup>2</sup>

The Newark Supergroup, a name proposed herein for adoption by the U.S. Geological Survey, includes the largely continental clastic rocks ("red beds") and interbedded basaltic flow rocks of Late Triassic and Early Jurassic age that crop out in discrete elongate basins parallel to the Appalachian orogen in eastern North America (fig. 1). The term "Newark Supergroup" was introduced by Van Houten (1977), referring to an unpublished manuscript by Olsen, to replace "Newark Group" (Redfield, 1856), a term that had been widely used but frequently misapplied in a time-stratigraphic sense (Klein, 1962). The use of Newark Supergroup preserves a well-established name (North American Stratigraphic Code, art. 7: c) which has increasingly been applied outside the U.S. Geological Survey to the rocks in all of the exposed basins (Geological Society of America, 1983, p. 156). The Newark Supergroup is a formal assemblage of related groups and formations (North American Stratigraphic Code, art. 29) with close lithologic and structural relationships that are implied through use of the supergroup designation. The term was clearly redefined by Olsen in 1978, but was expanded to include subsurface red beds of early Mesozoic age beneath the Atlantic Coastal Plain and Continental Shelf. As these subsurface rocks are poorly understood and apparently of diverse age, lithology, and origin, the term Newark Supergroup is here restricted to rocks that crop out, although we recognize that coeval strata are certainly concealed at depth beneath the Coastal Plain.

The Newark Supergroup strata in the exposed basins of eastern North America have variously been considered to be partly or solely of Early Jurassic age (Rogers, 1842; Lyell, 1847; Redfield, 1856), Permo-triassic (Emmons, 1857), Jurassic or Late Triassic (Fontaine, 1883), then solely of Late Triassic age, at first based on rare vertebrate and plant fossils (Ward, 1891; Eastman, 1913) and subsequently on vertebrate and plant fossils (Reeside and others, 1957) and radiometric ages of intercalated igneous rocks (Armstrong and Besancon, 1970). Some of the basins, however, have been determined to contain Lower Jurassic as well as Upper Triassic strata, as evidenced by spores, pollen, and well-preserved vertebrate remains in lacustrine mudstones (Cornet and others, 1973; Cornet, 1977; Olsen, 1978; Olsen and others, 1982) interbedded with basalt flows. The Lower Jurassic flows and interbedded strata can be considered informally as the "upper" Newark Supergroup and the Upper Triassic rocks as the "lower" Newark Supergroup.

The basins with only Upper Triassic rocks (with Group names where used) are the Wadesboro-Sanford-Durham (Chatham Group of Emmons, 1857)(1, 2, 3 on fig. 1); Davie County(4); Dan River and Danville (Dan River Group of Thayer, 1970)(5); Scottsburg(6); basins north of Scottsburg basin(7); Farmville(8); Richmond (Tuckahoe and Chesterfield Groups of Shaler and Woodworth, 1899)(9); Taylorsville(10); Scottsville(11),

<sup>1</sup>U.S. Geological Survey.

<sup>2</sup>Yale University, New Haven, Conn.




FIGURE 1.—Exposed basins of the Newark Supergroup in eastern North America.

and Barbourville (Culpeper Group of Lindholm, 1979)(12). The basins where Upper Triassic rocks are overlain by Lower Jurassic rocks are: the Culpeper (Culpeper Group of Lindholm, 1979)(13); Gettysburg (Conewago Group of Ashley, 1931)(14); Newark(15); Pomperaug(16); Hartford with Cherry Valley outlier (Meriden Group of Krynine, 1950)(17); Deerfield(18); Fundy or Minas (Fundy Group of Klein, 1962)(19); and Chedabucto(20).

Older Mesozoic strata of the lower Newark Supergroup (Upper Triassic, middle and upper Carnian), which are commonly coal-bearing, are preserved in the southern basins (1-10, fig. 1). Strata in two small, centrally located basins (11, 12, fig. 1) are mainly conglomerates and red beds that apparently lack diagnostic fossils but resemble Upper Triassic (upper Carnian and middle and upper Norian) rocks in adjacent basins to the north. Strata from the northern basins contain intercalated basalt flows and younger strata of the upper Newark Supergroup (13-18, fig. 1), span Late Triassic (Carnian and Norian) through Early Jurassic (Hettangian to Toarcian) time, and in the Hartford Basin (17, fig. 1), perhaps extend into Middle Jurassic (Bajocian) time. In the extreme northeast, the Fundy (Minas) Basin (19, fig. 1) is anomalous to this regional pattern because it contains Upper and possibly Middle Triassic (Ladinian) strata at the base and Lower Jurassic strata and basalt flows of the upper Newark Supergroup at the top.

As Olsen (1978) pointed out: " \* \* \* Raising the rank of the term Newark to Supergroup preserves the original and familiar meaning of Redfield's designation, allows the formations of individual basins to be included in specific groups while remaining in a strictly rock-stratigraphic hierarchy, and permits the maximum amount of flexibility for future subdivision. \* \* \*"

#### REFERENCES CITED

- Armstrong, R. L., and Besancon, J., 1970, A Triassic time scale dilemma: K-Ar dating of Upper Triassic mafic igneous rocks, eastern U.S.A. and Canada and post-Triassic plutons, western Idaho, U.S.A.: *Eclogae Geologicae Helveticae*, v. 63, p. 15-28.
- Ashley, G. H., 1931, A syllabus of Pennsylvania geology and mineral resources: Pennsylvania Topographic and Geologic Survey, 4th Series, Bulletin no. G-1, 159 p.
- Calver, J. L., 1963, Geologic map of Virginia: Virginia Department Conservation and Economic Development, Charlottesville, Scale 1:500,000.
- Cornet, Bruce, 1977, The palynostratigraphy and age of the Newark Supergroup: Unpublished Ph.D Thesis, Pennsylvania State University, 506 p.
- Cornet, Bruce, Traverse, Alfred, and McDonald, N. G., 1973, Fossil spores, pollen and fishes from Connecticut indicate Early Jurassic age for part of the Newark Group: *Science*, v. 182, p. 1243-1246.
- Eastman, C. R., 1913, Notes on Triassic fishes belonging to the Families Catopteridae and Semionotidae: *Annals of the Carnegie Museum*, v. 9, p. 139-148.
- Emmons, E. E., 1857, *American Geology* Pt. VI (III); X, 152, Albany.
- Fontaine, W. M., 1883, Contributions to the knowledge of the older Mesozoic flora of Virginia: U.S. Geological Survey Monograph 6, 144 p.

- Geological Society of America, 1983, Symposium: Advances in Paleontology and paleoecology—Newark Supergroup (Early Mesozoic): North-eastern Section, Abstracts with programs, v. 15, no. 3, p. 156.
- Hitchcock, Edward, 1832, Report on the geology of Massachusetts: American Journal of Science, v. 22, p. 1-70.
- Jansa, L. F., and Wade, J. A., 1975, Geology of the continental margin off Nova Scotia and Newfoundland in Volume 2, Offshore geology of Eastern Canada: Geological Survey of Canada Paper 74, 30, p. 51-105, Ottawa.
- King, P. B., and others, 1944, Tectonic map of the United States: National Research Council, Division of Geology and Geography, Committee on Tectonics, American Association of Petroleum Geologists, 2 sheets, scale 1:2,500,000.
- Klein, G. DeV., 1962, Triassic sedimentation, Maritime Provinces, Canada: Geological Society of America Bulletin, v. 73, p. 1127-1146.
- Krynine, P. D., 1950, Petrology, stratigraphy, and origin of the Triassic sedimentary rocks of Connecticut: Geological and Natural History Survey Bulletin no. 73, 247 p.
- Lindholm, R. C., 1979, Geologic history and stratigraphy of the Triassic-Jurassic Culpeper basin, Virginia: Geological Society of America Bulletin, (pt. 2, microfiche), v. 90, no. 11, p. 1702-1736.
- Lyell, Sir Charles, 1847, On the structure and probable age of the coal field of the James River near Richmond, Virginia: Quarterly Journal of the Geological Society of London, v. 3, p. 261-280.
- North American Commission on Stratigraphic Nomenclature, 1983, North American Stratigraphic Code: American Association of Petroleum Geologists Bulletin, v. 67, no. 5, p. 841-875.
- Olsen, P. E., 1978, On the use of the term Newark for Triassic and Early Jurassic rocks of eastern North America: Newsletters on Stratigraphy, v. 7, no. 2, p. 90-95.
- Olsen, P. E., McCune, A. R., Thomson, R. S., 1982, Correlation of the early Mesozoic Supergroup by vertebrates, principally fishes: American Journal of Science, v. 282, p. 1-44.
- Redfield, W. C., 1856, On the relations of the fossil fishes of the sandstone of Connecticut and the Atlantic States to the Liassic and Oolitic periods: American Journal of Science (ser. 2) 22, p. 357-363.
- Reeside, J. B., Jr., (Chairman, Triassic Subcommittee) and others, 1957, Correlation of Triassic formations of North America: Geological Society of America Bulletin, v. 68, p. 1451-1514.
- Rogers, W. B., 1842, Report of the progress of the Geological Survey of the State of Virginia for the year 1841: 12 p., Richmond, Va. (reprinted in Geology of the Virginias, 1884, p. 537-546.)
- Shaler, N. S., and Woodworth, J. B., 1899, Geology of the Richmond Basin, Virginia: U.S. Geological Survey Annual Report 19, 1897-1898, pt. 2, p. 385-515.
- Thayer, P. A., 1970, Stratigraphy and geology of the Dan River Triassic basin, North Carolina: Southeastern Geology, v. 12, p. 1-37.
- Van Houten, F. B., 1977, Triassic-Liassic deposits of Morocco and Eastern North America: Comparison: American Association of Petroleum Geologists Bulletin, v. 61, no. 1, p. 79-99.
- Ward, L. F., 1891, The plant-bearing deposits of the American Triassic: Science, v. 18, p. 287-288.