

PERIODICITY OF LAKE-LEVEL CYCLES IN THE LATE TRIASSIC LOCKATONG  
FORMATION OF THE NEWARK BASIN (NEWARK SUPERGROUP, NEW JERSEY AND  
PENNSYLVANIA)

P.E. Olsen<sup>1</sup>

Peabody Museum, Box 6666, 170 Whitney Avenue, New  
Haven, Connecticut 06511.

The short (ca. 5m) sedimentary cycles which Van Houten first described from the Lockatong Formation of the Newark Basin were produced by the climate controlled rise and fall of large lakes. During their high stands many of these lakes covered over 7000 km<sup>2</sup> and were over 100 m deep, but during their low stand they were reduced to playas or dried out completely. Annual laminations (varves) present in many short cycles show that the lakes which produced them oscillated in depth with a periodicity of about 21 800 years. Lockatong short cycles make up longer, compound cycles, 25 m and 100 m thick, which reflect periodic changes in the magnitude of high-stands of the 21 800 year lake-level cycles. These lower frequency cycles show a 101 400 and a 418 000 periodicity, respectively. These periodicities are within the tight independent constraints placed on the duration of these cycles by radiometric dates and sediment thickness. During the Triassic the Newark Basin was located at about 15 N, paleolatitude and the periodicities and the ratios of the short and compound cycles are in very close agreement with the Milankovitch predictions for that paleolatitude.

During the early Mesozoic, Pangea rested on the equator (Fig. 1) and a long series of rift valleys formed around the zone which was to become the Atlantic Ocean. These rift valleys filled with thousands of meters of sediments and igneous rocks and the North American contingent of what remains of these deposits is called the Newark Supergroup (1). The exposed portions of the Newark Supergroup consist largely of lacustrine deposits and these rocks preserve a record of changing climate spanning some 35 million years over a distance of over 2000 kilometers. The largest and thickest sequences are preserved in the Newark




Figure 1 Lockatong Formation in the Newark Basin : A, Predrift (Late-Triassic, Early Jurassic) position of rift basins (Newark Supergroup consists of basins exposed in eastern North America - continental positions from (6) - geology adapted from (1,6,7)) ; B, distribution of Lockatong Formation in Newark Basin - intrusions omitted (a, position of sections in Weehawken area (sections A-F in Figure 5); b, Delaware River section; c, Rushland I quarry section; d, Eureka Quarry section); C, major cities around Newark Basin (a, New York City; b, Trenton; c, Philadelphia).

Basin division of the Newark Supergroup in New York, New Jersey and Pennsylvania. Van Houten (2,3) was one of the first to recognize that much of the Newark Basin record is cyclic and that this cyclicality reflected periodically changing climate. He suggested that this climate change was in turn controlled by the precession of the equinoxes. This paper is essentially a reexamination, extension, and confirmation of Van Houten's original work.

There are 9 formations in the Newark Basin section (4,5) (Fig. 2) and of these the Late Triassic Lockatong and overlying Passaic formations comprise the longest continuous lacustrine record. The Lockatong (Figs 1 and 2) consists almost entirely of a spectrum of thin (ca. 5 m) sedimentary cycles, the end




Figure 2 Formations of the Newark Basin and their ages : A, radiometric scales for the Late Triassic (Late Carnian is here assumed to be 1/3 of Carnian time and the Mid-Late Carnian and Triassic-Jurassic boundaries are correlated with the basin section by biostratigraphic data); B, system and stage boundaries based on biostratigraphy, radiometric dates, and paleomagnetism; C, diagram of Newark Basin section (black horizontal lines and black bars represent gray and black units of the Triassic portion of the section and white areas and bars are principally red beds) ; D, formations of the Newark Basin (from 4,5).

members of which Van Houten (2,3) termed detrital and chemical (Fig. 3). The vertical sequence of fossils and sedimentary structures found in the cycles show that they owe their origin to the rise and fall of lakes. Each detrital cycle averages 5.7 m and can be divided into three lithologically distinct divisions (Fig. 3) (7) : division 1 consists of platy to massive gray mudstone sometimes showing current bedding deposited during lake transgression; division 2 is a fine calcareous black mudstone often platy to microlaminated and organic rich, deposited during lake high-stand; and division 3 is made up of platy to massive gray mudstone deposited during lake regression, low-stand, and lake bottom exposure. Chemical cycles are, on the average thinner than detrital cycles (4.5 m) and consist of

a lower platy gray mudstone and an upper much more massive gray or red mudstone often rich in sprays of analcime and dolomite crystals and pseudomorphs after gypsum, glauberite and halite - hence Van Houten's epithet chemical cycles.

Van Houten (2,3) reasoned that the vertical sequence of bed types with short cycles reflected the expansion and contraction of lakes. Detrital cycles were produced by lakes which were deep enough to have reached their outlet and were hence through-flowing. Chemical cycles, in contrast, were produced by lakes which never reached their outlet and therefore they concentrated salts during their entire existence. Van Houten argued that all these lakes were relatively shallow and that their depths were governed by changes in precipitation and thus the individual short cycles record regional cyclic changes in precipitation.




Figure 3 Major features of Locketong detrital cycles. This cycle is marked SH in Figure 9.

The alternative explanation to changing climate is that the short sedimentary cycles were produced by the filling in of small shallow lakes by sediments, the differences between the cycles being due to local hydrographic differences, and the development of the lake basins being due to local tectonics. Chemical cycles might then form geographically adjacent to




Figure 4 Detrital cycles W-5 (above) and W-6 (below) at excavation at Weehawken, New Jersey showing distribution of major fossils and their preservation : "deep tailed swimmer" and Tanytrachelos are small reptiles, and Diplurus, Turseodus, Semionotus, Synorichthys, and Cionichthys are fishes. This is locality A in Figure 5.

detrital cycles. Information on the lateral extent of cycles is needed to select between these two explanations.

Over the last six years I have been attempting to trace individual Lockatong cycles laterally. This is made difficult principally because of : 1) the scarcity of large, continuous exposures; 2) the presence of lateral facies changes; and 3) the general similarity of all the sedimentary cycles. Detailed investigations of single sections, however, have shown that each cycle can be uniquely defined by the details of the vertical changes in the kinds, preservation, and diversity of fossils (Figs 4 and 5). It has become evident that it is a property of detrital cycles that these vertical changes seen in single cycles remain consistent laterally (Fig. 5). This property allows for individual cycles to be identified in vertical sections of many cycles and allows individual cycles to be traced over large distances (Figs 5 and 6). I first worked out a cycle-by-cycle




Figure 5 Lateral extent of Lockatong detrital cycles from Weehawken to Fort Lee, New Jersey: A, detailed correlation by the vertical distribution of fossil fishes of cycles W-5 (above) and W-6 (below) [abbreviations of vertebrate fossils - D, "deep tailed swimmer", T, *Tanytrachelos*, Dp, *Diplurus*, Tu, *Turseodus*, Sy, *Synorichthys*, S, *Semionotus*, and C, *Cionichthys*] B, correlation of all exposed cycles laterally based on the vertical distribution of fossil fishes. Localities of sections A - F shown in Figure 1 and given in detail in (8).

stratigraphy for 18 cycles in the Hoboken to Fort Lee area (7) where outcrops are closely spaced (about 1 per 2 km for 20 km (Figs 5 and 6)). Recently, a large series of new outcrops have been identified in the central and southern Newark Basin which permit the cycles found in the Hoboken - Fort Lee area to be traced over most of the basin. In addition, these newly found exposures allow the same cycles originally worked on by Van Houten to be traced over large areas of the central Newark Basin (Fig.6). It is now apparent that many individual detrital cycles can be traced over most of the Newark Basin and that these cycles retain their particular faunal identities over these areas. It is also apparent that the short sedimentary cycles of the Lockatong reflect changes in the depths of lakes which covered much, if not all, of the Newark Basin and that changes in precipitation and evaporation rates are the most plausible cause of the lake-level change.




Figure 6 Correlation across Newark Basin of detrital cycles of Weehawken member of lower Lockatong (see Figure 7) showing distribution of most abundant fish. Abbreviations of fish taxa as follows : S, Semionotus, T, Turseodus, Sy, Synorichthys, D, Diplurus, and DI, large coelacanth. Positions of sections shown in Figure 1.

A consequence of the ability to trace individual cycles laterally is that lithological and paleontological features which are associated with various degrees of lateral continuity can be identified. Those cycles fitting Van Houten's detrital category prove to be traceable individually over much longer distances than those fitting the chemical category which are restricted to the central 60 km of the basin. Furthermore, those detrital cycles which contain a microlaminated division 2 are traceable over the largest distances with the least change. The microlaminated portions of these cycles contain the best preserved vertebrate remains. The very fine laminae and the fine preservation of fossil fish and other delicate organisms in the microlaminated portions of division 2 show that deposition of these units could only have occurred below wave base (7,9).

The lateral continuity of the microlaminated portions of division 2 allows for an estimation of the minimum extent and depth of the Lockatong lakes during their maximum transgression. I have developed a new method of estimating the depths of ancient lakes which uses the familiar relationship between the maximum fetch of a water body, wind speed and the depth of wave mixing (9). Using this method, the fact that the microlaminated (i.e. varved) fish-bearing portion of individual Lockatong cycles cover over 7000 km<sup>2</sup> means that the lake which deposited that unit must have covered more than 7000 km<sup>2</sup> and must have been more than 100 m deep. The fact that the portions of Lockatong cycles which were deposited during lake low stand can also be traced over the extent of the formation illustrates that the lakes actually did become very shallow and frequently dried out entirely. Thus, those detrital cycles with a microlaminated division 2 resulted from tremendous fluctuations in water depth, similar to or greater in magnitude than the changes known to have occurred in many of the lakes of the Great Basin in the western United States (10,11) and the Great Lakes of Africa (12,13,14).

Detrital and chemical cycles do not occur randomly in vertical section. Clusters of 1 to 4 detrital and 1 to 4 chemical cycles make up 25 m cycles (added up to a total of 4 to 6 short cycles) (Fig. 7) and clusters of these 25 m cycles alternately dominated by chemical or detrital cycles clusters make up 100 m cycles (Fig. 7) (2,3). The pattern of 100 m cycles allows the Lockatong formation to be broken up tentatively into 11 members (Fig. 7). This pattern of 100 m cycles continues into the overlying Passaic Formation. In the latter, however, the chemical cycles disappear and their position is taken by red beds (Figs 2 and 7).

The classification of sedimentary cycles types developed by Van Houten and followed in my own previous papers is






Figure 7 A, tentative division of the Lockatong Formation into 100 m members as exposed along the Delaware River in the Hunterdon Plateau Fault Block: Black horizontal bars represent portions of the Lockatong dominated by detrital cycles, white bars indicate portions of Lockatong made up mostly of red chemical cycles or red beds, gray bars indicate portions made up mostly of gray chemical cycles. Black vertical bars indicate portions of Lockatong shown in Figures 9 and 10. B, measured section of McLaughlin's (17) First Thin Red and the Skunk Hollow Member of this report. C, key for B.

essentially a typological one. The method of classification forces cycles to be identified either as chemical or as detrital. The operational definition of chemical cycles requires that evaporitic minerals, especially zeolites and dolomite, be present in the cycle. Thus, even if in all other respects a cycle without evaporitic minerals looks exactly other chemical cycles and is the lateral equivalent of a "true" chemical cycle it is classified as detrital because it lacks the essential aspect of a chemical cycle. This purely semantic problem produces confusion in interpretation when by definition, chemical cycles turn into detrital cycles laterally. In addition, in this sort of typological classification system all the information inherent in intermediate cycles is lost. This is the nature of typological classifications.

As suggested by Van Houten (2,3) and Olsen (7) various features of the sedimentary fabric of the rock types which make up Locketong short cycles have characteristics which can be interpreted in terms of water depth, at least in a relative sense. Thus, division 2 of detrital cycles is interpreted as having been deposited in deeper water than divisions 1 or 3 because of the absence of indications of exposure to the air, presence of undisturbed laminations, lack of roots, etc. The vertical sequences of sediment fabric types through a cycle shows directional change in several characters which reflect the expansion and contraction of the lakes in which they were deposited. I have developed a tentative classification of sedimentary fabrics within Locketong cycles with seven categories (Fig. 8) ranked from 0 to 6 in order of the relative depth of water in which they were deposited. This sequence is seen in vertical succession through typical detrital cycles as 1 2 3 4 5 4 3 2 1 0 (from the bottom up). Walther's law suggests that these same




Figure 8 Ranks of sedimentary fabrics of mudstones and siltstones of Locketong Formation arranged in order of the interpreted relative depth of water in which they were deposited and their relationship to organic carbon content and fossil fish preservation. Rank 0 (not shown) is a microbrecciated and disrupted massive mudstone sometimes with evaporitic minerals.

fabrics might be reflected in lateral changes from the center of the basin outwards, as well, and for the most part they are (7).

When the rank of the fabrics of beds which compose Lockatong short cycles are graphed against the measured section (Figs 9 and 10), the rank can be read as an indicator of water depth and hence a proxy indicator of (at least) local climate. The curve produced by this method is analogous to curves produced




Figure 9 Section of First Thin Red and Skunk Hollow members of Lockatong along Delaware River (see Figure 7) showing the classification of cycle type according to Van Houten (2,3), the application of depth ranks based on Figure 8, and the duration of short sedimentary cycles. Duration is equal to  $0.24 \text{ mm} \times \text{cycle thickness}$  and this value is placed at base of cycle. SH is cycle drawn in Figure 3.